

Christie, Mark, Miller, Paul K., Barry, Timothy and Dewhurst, Susan (2012) The case for community conservation in cardiovascular health promotion in the North West: Evidence from the Lancaster 'Greenfingers' initiative. *The Cumbria Partnership Journal of Research, Practice and Learning*, 2 (1). pp. 22-26.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/1415/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
 - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

The case for community conservation in cardiovascular health promotion in the North West: Evidence from the Lancaster 'Greenfingers' initiative

Mark Christie, Paul K. Miller, Tim Barry & Susan Dewhurst

Abstract

In this paper, a case is made for the greater promotion of regular cardiovascular exercise in the North West through the potential of a (hitherto under-utilised) resource that North Lancashire and Cumbria have in abundance: rural environments. Drawing upon extant literature and indicative data from pilot research from a recent community conservation initiative in Lancaster, it is argued that the embedded activities can encourage precisely the kinds of optimal, affordable and sustainable exercise that could play an important role in the region's cardiovascular health.

Keywords

CVD; cardiovascular disease; conservation; fitness; Green Gym; exercise referral; health promotion.

1. Introduction

According to the British Heart Foundation Health Promotion Research Group:

Diseases of the heart and circulatory system...are the main cause of death in the UK and account for almost 191 000 deaths each year – one in three of all deaths. (Scarborough et al., 2010, p. 14)

Moreover, between 2006 and 2008, cardiovascular disease (henceforth CVD) accounted for 28% of all premature deaths (i.e. deaths under the age of 75 years) in men and 20% women (Ibid.). Among the nine English NHS regions, the age-standardised death rates among the under-75 age groups in the North West (25.46 per 100 000) were higher than all other regions bar the North East (26.35 per 100 000) and well above the national average of 19.69 per 100 000 (Scarborough, et al., 2010, p. 29). The specific picture in the broad North Lancashire and Cumbria area is a little more mixed;

some Local Authority districts such as Copeland (28.12) Lancaster (24.63) Carlisle (21.21) show premature CVD death rates above the national average, while others such as South Lakeland (14.52) are well below (Ibid.). There is a profusion of compelling evidence that regular aerobic exercise, i.e. that which increases breathing and heart rate, is a powerful preventative against CVD (see, for example, the seminal meta-analysis compiled by Berlin & Colditz, 1990). A corollary of this is the UK Chief Medical Officer's recommendation that adults aged 19-64 years should exercise at a moderate (or higher) intensity for thirty minutes or more at least five times per week (Department of Health, 2004), recently rephrased as:

Adults should aim to be active daily. Over a week, activity should add up to at least 150 minutes (2½ hours) of moderate intensity activity in bouts of 10 minutes or more. (Department of Health, 2011, p. 1).

Sport England's rolling *Active People Survey* measures the exercise uptake situation in the North West of England as falling far short of these recommended guidelines; between October 2009 and October 2010, only 15.2% of Cumbrian respondents, and 11.6% of Lancastrians, reported having taken part in any 'sport or physically active recreation' on 20 or more of the last 28 days while 47.2% and 49.4%, respectively, reported having taken part in none whatsoever (Sport England, 2011)ⁱ.

2. Cardiovascular Exercise and the 'Green Gym' Concept

A number of initiatives have, to date, been trialled to increase public engagement with basic cardiovascular exercise at national and local levels, including primary care 'exercise referrals'. While showing clear short-term benefits, the long-term value of these referrals remains in dispute (Riddoch, Puig-Ribera, & Cooper, 1998), largely as an output of difficulties in sustaining participant commitment to exercise post-intervention (Morgan, 2005). Among the most recent initiatives to emerge from outside the NHS itself, meanwhile, is the 'Green Gym'. First piloted in 2000 by the British Trust for Conservation Volunteers (BTCV), a green gym is a scheme through which members of the public are invited to take part in physically-active conservation activities with a view to concurrently improving their own health and the public environment. The basic principle here is not new; a range of authors (Ferris, Norman, & Sempik, 2001; Woolley, 2004) have stressed the benefits that green spaces and community gardens can offer in terms of general health and wellbeing. The provision of access to activity in natural environments, it is argued, facilitates regular exercise and communal interaction, with tangible benefits in physical and psychosocial domains (Maas & Verheij, 2007). A recently-published independent evaluation of a range of BTCV Green Gym initiatives (Yerrell, 2008) has, however, *empirically* illustrated the fitness

benefits gained from working outdoors in this manner. Reported activities in which volunteers participated were varied, but findings suggested that they were largely of sufficient duration and intensity to improve general cardiovascular fitness amongst participants and significant improvements in physical health scores after six months of regular involvement are also reported. Crucially, 72% of original participants tended to remain regularly involved after this initial six month period, significantly outstripping sustained adherence figures for more orthodox 'exercise referral' models (typically in the 25-30% range, as reported by Taylor, Doust, & Webborn, 1998), largely due to the concomitant community-building aspect endemic in Green Gym initiatives. Furthermore, levels of physical activity among participants are also reported to have increased in their lives away from the projects themselves (Yerrell, 2008).

Although, to date, the corpus of evidence in terms of the direct and sustainable cardiovascular health benefits of Green Gym initiatives is still not fully robust, there is little doubt that the range of research completed suggests very encouraging outcomes. This is reflected in the fact that, by 2008, 10% of all participants in BTCV green gym initiatives were directly referred by local health services (Yerrell, 2008).

2.1. Case: The 'Greenfingers' Initiative in Lancaster

The first run of the 'Greenfingers' initiative, which began in October 2010, was promoted in partnership with Lancaster City Council as a community-based project involving the clearance of the non-native, invasive rhododendron from a section of Lancaster's Williamson Park, and ultimately its replacement with native plant species. Tasks involved a range of physical activities, including pruning, sawing, digging, chopping, clearing and general tidying of the designated area of the park, approximately an acre and a half in size. Following an initial nine week exploratory period, during which the majority

of volunteers were recruited, the project was extended for a further total of six months.

3. Research Design

The current position of the American College of Sports Medicine for development and maintenance of physical fitness and health (Garber et al., 2011) is that exercise training above and beyond activities of daily living should include cardiovascular component working at 'moderate intensity', i.e. a heart rate of 64-76% of aged predicted maximum heart rate (HR_{max} , which is defined as 220 minus the participant age in years), for a total of ≥ 150 min.wk⁻¹. There is substantial evidence to suggest that working at this exercise volume is associated with lower the rates of CVD in many populations (Lee, Rexrode, Cook, Manson, & Buring, 2001; Manson et al., 2002; Sesso et al., 2000). In the present pilot study, heart rate was recorded from a sample of two populations ('young' and 'older') participating in the Greenfingers initiative to monitor whether the workload was sufficient to induce physical fitness and health benefits. 7 young males aged between 19-28 (21.9 ± 3.3) years and 5 older individuals (3 males and 2 females) aged between 42 and 68 (59.0 ± 10.0) years volunteered for the study, all were healthy and Lancaster based. Young volunteers acted also as activity leadersⁱⁱ.

The physical activity included a variety of outdoor tasks which included sawing, clearing and pruning and lifting for a period of 55 minutes. Heart rate was recorded every five minutes using Polar Heart Rate Monitors. Resting heart rate was recorded at 0 min and 5 min, passive recovery was given before the final heart rate reading at 60 min. For both the absolute heart rate response (bts.min⁻¹) and HR_{max} (%) t unpaired Students T-Test were performed on the mean of the young and older data sets (SPSS, v.15).

4. Results

Figure 1: Heart Rate response (bts.min⁻¹) in 5 older subjects and 7 younger subjects. Values are mean \pm SE. * indicates significant difference ($p < 0.05$) between the mean of the groups.





Figure 2: Percentage of HR_{max} (maximum heart rate calculated from 220 -age) in 5 older subject and 7 younger subjects. Values are mean \pm SE. The grey band indicates the moderate intensity training zone suggested by Garber et al. (2011) of 64- 76% of HR_{max} .



Despite the absolute heart rate data being significantly higher in the older group, when this data is normalised using the percentage of HR_{max} , as shown in Figure 2, there is no difference between the intensity that the groups worked at (a mean of 66.4 ± 4.4 % HR_{max} compared to 71.3 ± 6.5 % HR_{max} , young and older group respectively. After 10 minutes, both groups were working at the exercise intensity defined by the ACSM (Garber, et al., 2011) as 'moderate' (between 64 and 76 % HR_{max} , shown by the grey band).

The young group, however, remained at the lower end of the banding and never increased above this moderate intensity while the older group, on occasion, were working at an intensity which would be classified as 'vigorous' (Ibid.).

5. Discussion

The main finding of the present Greenfingers pilot study was that the intensity of the work undertaken was sufficient to consistently raise heart rate into the moderate intensity zone, which has been shown to induce both physical fitness and health benefits in both the young and older groups. The ACSM (Garber, et al., 2011) suggest that physical exertion should be either of a moderate intensity, vigorous intensity or combination of moderate and vigorous intensity. The data from the Greenfingers initiative suggests that this form of community conservation activity produces the former in younger populations, and the latter in older participantsⁱⁱⁱ.

Although of small-scale, the pilot study findings clearly indicate that the activities involved in community conservation work can, providing that work schemes are appropriately tailored, produce optimal work for cardiovascular fitness. Anecdotal evidence collected from participants during the project also indicated increased motivation for engaging with regular physical activity outside of the scheme itself, while existing literature (Yerrell, 2008) suggests that these types of scheme may well produce much higher levels of sustained adherence than exercise referrals to leisure centres. Indeed, several participants in Greenfingers itself continued to be involved within Williamson Park as volunteers following the end of the scheme. As with BTCV Green Gyms, Greenfingers was inexpensive to establish and operate, run by volunteers and free to join (with, therefore, no associated cost for referrals). Locally, BTCV has now run projects in Preston and Carlisle, and a counterpart 'Fit to Grow' scheme, promoted by Carlisle City Council, is currently being

established in Currock, Cumbria. The Federation of City Farms and Community Gardens (2011) note that local authorities and health organisations often own land that is derelict and/or underused, which could well become the focus for active participation projects of this nature in the future. Health and leisure partnerships, linking with community groups, could potentially, therefore, unlock less conventional funding streams, including biodiversity, volunteering and community development grants.

In a region with substantial tracts of rural landscape, thus, community conservation projects would appear to provide an economical and effectual means of boosting public engagement with cardiovascular activity. More robust and extensive data, regarding a range of health indicators, will be collected to build a stronger evidence base when Greenfingers begins its second phase in October 2011. As part of an enhanced local body of knowledge, such research could prove vital in making the region's case for the funding of similar community projects (and related research) through new GP consortia and so forth, with substantial benefits for public health and environment alike.

Notes

ⁱ Some caution should be exercised in approaching these statistics. Given the self-reported nature of the data, it is fair to assert they may well exaggerate the levels of physical activity actually undertaken.

ⁱⁱ All subjects completed an informed consent form and ethical approval was granted by the University of Cumbria Ethics Committee.

ⁱⁱⁱ In addition to performing cardiovascular exercise, the ACSM (Garber, et al., 2011) also recommends that healthy adults should engage in physical activity that incorporates resistance exercises, and exercises to improve neuromotor control. Although not measured in the present study, the green gym activities included lifting (and in some instances heavy lifting), bending and cutting, all of which would task the neuromuscular system.

Affiliations

Mark Christie, Senior Lecturer in Sport and Physical Activity Development, Faculty of Health & Wellbeing, University of Cumbria.

Dr. Paul K. Miller, Senior Lecturer in Social Psychology, Faculty of Health & Wellbeing, University of Cumbria.

Tim Barry, Principle Lecturer in Physiology, Faculty of Health & Wellbeing, University of Cumbria.

Dr. Susan Dewhurst, Senior Lecturer in Physiology, Faculty of Health & Wellbeing, University of Cumbria.

Contact

Mark Christie, mark.christie@cumbria.ac.uk

References

- Berlin, J., & Colditz, G. (1990). A meta-analysis of physical activity in the prevention of coronary heart disease. *American Journal of Epidemiology*, *132*, 612 - 628.
- Department of Health. (2004). At least five a week: evidence on the impact of physical activity and its relationship to health: a report from the Chief Medical Officer. London: Department of Health.
- Department of Health. (2011). Factsheet 4: Physical activity guidelines for Adults (19–64 years). London: Department of Health.
- Federation of City Farms and Community Gardens. (2011). Gardening & Health. Retrieved 18/08, 2011, from <http://www.farmgarden.org.uk/home/local-food-project/growing-trends/gardening-a-health>
- Ferris, J., Norman, C., & Sempik, J. (2001). People, land and sustainability: Community gardens and the social dimension of sustainable development. *Social Policy and Administration*, *35*(5), 559-568.
- Garber, C. E., Blissmer, B., Deschenes, M. R., A., F. B., Lamonte, M. J., Lee, I. M. (2011). Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Medicine & Science in Sports & Exercise*, *43*(7), 1334-1359.
- Lee, I. M., Rexrode, K. M., Cook, N. R., Manson, J. E., & Buring, J. E. (2001). Physical activity and Coronary Heart Disease, in Women: is "no pain, no gain" passe? . *Journal of the American Medical Association*, *285*(11), 1447-1454.
- Maas, J., & Verheij, R. A. (2007). Are health benefits of physical activity in natural environments used in primary care by general practitioners in The Netherlands? *Urban Forestry & Urban Greening*, *6*(4), 227-233.
- Manson, J. E., Greenland, P., LaCroix, A. Z., Stefanick, M. L., Mouton, C. P., Oberman, A. (2002). Walking compared with vigorous exercise for the prevention of cardiovascular events in women. *New England Journal of Medicine*, *347*(10), 716-725.
- Morgan, O. (2005). Approaches to increase physical activity: reviewing the evidence for exercise-referral schemes. *Public Health*, *119*(5), 361-370
- Riddoch, C., Puig-Ribera, A., & Cooper, A. (1998). Effectiveness of physical activity promotion schemes in primary

care: a review. London: Health Education Authority.

[=a3e2e2c4-9e39-4392-9e74-6cd51e750694&version=5](#)

Scarborough, P., Bhatnagar, P., Wickramasinghe, K., Smolina, K., Mitchell, C., & Rayner, M. (2010). *Coronary heart disease statistics 2010*. London: British Heart Foundation.

Sesso, H. D., Stampfer, M. J., Rosner, B., Hennekens, C. H., Gaziano, J. M., Manson, J. E. (2000). Systolic and diastolic blood pressure, pulse pressure, and mean arterial pressure as predictors of cardiovascular risk in men. *Hypertension*, *36*, 801-807.

Sport England. (2011). Sport and active recreation (NI8), by frequency and key demographics. Retrieved 26/08, 2011, from http://www.sportengland.org/research/active_people_survey/idoc.ashx?docid

Taylor, A., Doust, J., & Webborn, N. (1998). Randomised controlled trial to examine the effects of a GP exercise referral scheme in Hailsham. East Sussex, on modifiable coronary heart disease risk factors. *Journal of Epidemiology and Community Health*, *52*, 595–601.

Woolley, H. (2004). *The value of public space: how high quality parks and public spaces create economic, social and environmental value*. London: University of Sheffield/CABE.

Yerrell, P. (2008). *National Evaluation of BTCV's Green Gym*. Oxford: School of Health and Social Care, Oxford Brookes University.