

Dewhurst, Susan, Nelson, Norah, Dougall, Paul K. and Bampouras, Theodoros (2014) Scottish country dance: benefits to functional ability in older women. *Journal of Aging & Physical Activity*, 22 (1). pp. 146-153.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/1539/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
 - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

Scottish Country Dance: Benefits to Functional Ability in Older Women

Susan Dewhurst, Norah Nelson, Paul K. Dougall, and Theodoros M. Bampouras

The effects of long-term participation in Scottish country dance on body composition, functional ability, and balance in healthy older females were examined. Participants were grouped into dancers and physically active nondancers (ages 60–70 and 70–80 for both groups). Physical activity, body composition (body-mass index, skinfold thickness, waist-to-hip ratio), functional ability (6-min walk distance, 6-m walk time, 8-ft up-and-go time, lower body flexibility, shoulder flexibility), and static balance were measured. Younger dancers and physically active nondancers had similar 6-min walk distance, 6-m walk time, and 8-ft up-and-go time results; however, while older dancers performed similarly to younger dancers, older physically active nondancers performed poorer than their younger counterparts ($p < .05$). Body composition and static balance were the same for all groups. Regular physical activity can maintain body composition and postural stability with advancing age; however, Scottish country dance can delay the effects of aging on locomotion-related functional abilities.

Keywords:

Regular physical activity is associated with numerous health benefits for individuals of all ages. Engaging in the recommended frequency, intensity, and duration of exercise proposed by the American College of Sports Medicine and the American Heart Association decreases all causes of mortality by reducing the risk of developing chronic disease (Garber et al., 2011). While the biological aging process cannot be prevented, regular physical activity can mitigate the potentially disabling effects of aging (American College of Sports Medicine et al., 2009). For older adults, the aim of exercise interventions is not only to reduce the risk of chronic disease, but also to maintain functional ability, allowing for independent aging.

The ability to perform everyday living tasks such as rising from a seated to a standing position, walking at a speed fast enough to successfully cross the road, stepping onto or avoiding a raised object, and simply maintaining an upright postural position are key indicators of functional ability and, consequently, independent living. Neuromuscular degeneration is a key contributor to a reduced ability to perform these everyday tasks with increasing age. Fortunately, the aging neuromuscular system allows for regeneration following a period of resistance training or neuromotor training (for detailed reviews, see Chou, Hwang, & Wu, 2012; Granacher, Zahner, & Gollhofer, 2008; Peterson, Rhea, Sen, & Gordon, 2010). However, surveys show that with aging, there is a greater preference for lower intensity,

predominantly cardiorespiratory physical activities such as walking, with <15% of 65- to 75-year-old adults participating in strength training reducing to <9% for adults over 75 years old (Schoenborn & Adams, 2010).

Scottish country dance is an activity that is hugely popular in older individuals—not only in Scotland, but also in North America, Australia, and New Zealand. The cultural resonance and social benefits that come with group dance make this an appealing activity for the older age group (Cooper & Thomas, 2002; Lima & Vieira, 2007; Wikstrom, 2004). Scottish country dance, like other traditional dance forms such as square dancing and folk dancing, articulates choreographed multidirectional movements incorporating many elements of neuromuscular control. It consists of a number of set dances in pairs and groups, with varying degrees of intensity. Participants perform various vertical and lateral movements, turning and spinning around in different directions to the rhythms of both slow and faster tempos.

In recent years, evidence is emerging on the physical benefits of various forms of dance to an older population (for detailed review, see Keogh, Kilding, Pidgeon, Ashley, & Gillis, 2009). Intervention studies have found that after 8–12 weeks participation in dance, there are increases in aerobic power, muscle endurance, muscle strength, and flexibility (Eyigor, Karapolat, Durmaz, Ibisoglu, & Cakir, 2009; Hopkins, Murrah, Hoeger, & Rhodes, 1990; Machado, Schettino, & Pereira, 2010; McKinley et al., 2008). Specific to functional ability, improved walking, change of direction, timed up-and-go, and stair climb speed have been reported, along with favorable improvements in both static and dynamic balance (Federici, Bellagamba, & Rocchi, 2005; Hackney, Kantorovich, & Earhart, 2007; Hopkins, Murrah, Hoeger, & Rhodes, 1990; Jeon, Choe, & Chae, 2000; Keogh,

Comment [A1]: Please provide 3 or 4 keywords/phrases for indexing purposes that are not included in the title.

Dewhurst and Bampouras are with the Faculty of Health and Wellbeing, University of Cumbria, Lancaster, UK. Nelson is with the School of Culture and Lifestyle, University of Derby, Buxton, UK. Dougall is with the Dept. of Sport, Culture and the Arts, University of Strathclyde, Glasgow, UK.

Kilding, Pidgeon, Ashley, & Gillis, 2012; McKinley et al., 2008; Shigematsu et al., 2002; Sofianidis, Hatzitaki, Douka, & Grouios, 2009; Song, June, Kim, & Jeon, 2004; Young, Weeks, & Beck, 2007). Similarly, cross-sectional studies found that older dancers had greater aerobic power, muscle strength, muscle endurance, muscle power, balance, and walking speed with additional findings of increased bone-mineral content compared with older nondancers (Uusi-Rasi et al., 1999; Verghese, 2006; Zhang, Ishikawa-Takata, Yamazaki, Morita, & Ohta, 2008).

This study assessed the functional ability of habituated Scottish country dancers, in comparison with individuals who participated in a variety of physical activities, to investigate whether Scottish country dance may improve the performance of various functional ability tasks.

Methods

Participants

Sixty healthy older females participated in the study after giving written informed consent. The sample consisted of Scottish country dancers ($n = 26$, age 67.6 ± 6 years, body mass 66.0 ± 9.3 kg, stature 160.2 ± 6.6 cm) and physically active nondancers ($n = 34$, age 71.3 ± 6 years, body mass 64.7 ± 10.6 kg, stature 157.1 ± 6.5 cm), which were split into two subgroups: a younger one (60–70 years) and an older one (70–80 years). Participants had no known neuromuscular disorders and had to be considered as medically stable, according to the criteria described by Greig et al. (1994). All participants were engaging in some form of physical activity at least three times a week. The dancers had a minimum of 10 years participatory involvement in Scottish country dancing. Physically active nondancers participated in a variety of activities, most commonly brisk walking, golf, bowls, and gardening. Ethical approval was obtained from the ethics committee of the University of Strathclyde, and all procedures followed were in accordance with the 1975 Declaration of Helsinki.

Physical Activity Assessment

Current levels of physical activity were measured using the Rapid Assessment of Physical Activity (RAPA) scale, which is an easy-to-use, valid measure of physical activity in older adults (Topolski et al., 2006). This scale consists of nine tick-box responses relating to level and intensity of usual physical activity, as well as additional strength and flexibility activities. RAPA 1 measures cardiorespiratory-based physical activity (scoring 1–7 points, with a maximum score of 7), RAPA 2a and 2b measure strength- and flexibility-based physical activity, respectively (yes/no answer).

Body-Composition Assessment

Body-mass index (BMI) was calculated from height and body mass. Skinfold measurements were taken in

triplicate using Harpenden skinfold calipers (Burgess Hill, UK) at four sites: biceps, triceps, subscapular, and suprailiac. The average value from each site was used, and the sum of the four sites (Sum4SF) is reported.

Waist and hip circumferences were measured in duplicate with an anthropometric tape while the subjects were wearing light clothing. Waist circumference was measured at the minimum circumference between the iliac crest and the rib cage. Hip circumference was measured at the maximum protuberance of the buttocks. Waist-to-hip ratio (waist:hip) was also calculated.

Functional Ability Assessment

A range of functional ability tests, as described by Rikli and Jones (2012) and Thomas, De Vito, and Macaluso (2007), were used to assess a number of parameters important in everyday living tasks.

6-Min Walk Test.

To estimate cardiorespiratory fitness, participants were instructed to walk at their normal walking pace for 6 min around a predetermined course in an indoor sports hall. Total distance covered was recorded. This test has been shown highly reliable for this population, with an interclass correlation of .91 (Rikli & Jones, 2012).

6-m Walk Time.

To assess maximal walking speed, participants started from a static standing position and walked as fast as they could to the end of a 9-m course (Thomas, De Vito, & Macaluso, 2007). Visible markers were placed at the start, 6 m, and 9 m. Time taken from start to 6 m was recorded using a stopwatch (Seiko, Tokyo, Japan). The test was done in triplicate, with the best score used for further analysis. Although reliability data are not available for this particular test, gait speed tests are highly reliable (Steffen, Hacker & Mollinger, 2002) with a similar distance gait speed test (7.62 m) yielding a correlation coefficient of $>.90$ (Bohannon, 1997).

8-ft Up-and-Go Time.

To assess the integration of neuromuscular parameters (power, speed, agility, and dynamic balance; Rikli & Jones, 2012), time taken to raise from a seated position, walk 2.44 m (8 ft), turn, and return to the seated position was recorded. This test was done in triplicate with the best score used for further analysis. The reliability of this test in a similar population was high (intraclass correlation coefficient = .9; Rikli & Jones, 2012).

Chair Sit-and-Reach.

To assess back and hamstring flexibility, while sitting on a chair with the legs stretched out in front, the volunteers were asked to reach down toward their toes. The distance between the extended fingers and the tip of the toes was measured. This was measured in triplicate for each side, and the average for each side was used for further analysis. Similar to the previous tests, this test was shown to have high reliability, as indicated by a high intraclass correlation coefficient of .96 (Rikli & Jones, 2012).

Comment [A2]: Please provide model of stopwatch. Also, is Seiko S-Yard Co Ltd the correct manufacturer, as listed here? That appears to be a golf club division rather than the stopwatch manufacturing division (Seiko Watch Corporation, Tokyo, Japan).

Discussion

The main finding of the current study was that dancers aged 60–70 years old had a similar functional ability (6-min walk distance, 6-m walk time, and 8-ft up-and-go time) as physically active nondancers of similar age. However, while older dancers (70–80 years) did not differ in performance to younger dancers, older nondancers showed a poorer performance when compared with their younger counterparts. Therefore, Scottish country dancing can serve as a beneficial physical activity in adequately stressing functional abilities of the older individuals. The current results concur with recent findings by Keogh et al. (2012) and Machado et al. (2010), who reported improvements in functional abilities and explosive strength of older individuals who participated in dancing.

All groups participated in the same amount of physical activity per week. The median response to the cardiorespiratory physical activity score was 6 on the RAPA scale of 1–7, worded as “I do 30 minutes or more a day of moderate physical activities, 5 or more days a week.” This meets the recommendation of the American College of Sports Medicine et al. (2009) for cardiorespiratory exercise in healthy older adults, quantified as being moderate-intensity exercise for >150 min/week or vigorous-intensity exercise >75 min/week. Strength-based physical activity had a much lower participation, as only 17% of the women reported doing activities to increase muscle strength once a week or more, whereas 54% responded positively to participating in activities to improve flexibility once a week or more. The strength results compare favorably with findings by Schoenborn & Adams (2010), who reported a similarly low percentage of participation in strength training (<15%) for older adults age 65–75 years. As there was no difference between the groups in the physical activity levels, it may be assumed that the differences in the functional ability tasks are due to the inclusion of Scottish country dance to the weekly physical activity of the young and older dancers.

Body-composition measures (BMI, Sum4SF, waist and hip circumference, waist:hip) were not different in dancers and physically active nondancers or between the younger and older age groups. When the BMI and waist circumference were categorized according to the National Institutes of Health (1998) guidelines, all groups fell into the lower end of the overweight category (BMI 25.0–29.9), with all groups having a waist circumference below the high disease-risk cutoff of ≥ 88 cm. Waist:hip ratio gives an index of subcutaneous and intra-abdominal adipose tissue. All groups fell below the abdominal obesity cutoff of ≥ 0.85 presented by the World Health Organization (2011). The findings agree with literature suggesting that meeting the physical activity guidelines has favorable effect on body composition, which in turn reduces the risk of developing chronic disease (for review, see American College of Sports Medicine et al., 2009).

The 6-min walk distance is a well-established measure of cardiorespiratory fitness, replicating efforts

commonly performed by healthy individuals during daily activities (Mangan & Judge, 1994). All groups were meeting the physical activity guidelines for cardiorespiratory fitness which is reflected with favorable comparisons to results reported from other community dwelling older populations (Bautmans, Lambert, & Mets, 2004; Lusardi, Pellecchia, & Schulman, 2003; Steffen et al., 2002). Of particular interest is that the 6-min walk distance, and therefore cardiorespiratory fitness, did not decrease in the older age group in the dancers, but it did in the physically active nondancers, suggesting that Scottish country dance may provide enough of a stimulus to delay the inevitable effects of aging on the cardiorespiratory system in individuals in their eighth decade of life. Indeed, the ES is very high for all comparisons, indicating a substantial difference rather than just a statistical one (Cumming, 2011). In addition, the younger dancing and nondancing groups, as well as the older dancing groups, were above the norms (by 5%, 4%, and 9%, respectively) provided by Rikli & Jones (1999), while the older control group was below (by 3%) those norms for this test.

Neuromuscular function is reflected in the 6-m walking time test and the 8-ft up-and-go. The 6-m walking time test has been found to improve with increased strength (Schlicht, Camaione, & Owen, 2001) while the 8-ft up-and-go test provides the neuromuscular system with a variety of challenges, including generation of leg force to lift the individual off the chair without using their arms, assume a balanced upright position, walk at speed, change direction, and walk at speed again while turning to resume a seated position (Rikli & Jones, 2012). The performance of these tasks is compromised with age, as the individuals may be working close to their maximum capacity. For example, simply rising from the chair has been estimated to require almost 100% of a healthy older individual's leg torque (Alexander, Schultz, Ashton-Miller, Gross, & Giordani, 1997). Our results for the 6-m walk test indicate slower speeds as compared with the speeds reported by Thomas et al. (2007). On the contrary, results for 8-ft up-and-go compare very favorably with norms reported by Rikli and Jones (1999), with all four groups scoring faster 8-ft up-and-go times than the norm. While for the physically active nondancers, both the 6-m walking speed and 8-ft up-and-go test performances was poorer in the older age group compared with the younger group (with large to high ES [Cumming, 2011], indicating practically significantly lower scores, too), this was not the case for the dancers. The younger and older dancers had similar performance for both assessments, suggesting that in addition to being of benefit to cardiorespiratory health, Scottish country dance may reduce the age-induced degeneration of the neuromuscular system. It is of interest to note that older dancers performed significantly better than the older physically active nondancers in the locomotion-related functional tests.

It is suggested that this reflects specific adaptations to the activity in question. Scottish country dance routines continually stresses the neuromuscular system with their skipping, turning, and hopping elements, and while no

direct comparison can be made to resistance training, it appears that Scottish country dance may be an effective way to train the neuromuscular system without the need of additional specific strength-based training.

Back and hamstring flexibility was measured from the chair sit-and-reach test, which was not different between the groups, suggesting that Scottish country dancing has no additional benefits to lower back or hamstring flexibility. This flexibility appears to have been maintained from the younger to the older groups, possibly due to over half of the participants engaging in flexibility-based exercises at least once a week. Interestingly, although the scores for both younger groups are higher than the norm (Rikli & Jones, 1999), the dancers' scores are below the norm for the older group, while just over the norm for the physically active control.

Upper body flexibility was estimated from the back scratch. The left-hand back scratch test showed a decrease in performance in the older physically active nondancers when compared with the younger physically active nondancers. This was not seen between younger and older dancers. Interestingly, the right back scratch did not show this with all groups having similar performances. As by far most participants were right handed, it is assumed that flexibility was better maintained in the dominant side, as the preferred side for many everyday functions (e.g., combing hair, putting on clothes, etc.; Rikli & Jones, 2012).

It is well established that aging affects balance in older individuals (for review see Maki & McLroy, 1996). However, in studies where physically active older participants were compared with non-physically active counterparts, physical activity was shown to have a positive effect on balance (Howe, Rochester, Neil, Skelton, & Ballinger, 2011; Perrin, Gauchard, Perrot, & Jeandel, 1999). Indeed, better balance was found with increased length of engagement in physical activity (Perrin, Gauchard, Perrot, & Jeandel, 1999). Balance in older individuals is an important factor for their well-being, as improved balance reduces the risk of falling (Choy, Brauer, & Nitz, 2003; Dewhurst et al., 2007; Howe, Rochester, Neil, Skelton, & Ballinger, 2011). The present findings concur with the literature, as balance was not different between the younger and older age groups for both Scottish country dancers and physically active nondancers, suggesting that physical activity levels were sufficient in both groups to ensure maintenance of postural control. A recent Cochrane review (Howe, Rochester, Neil, Skelton, & Ballinger, 2011) called for more research on activities that can improve balance in older populations. The present study posits that it may be possible to maintain balance in older adults not only through general physical activity, but also through more specific activities.

In summary, these findings advocate the participation in Scottish country dance in combating the inevitable degenerative effects of aging in the performance of functional ability tasks when compared with age-matched individuals participating in other forms

of cardiorespiratory physical activity. Our results agree with recommendations made by Keogh et al. (2009) regarding the benefits of dance in older women. Both dance and regular physical activity appear to maintain favorable body composition and prevent a decrease in balance with older age.

References

- Alexander, N.B., Schultz, A.B., Ashton-Miller, J.A., Gross, M.M., & Giordani, B. (1997). Muscle strength and rising from a chair in older adults. *Muscle & Nerve*, 5(Supplement), S56-S59. [PubMed doi:10.1002/\(SICI\)1097-4598\(1997\)5+<56::AID-MUS14>3.0.CO;2-X](#)
- American College of Sports Medicine, Chodzko-Zajko, W.J., Proctor, D.N., Fiatarone Singh, M.A., Minson, C.T., Nigg, C.R., . . . Skinner, J.S. (2009). American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Medicine and Science in Sports and Exercise*, 41(7), 1510-1530. [PubMed doi:10.1249/MSS.0b013e3181a0c95c](#)
- Bautmans, I., Lambert, M., & Mets, T. (2004). The six-minute walk test in community dwelling elderly: Influence of health status. *BMC Geriatrics*, 4, 6. [PubMed doi:10.1186/1471-2318-4-6](#)
- Bohannon, R.W. (1997). Comfortable and maximum walking speed of adults aged 20-79 years: reference values and determinants. *Age and Ageing*, 26, 15-19. [PubMed doi:10.1093/ageing/26.1.15](#)
- Chou, C.H., Hwang, C.L., & Wu, Y.T. (2012). Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: A meta-analysis. *Archives of Physical Medicine and Rehabilitation*, 93(2), 237-244. [PubMed doi:10.1016/j.apmr.2011.08.042](#)
- Choy, N.L., Brauer, S., & Nitz, J. (2003). Changes in postural stability in women aged 20 to 80 years. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 58(6), 525-530. [PubMed doi:10.1093/gerona/58.6.M525](#)
- Cooper, L., & Thomas, H. (2002). Growing old gracefully—Social dance in the third age. *Ageing and Society*, 22(6), 689-708. [doi:10.1017/S0144686X02008929](#)
- Cumming, G. (2011). *Understanding new statistics: Effect sizes, confidence intervals, and meta-analysis*. London, UK: Routledge.
- Dewhurst, S., Riches, P.E., & De Vito, G. (2007). Moderate alterations in lower limbs muscle temperature do not affect postural stability during quiet standing in both young and older women. *Journal of Electromyography and Kinesiology*, 17(3), 292-298. [PubMed doi:10.1016/j.jelekin.2006.03.002](#)
- Eyigor, S., Karapolat, H., Durmaz, B., Ibisoglu, U., & Cakir, S. (2009). A randomized controlled trial of Turkish folklore dance on the physical performance, balance, depression and quality of life in older women. *Archives of Gerontology and Geriatrics*,

- 48(1), 84–88. [PubMed](#)
[doi:10.1016/j.archger.2007.10.008](https://doi.org/10.1016/j.archger.2007.10.008)
- Federici, A., Bellagamba, S., & Rocchi, M.B. (2005). Does dance-based training improve balance in adult and young old subjects? A pilot randomized controlled trial. *Aging Clinical and Experimental Research*, 17(5), 385–389. [PubMed](#)
[doi:10.1007/BF03324627](https://doi.org/10.1007/BF03324627)
- Garber, C.E., Blissmer, B., Deschenes, M.R., Franklin, B.A., Lamonte, M.J., Lee, I.M., . . . American College of Sports Medicine. (2011). American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Medicine and Science in Sports and Exercise*, 43(7), 1334–1359. [PubMed](#)
[doi:10.1249/MSS.0b013e318213fefb](https://doi.org/10.1249/MSS.0b013e318213fefb)
- Granacher, U., Zahner, L., & Gollhofer, A. (2008). Strength, power, and postural control in seniors: Considerations for functional adaptations and for fall prevention. *European Journal of Sport Science*, 8(6), 325–340. [PubMed](#)
[doi:10.1080/17461390802478066](https://doi.org/10.1080/17461390802478066)
- Greig, C.A., Young, A., Skelton, D.A., Pippet, E., Butler, F.M., & Mahmud, S.M. (1994). Exercise studies with elderly volunteers. *Age and Ageing*, 23(3), 185–189. [PubMed](#)
[doi:10.1093/ageing/23.3.185](https://doi.org/10.1093/ageing/23.3.185)
- Hackney, M.E., Kantorovich, S., & Earhart, G.M. (2007). A study on the effects of argentine tango as a form of partnered dance for those with Parkinson disease and the healthy elderly. *American Journal of Dance Therapy*, 29(2), 109–127. [PubMed](#)
[doi:10.1007/s10465-007-9039-2](https://doi.org/10.1007/s10465-007-9039-2)
- Hopkins, D.R., Murrain, B., Hoeger, W.W., & Rhodes, R.C. (1990). Effect of low-impact aerobic dance on the functional fitness of elderly women. *The Gerontologist*, 30(2), 189–192. [PubMed](#)
[doi:10.1093/geront/30.2.189](https://doi.org/10.1093/geront/30.2.189)
- Howe, T.E., Rochester, L., Neil, F., Skelton, D.A., & Ballinger, C. (2011). Exercise for improving balance in older people. *Cochrane Database of Systematic Reviews*, (11), CD004963. [PubMed](#)
[doi:10.1002/14651858.CD004963.pub3](https://doi.org/10.1002/14651858.CD004963.pub3)
- Jeon, M.Y., Choe, M.A., & Chae, Y.R. (2000). Effect of Korean traditional dance movement training on balance, gait and leg strength in home bound elderly women. *Journal of Korean Academy of Nursing*, 30(3), 647–658.
- Keogh, J.W.L., Kilding, A., Pidgeon, P., Ashley, L., & Gillis, D. (2009). Physical benefits of dancing for healthy older adults: A review. *Journal of Aging and Physical Activity*, 17(4), 479–500. [PubMed](#)
- Keogh, J.W.L., Kilding, A., Pidgeon, P., Ashley, L., & Gillis, D. (2012). Effects of different weekly frequencies of dance on older adults' functional performance and physical activity patterns. *European Journal of Sports and Exercise Science*, 1(1), 14–23.
- Lima, M.M.S., & Vieira, A.P. (2007). Ballroom dance as therapy for the elderly in Brazil. *American Journal of Dance Therapy*, 29(2), 129–142. [PubMed](#)
[doi:10.1007/s10465-007-9040-9](https://doi.org/10.1007/s10465-007-9040-9)
- Lusardi, M.M., Pellecchia, G.L., & Schulman, M.K. (2003). Functional performance in community living older adults. *Journal of Geriatric Physical Therapy*, 26(3), 14–22. [PubMed](#)
[doi:10.1519/00139143-200312000-00003](https://doi.org/10.1519/00139143-200312000-00003)
- Machado, M., Schettino, L., & Pereira, R. (2010). Development of muscular explosive force in older women: Influence of a dance-based exercise routine. *Medicina Sportiva*, 4, 209–213.
- Maki, B.E., & McIlroy, W.E. (1996). Postural control in the older adult. *Clinics in Geriatric Medicine*, 12(4), 635–658. [PubMed](#)
- Mangan, D., & Judge, J. (1994). Reliability and validation of the six-minutes walk. *Journal of the American Geriatrics Society*, 42, SA73.
- McKinley, P., Jacobson, A., Leroux, A., Bednarczyk, V., Rossignol, M., & Fung, J. (2008). Effect of a community-based argentine tango dance program on functional balance and confidence in older adults. *Journal of Aging and Physical Activity*, 16(4), 435–453. [PubMed](#)
- National Institutes of Health. (1998). Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults—The evidence report. *Obesity Research*, 6(Suppl. 2), 51S–209S. [PubMed](#)
[doi:10.1002/j.1550-8528.1998.tb00690.x](https://doi.org/10.1002/j.1550-8528.1998.tb00690.x)
- Perrin, P.P., Gauchard, G.C., Perrot, C., & Jeandel, C. (1999). Effects of physical and sporting activities on balance control in elderly people. *British Journal of Sports Medicine*, 33(2), 121–126. [PubMed](#)
[doi:10.1136/bjism.33.2.121](https://doi.org/10.1136/bjism.33.2.121)
- Peterson, M.D., Rhea, M.R., Sen, A., & Gordon, P.M. (2010). Resistance exercise for muscular strength in older adults: A meta-analysis. *Ageing Research Reviews*, 9(3), 226–237. [PubMed](#)
[doi:10.1016/j.arr.2010.03.004](https://doi.org/10.1016/j.arr.2010.03.004)
- Rikli, R.E., & Jones, C.J. (2012). Development and validation of criterion-referenced clinically relevant fitness standards for maintaining physical independence in later years. *The Gerontologist*, 53(2), 255–267. [PubMed](#)
[doi:10.1093/geront/gns071](https://doi.org/10.1093/geront/gns071)
- Rikli, R.E., & Jones, C.J. (1999). Functional fitness normative scores for community-residing older adults, ages 60–94. *Journal of Aging and Physical Activity*, 7, 162–181. [PubMed](#)
[doi:10.1093/geront/gns071](https://doi.org/10.1093/geront/gns071)
- Schlicht, J., Camaione, D.N., & Owen, S.V. (2001). Effect of intense strength training on standing balance, walking speed, and sit-to-stand performance in older adults. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 56(5), M281–M286. [PubMed](#)
[doi:10.1093/gerona/56.5.M281](https://doi.org/10.1093/gerona/56.5.M281)
- Schoenborn, C.A., & Adams, P.E. (2010). Health behaviors of adults: United States, 2005–2007. *Vital and Health*

- Statistics. Series 10, Data from the National Health Survey, 245*, 1–132. [PubMed](#)
- Shigematsu, R., Chang, M., Yabushita, N., Sakai, T., Nakagaichi, M., Nho, H., & Tanaka, K. (2002). Dance-based aerobic exercise may improve indices of falling risk in older women. *Age and Ageing, 31*(4), 261–266. [PubMed doi:10.1093/ageing/31.4.261](#)
- Sofianidis, G., Hatzitaki, V., Douka, S., & Grouios, G. (2009). Effect of a 10-week traditional dance program on static and dynamic balance control in elderly adults. *Journal of Aging and Physical Activity, 17*(2), 167–180. [PubMed](#)
- Song, R., June, K.J., Kim, C.G., & Jeon, M.Y. (2004). Comparisons of motivation, health behaviors, and functional status among elders in residential homes in Korea. *Public Health Nursing (Boston, Mass.), 21*(4), 361–371. [PubMed](#) [doi:10.1111/j.0737-1209.2004.21410.x](#)
- Steffen, T.M., Hacker, T.A., & Mollinger, L. (2002). Age- and gender-related test performance in community-dwelling elderly people: Six-minute walk test, Berg balance scale, timed up & go test, and gait speeds. *Physical Therapy, 82*(2), 128–137. [PubMed](#)
- Thomas, E.E., De Vito, G., & Macaluso, A. (2007). Speed training with body weight unloading improves walking energy cost and maximal speed in 75- to 85-year-old healthy women. *Journal of Applied Physiology, 103*(5), 1598–1603. [PubMed](#) [doi:10.1152/jappphysiol.00399.2007](#)
- Topolski, T.D., LoGerfo, J., Patrick, D.L., Williams, B., Walwick, J., & Patrick, M.B. (2006). The rapid assessment of physical activity (RAPA) among older adults. *Preventing Chronic Disease, 3*(4), A118. [PubMed](#)
- Uusi-Rasi, K., Sievanen, H., Vuori, I., Heinonen, A., Kannus, P., Pasanen, M., . . . Oja, P. (1999). Long-term recreational gymnastics, estrogen use, and selected risk factors for osteoporotic fractures. *Journal of Bone and Mineral Research, 14*(7), 1231–1238. [PubMed doi:10.1359/jbmr.1999.14.7.1231](#)
- Vergheze, J. (2006). Cognitive and mobility profile of older social dancers. *Journal of the American Geriatrics Society, 54*(8), 1241–1244. [PubMed](#) [doi:10.1111/j.1532-5415.2006.00808.x](#)
- Wikström, B.M. (2004). Older adults and the arts: The importance of aesthetic forms of expression in later life. *Journal of Gerontological Nursing, 30*(9), 30–36. [PubMed](#)
- World Health Organization. (2011). Waist circumference and waist-hip ratio: Report of a WHO expert consultation, Geneva, 8–11 December 2008. Geneva, Switzerland: World Health Organization.
- Young, C.M., Weeks, B.K., & Beck, B.R. (2007). Simple, novel physical activity maintains proximal femur bone mineral density, and improves muscle strength and balance in sedentary, postmenopausal Caucasian women. *Osteoporosis International, 18*(10), 1379–1387. [PubMed](#) [doi:10.1007/s00198-007-0400-6](#)
- Zhang, J.G., Ishikawa-Takata, K., Yamazaki, H., Morita, T., & Ohta, T. (2008). Postural stability and physical performance in social dancers. *Gait & Posture, 27*(4), 697–701. [PubMed](#) [doi:10.1016/j.gaitpost.2007.09.004](#)




Figure 1 — Functional ability parameters: (A) 6-min walk distance, (B) 6-meter walk time, (C) 8-ft up-and-go time, (D) chair sit-and-reach distance, (E) left back-scratch distance, and (F) right back scratch distance for groups age 60–70 and 70–80 years. For (E) and (F), positive scores indicate hand overlap, while negative scores indicate the hands are apart. Scottish country dancers are shown with the white bars, and physically active control are shown with the black bars. *Significant difference between groups ($p < .05$). Data are presented as $M \pm SD$.




Figure 2 — Static balance area (95% ellipse area) for the groups age 60–70 and 70–80 years old. Scottish country dancers are shown with the white bars, and physically active control are shown with the black bars. Data are presented as $M \pm SD$.

Table 1 Anthropometric and Body-Composition Measurements for All Groups, $M \pm SD$

Category	Scottish Country Dancers		Nondancers	
	60–70 years (<i>n</i> = 15)	70–80 years (<i>n</i> = 11)	60–70 years (<i>n</i> = 13)	70–80 years (<i>n</i> = 21)
Age (years)	63.5 ± 3.0	73.7 ± 2.7	65.2 ± 3.4	75.1 ± 2.5
Height (cm)	1.61 ± 0.06	1.58 ± 0.08	1.55 ± 0.07	1.58 ± 0.06
Body mass (kg)	68.6 ± 9.5	62.4 ± 8.6	62.7 ± 11.8	65.9 ± 9.8
Body-mass index (kg/m ²)	26.5 ± 3.6	24.9 ± 2.8	25.9 ± 4.5	26.5 ± 4.6
Sum of four skinfolds (mm)	71.4 ± 18.9	67.5 ± 22.9	61.9 ± 19.1	72.1 ± 19.2
Waist circumference (cm)	84.0 ± 7.4	79.7 ± 8.2	80.5 ± 11.3	85.6 ± 9.4
Hip circumference (cm)	104.4 ± 6.1	101.0 ± 5.9	100.6 ± 9.5	104.4 ± 9.0
Waist:hip ratio	0.80 ± 0.04	0.79 ± 0.05	0.80 ± 0.06	0.82 ± 0.05