

Bampouras, Theodoros, Marrin, Kelly, Sankey, Sean P. and Jones, Paul A. (2014) Test-retest reliability and sensitivity of the Concept2 Dyno dynamometer: practical applications. *Journal of Strength and Conditioning Research*, 28 (5). pp. 1381-1385.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/1647/>

*Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.*

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

**provided that**

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
- a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

**You may not**

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing [insight@cumbria.ac.uk](mailto:insight@cumbria.ac.uk).

**Manuscript number: R- 347412**

**Revision 1**

**TITLE PAGE**

**Title:**

Test-retest reliability and sensitivity of the Concept2 Dyno<sup>®</sup> dynamometer: practical applications

**Brief running Head:**

Test-retest reliability of Concept2 Dyno

**Authors:**

Theodoros M. Bampouras, MSc<sup>1</sup>, Kelly Marrin, MSc, PhD<sup>2</sup>, Sean P. Sankey, MSc, CSCS<sup>3</sup>, and Paul A. Jones, MSc, CSCS<sup>4</sup>

**Departments:**

<sup>1</sup> Faculty of Health and Wellbeing, Sport and Physical Activity

<sup>2</sup> Department of Sport and Physical Activity

<sup>3</sup> Faculty of Advanced Engineering and Sciences

<sup>4</sup> Centre for Rehabilitation and Human Performance Research

**Institutions**

<sup>1</sup> University of Cumbria

<sup>2</sup> Edge Hill University

<sup>3</sup> University of Bolton

<sup>4</sup> University of Salford

**Corresponding author:** Theodoros M. Bampouras, Faculty of Health and Wellbeing, Sport and Physical Activity, University of Cumbria, Bowerham Road, Lancaster, LA1 3JD, United Kingdom / +44 1524 526531 / theodoros.bampouras@cumbria.ac.uk

**BLIND TITLE PAGE**

**Title:**

Test-retest reliability and sensitivity of the Concept2 Dyno<sup>®</sup> dynamometer: practical applications.

## **ABSTRACT**

Strength assessment is often part of the objective periodical observation of teams, squads or large groups of athletes. Equipment that provides assessment that is mobile and is easy to use will reduce the impact on the athletes training and competitive calendar. However, any equipment used must be reliable to allow accurate monitoring of performance. The aim of this study was to examine the reliability of the Concept2 Dyno<sup>®</sup> dynamometer. Forty-six competitive athletes (males: n=36, age 23.3±6.8 years, height 1.80±0.09 m, body mass 82.3±15.6 kg; females, n=10, age 20.7±1.4 years, height 1.65±0.09 m, body mass 62.7±11.8 kg), with a strength training background of more than 2 years, performed a familiarisation session and three experimental sessions with one week intervening each. Each experimental session consisted of three maximal efforts of seated chest press (CPress), seated row (SRow), and seated leg press (LPRESS) exercises. Reliability was assessed examining systematic bias, intraclass correlation coefficient (ICC), coefficient of variation (CV) and 95% limits of agreement (95%LoA) between sessions. No systematic bias was found for any of the exercises. ICC were high (0.89–0.98) with relatively low CV (6.2%–4.3%). Finally, 95%LoA indicated that subsequent testing could underestimate by a factor of 0.87 or overestimate by a factor of 1.17, on average. These results indicate that Concept2 Dyno<sup>®</sup> dynamometer is reliable and can be used in the field to efficiently monitor strength performance. Coaches and researchers should use ‘analytical goals’ to help decide as to the use of Concept2 Dyno<sup>®</sup> for their purposes.

**Keywords:** portable dynamometer, strength testing, sport-specific testing, test-retest,

## INTRODUCTION

Strength testing is a commonly utilised procedure to monitor adaptations to training interventions (7,9,13,18) or to provide an indication of any muscular weaknesses (7,11). Various methods have been used to assess strength, including isokinetic dynamometers (e.g. 11), force platforms (e.g. 18) or weights (e.g. 9). Technological developments have resulted in portable dynamometers, enabling strength testing to be conducted in the field, allowing higher test efficiency and functionality (3,15,21).

One portable dynamometer that allows assessment of three common multi-joint exercises (chest press, seated row and leg press) is the Concept2 Dyno<sup>®</sup>. The Concept2 Dyno<sup>®</sup> consists of an air-resisted flywheel, which responds to the user's efforts. The resistance can be manipulated by eight damper levers which control the air flow and increase the air resistance with more dampers open. The user adapts a seated position for all three exercises with their back (for the leg and chest press) and chest (for the seated row) supported. These positions mimic the position an individual would adapt to perform the abovementioned exercises in the respective exercise equipment.

The portability of the equipment, the familiarity of the design as well as the familiarity of the three exercises for people that do some resistance exercise, makes Concept2 Dyno<sup>®</sup> an appealing solution for strength assessment in the athletes own space. Indeed, this commercially available dynamometer is widely used by athletic clubs and the police force as well as by researchers for strength assessment (8,11).

However, in order for any strength testing to provide accurate and useful information, any assessment tool must demonstrate valid and reliable measures, additionally ensuring that changes in performance will be able to be detected (10). A number of studies have examined the reliability of portable strength testing devices comprising portable force platform (21) and wireless accelerometer (3), in order to allow for more sport-specific and efficient testing that would yield more informative results for athletes and coaches (15).

Notwithstanding the wide application of Concept2 Dyno<sup>®</sup>, such data does not exist for it. Although the exercises offered by Concept2 Dyno<sup>®</sup> have logical validity (as they have close resemblance with established respective exercises), the reliability and sensitivity of these exercises on Concept2 Dyno<sup>®</sup> should be examined to offer practical recommendations for strength assessment. Therefore, the aim of the present paper was to examine the reliability and sensitivity of the Concept2 Dyno<sup>®</sup>.

## **METHODS**

### **Experimental approach to the problem**

The reliability and sensitivity of the Concept2 Dyno<sup>®</sup> was assessed using a repeated measures design. All subjects were familiar with the exercises (chest press, seated row, leg press) allowed by the Concept2 Dyno<sup>®</sup>. All the exercises were performed on three sessions, one week apart each, following manufacturer's guidelines and ensuring strict adherence to form. In addition, a sub-sample consisting of the individuals who were performing the exercises with maximum resistance, was also analysed in the same way to enable inferences for using the Concept2 Dyno<sup>®</sup> with well-trained athletes. Reliability was assessed examining

differences, intraclass correlation coefficient, coefficient of variation and 95% limits of agreement between sessions while sensitivity was assessed with the standard error of measurement.

## **Subjects**

46 competitive athletes (males:  $n=36$ , age  $23.3 \pm 6.8$  years, height  $1.80 \pm 0.09$  m, body mass  $82.3 \pm 15.6$  kg; females,  $n=10$ , age  $20.7 \pm 1.4$  years, height  $1.65 \pm 0.09$  m, body mass  $62.7 \pm 11.8$  kg) free of any medical conditions or injuries in the 6 months prior to the investigation, agreed to participate in the study. The subjects competed in sports or events where strength and power was a significant aspect of successful performance. All subjects trained regularly with resistance (2-3 times per week) as part of their sport training programme for at least 2 years prior to the experiment. Although not specifically training using the Concept2 Dyno<sup>®</sup>, subjects were familiar and have been using the three exercises in their training programmes. Institutional ethical approval was granted and detailed information regarding the nature and purpose of the study was provided to prospective participants before they completed informed consent forms.

## **Procedures**

A repeated measures design was employed to determine the reliability and sensitivity of the Concept2 Dyno<sup>®</sup>. Participants attended the laboratory on four separate sessions. The first session ensured that subjects were familiar with the testing procedures and were able to maintain safe and controlled technique. The resistance level for each subject for each exercise was also determined in this session. This was established as the maximum resistance the

subject could move without any obvious fluctuations in velocity. Height was recorded to the nearest 0.01 m using a stadiometer (Harpenden, Burgess Hill, UK) and body mass was measured to the nearest 0.1 kg utilising calibrated balance scales (Seca, Birmingham, UK).

The three experimental sessions comprised of performing all three exercises of seated chest press (CPress), seated row (SRow), and seated leg press (LPress). The order of the exercises and the resistance level for each exercise were maintained the same for all sessions for each subject. For each exercise, the subjects performed three low-intensity repetitions and, immediately after, three maximal effort repetitions, according to manufacturer's guidelines. Execution form was maintained throughout. For the CPress, the subjects sat in the dynamometer seat with their back straight and the legs in a comfortable position. The handle bar was set at the same height as the subject's sternum. For the SRow, the subject sat in the dynamometer seat with their back straight and the anterior upper body touching the seat backrest. The height of the bar was the same as in the CPress. Finally, for the LPress, the subjects adopted a similar position as in the CPress, with the difference that they held their body stable by holding handles below the seat. For each effort, the weight 'pushed' or 'pulled' (in kg) was displayed on the equipment's screen (Dyno II, Nottingham, UK). The best score from the maximal effort repetitions was recorded and used for subsequent analysis.

The testing sessions were conducted at similar times of day ( $\pm$  2hrs) and under similar environmental conditions, to mediate the confounding effects of circadian rhythms and environmental influences on performance (2). All testing sessions for each subject took part within the same training phase. Subjects were instructed to refrain from strenuous exercise in the 48 hours preceding testing, and to ensure they were adequately hydrated and consumed the same diet prior to each testing session. The aforementioned controls minimised the



influence of extraneous variables on their strength performance, thus, enhancing this study's internal validity and reliability. One week intervened each experimental session.

### **Statistical analyses**

Descriptive statistics are reported as mean  $\pm$  SD unless otherwise stated. Normality of data was examined using the Shapiro-Wilk test and subsequently confirmed. Homoscedasticity of data was examined and found present for the LPress data, therefore all data was logarithmically transformed for consistency. Reliability was assessed according to suggestions by Atkinson and Nevill (1). A repeated measures analysis of variance (ANOVA) was used to assess systematic bias between the three testing occasions. The intraclass correlation coefficient ( $ICC_{3,1}$ ) was calculated from ANOVA statistics (19) as a measure of relative reliability (the degree at which the subjects maintain their rank in the sample).  $ICC_{3,1}$  was calculated between the first and subsequent sessions to examine whether reliability improved with more sessions (i.e. session 1 to 2 and session 2 to 3). In addition, coefficient of variation (CV) and 95% limits of agreement (95%LoA; 4) were calculated as measures of absolute reliability (the degree of variability in the repeated measures for each individual). CV was calculated as standard deviation / mean (17) and antilog was taken. Similar to the  $ICC_{3,1}$ , CV and 95%LoA were calculated between the first and subsequent sessions. Finally, standard error of measurement (SEM) was calculated as standard deviation  $\times$  square root (1- $ICC$ ) (20).

In addition, the same statistical analysis took place for a sub-sample of subjects (CPress, n = 28; SRow, n = 27; LPress, n = 37) that were able to execute the exercises with the maximum possible resistance (i.e. all damper levers open). Statistical tests were performed utilising the

statistical package for social sciences version 16.0 (SPSS Inc, USA). Significance level was set at  $P \leq 0.05$ .

## **RESULTS**

Descriptive statistics of all three exercises and trials for both full and sub-sample can be found in Table 1. There was no systematic bias present between trials for the logarithmically transformed CPress ( $P = 0.784$ ), SRow ( $P = 0.464$ ) or LPress ( $P = 0.195$ ) when the whole sample was considered. Similarly, no systematic bias was present for the sub-sample between trials for CPress ( $P = 0.955$ ), SRow ( $P = 0.799$ ) or LPress ( $P = 0.61$ ).

High  $ICC_{3,1}$  (0.89 – 0.98) and low CV (6.2% - 4.3%) values indicated reliable repetition of performance for all exercises and both sample. 95%LoA ranged was from -15% to 19%.

Finally, SEM values for all exercises indicated reasonable sensitivity for both samples. All statistics for  $ICC_{3,1}$ , CV, 95%LoA and SEM for the whole and sub-samples can be found in Table 2.

## **DISCUSSION**

The novel finding of the current study was that the Concept2 Dyno<sup>®</sup> produced acceptably reliable results on the seated chest press (CPress), seated row (SRow), and seated leg press (LPress) as indicated by the high ICC for both the whole and the sub-sample (subjects that utilised maximum resistance). Acceptable sensitivity was demonstrated with both the whole sample and the sub-sample, with the values produced indicated a small SEM.

The ability to assess strength in the field is important as it can be used to evaluate and monitor progress and subsequently inform training. An important aspect of any performance-measuring equipment is the ability to reproduce results under the same testing conditions, thus allowing attributing any observed changes to the training progress. The results of the current study showed that the Concept2 Dyno<sup>®</sup> produced similar scores on all three trials for all exercises for both samples examined in the study, as no bias was present in any of the above scores.

Various suggestions have been made for the value of ICC that indicates good reliability, making interpretation of ICC challenging. For example, Fleiss (6) proposed ‘excellent’ reliability with an ICC > 0.75, while Vincent and Weir (20) suggested ‘high’ reliability with an ICC > 0.90. The results of the current study produced a range of ICC of 0.89 – 0.98, indicating good to high reliability for all exercises. As with the CV, these high ICC are comparable with those found for field tests such as the 1 repetition maximum with chain-loaded bars (ICC 0.93-0.99; 14).

A CV value of 10% has been routinely used as a threshold for consistency in reliability studies, with lower CV values considered to indicate ‘low’ variability; however, there is a lack of justification for the use of this value (1). Variability of test-retest with the Concept2 Dyno<sup>®</sup> dynamometer yielded CV values ranging from 4.3% to 6.2%. These values are slightly higher than other methods of measuring strength in the field, such as 1 repetition maximum with chain-loaded bars (CV 2.5%; 14) or an accelerometer to assess loaded squat jumps (CV 1.8%-3.2%; 3). However, they are still sufficiently low to detect performance changes. It must be noted that the percentages provided here were derived from the CV

antilogs. Although this conversion results in a ratio value, the ratios were sufficiently small to be presented as percentages, and hence be easier to interpret.

Suggestions towards more sport-specific performance testing have been made (15,17) to allow for greater similarity to actual performance, and therefore, achieve more meaningful results (11,15). Notwithstanding the efficiency or specificity of a test, in order to be meaningful, its sensitivity is a crucial factor (10). Indeed, the ability of muscular function tests to detect performance changes has been questioned (16). The 95% LoA indicated that subsequent testing sessions can underestimate, on average, by a factor of 0.87 or overestimate, on average, by a factor of 1.17 (Table 2). Therefore, any changes in performance must be outside these limits to indicate progression, or indeed, decrease in performance. The use of 'analytical goals' (1) would help practitioners and researchers in making decisions as to the use of Concept2 Dyno® for their purposes.

In addition to the above, the SEM scores provide a threshold at which any change in performance score below the SEM cannot be interpreted as a real change, but rather, as random variation (e.g. from biological variation) of the test, assisting the coach in making informed decisions about the an athlete's improvement. The SEM values identified for both samples are sufficiently small to make the exercises sensitive enough to detect real changes in performance. For example, following a 4-week traditional resistance exercise programme, strength was significantly improved by 23.7% for chest press, 25% for seated row and 25.4% for leg press (12). These improvements compare favourably with the percent the SEM scores represent for each exercise (~4% - 4.5% for CPress, ~5% - 6% for SRow, and ~5% - 5.5% for LPress), indicating that Concept 2 Dyno® could serve as an assessment tool.

It has been well established that familiarisation of the subjects with the testing procedure, in order to avoid any variation in performance due to motivation or learning effects, is vital during any performance testing (10). Indeed, a review of 17 studies with three trials or more revealed that the CV can be as high as 1.3 times the CV obtained from comparing the subsequent trials (10). The ratio CV of the raw data (CV of trial 1-2 / CV of trial 2-3) obtained for the whole sample for CPress, SRow and LPress were 1.2, 0.9 and 0.90, respectively. Further, the ratio CV for the sub-sample CPress, SRow and LPress were 1.2, 0.96 and 1.07, respectively. As subjects underwent a familiarisation trial, it appears that a single session was sufficient to ensure consistent performance between trials. Therefore, we suggest that one familiarisation trial should be allowed before any assessment takes place in athletes.

The ability of the athlete to select the resistance level that is most appropriate for them prior to each exercise must be considered in relation to the practical implications it has. As the athlete's ability to generate force changes with training, it is possible that the resistance level would also need to change, to accommodate increases in strength or power development. For power development, in particular, the wide range of loads used to produce optimal power (5) may present another implication to the use of the dynamometer. It is suggested that changes in performance when using the dynamometer are assessed by comparing results using the same resistance level.

The exercises used (chest press, seated row, leg press) are common exercises performed by athletes in various training facilities, hence the close resemblance to these moves and the fact it measures the performance itself, offers logical validity. However, the present study did not

include any validity assessment of the Concept2 Dyno®. As a result, the findings of the present study relate to performance measurements obtained by the Concept2 Dyno® only and the performance scores from it cannot be compared to other isoinertial strength measures using e.g. free weights.

## **PRACTICAL APPLICATIONS**

Concept2 Dyno® is a reliable strength testing equipment for competitive athletes who utilise strength training as part of their programme. All three exercises were found to have good reliability (high ICC and low CV) and acceptable sensitivity, making the dynamometer sufficiently sensitive to detect small changes in athletes' performance. Performance in leg press should be assessed using the coefficient of variation rather the raw score. In addition, care needs to be given to maintaining execution form throughout performance as well as maintaining the same execution form for repeat performances. From knowledge of the mechanics behind the Concept2 Dyno®, i.e. fluid resistance, validity may develop in sports that involve motion that is loaded in a similar manner (i.e. swimming, water polo). However, there may also be benefits for athletes over a broader spectrum of sports where such assessment and exercise provides additional challenges considering variation of load.

## REFERENCES

1. Atkinson, G, and Nevill, AM. Statistical methods for assessing measurement error (reliability) in variables relevant to sports medicine. *Sports Med* 26: 217-238, 1998.
2. Atkinson, G, and Reilly, T. Circadian variation in sports performance. *Sports Med* 21: 292-312, 1996.
3. Bampouras, TM, Relph, NS, Orme, D, and Esformes, JI. Validity and reliability of the Myotest Pro wireless accelerometer in squat jumps. *Isokinetics and Exercise Science*, 21: 101-105, 2013.
4. Bland, JM, and Altman, DG. Measuring agreement in method comparison studies. *Stat Methods Med Res* 8: 135-160, 1999.
5. Cormie, P, McBride, JM, and McCaulley, GO. Power-time, force-time, and velocity-time curve analysis during the jumps squat: impact of load. *J Appl Biomech* 24: 112-120, 2008.
6. Fleiss, JL. Design and analysis of clinical experiments. New York, NY: John Wiley & Sons, 1986.
7. Gioftsidou, A, Ispirlidis, I, Pafis, G, Malliou, P, Bikos, C, and Godolias, G. Isokinetic strength training program for muscular imbalances in professional soccer players. *Sport Sci Health* 2:101-105, 2008.
8. Graham-Smith, P, Burgess, K, and Ridler, A. The relationship between strength, power, flexibility, anthropometry and technique and 2000 m and 5000 m rowing ergometer performance. In: Kinanthropometry X. Marfell-Jones, M., and Olds, T, eds. New York: Routledge, 2007. pp. 135-150.
9. Hoffman, JR, Kang, J. Strength changes during an in-season resistance-training program for football. *J Strength Cond Res* 17: 109-114, 2003.

10. Hopkins, WG, Schabert, EJ, and Hawley, JA. Reliability of power in physical performance tests. *Sports Med* 31: 211-234, 2001.
11. Jones, P, and Bampouras, TM. A comparison of isokinetic and functional methods of assessing bilateral strength imbalance. *J Strength Cond Res* 24: 1553–1558, 2010.
12. Kim E, Dear A, Ferguson SL, Seo D, Bemben MG. Effects of 4 weeks of traditional resistance training vs. superslow strength training on early phase adaptations in strength, flexibility, and aerobic capacity in college-aged women. *J Strength Cond Res*. 25(11): 3006-3013, 2011.
13. Marrin, K, Bampouras, TM. Anthropometric and physiological changes in elite female water polo players during a training year. *Serbian J Sports Sci* 2: 75–84, 2008.
14. McCurdy, K, Langford, G, Jenkerson, D, and Doscher, M. The validity and reliability of the 1RM bench press using chain-loaded resistance. *J Strength Cond Res* 22: 678-683, 2008.
15. Muller, E, Benko, U, Raschner, C, and Schwameder, H. Specific fitness training and testing in competitive sports. *Med Sci Sports Exerc* 32: 216–220, 2000.
16. Murphy, AJ, and Wilson, GJ. The ability of tests of muscular function to reflect training-induced changes in performance. *J Sports Sci* 15: 191–200, 1997.
17. Sale, DG. Testing strength and power. In: MacDougall JD, Wenger HA, Green HJ, eds. *Physiological Testing of the High Performance Athlete*. 2nd ed. Champaign, IL: Human Kinetics, 1991. pp. 21 – 106.
18. Sankey, SP, Jones, PA, Bampouras, TM. Effects of two plyometric training programmes of different intensity on vertical jump performance in high school athletes. *Serbian J Sports Sci* 2: 123–130, 2008.



19. Shrout, PE, and Fleiss, JP. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull* 86: 420-428, 1979.
20. Vincent, WJ, and Weir, JP. *Statistics in Kinesiology*. Champaign, IL: Human Kinetics Books, 2012.
21. Walsh, MS, Ford, KR, Bangen, KJ, Myer, GD, and Hewett, TE. The validation of a portable force plate for measuring force-time data during jumping and landing tasks. *J Strength Cond Res* 20: 730-734, 2006.

## Tables Legends

Table 1. Descriptive statistics (mean  $\pm$  SD) for the seated bench press (CPress), seated bench pull (SRow), and seated leg press (LPress) from the three trials for both the whole and sub-sample. The sub-sample descriptives refer to the subjects that executed the exercises with maximum resistance on the dynamometer.

Table 2. Reliability and sensitivity statistics (intraclass correlation coefficient, ICC<sub>3,1</sub>; coefficient of variation, CV; range of 95% limits of agreement, 95% LoA (range); standard error of measurement (SEM)) for all exercises between trials, for both the whole and sub-samples. The sub-sample statistics refer to the subjects that executed the exercises with maximum resistance on the dynamometer.

Table 1.

Exercise	Whole sample			Sub-sample		
	Trial			Trial		
	1	2	3	1	2	3
CPress (kg)	75.6 ±	76.1 ±	75.8 ±	88.0 ±	87.8 ±	88.1 ±
	22.0	22.2	22.6	16.0	17.0	17.1
SRow (kg)	71.5 ±	71.4 ±	70.9 ±	83.1 ±	82.6 ±	82.7 ±
	19.5	20.3	21.0	13.8	16.0	16.8
LPress (kg)	177.7 ±	175.4 ±	173.2 ±	186.4 ±	183.6 ±	179.8 ±
	37.4	37.4	39.6	33.7	34.8	36.5

Table 2.

Trial	Whole sample								Sub-sample							
	ICC		CV (%)		95% LoA (range)		SEM (kg)		ICC		CV (%)		95% LoA (range)		SEM (kg)	
	1-2	2-3	1-2	2-3	1-2	2-3	1-2	2-3	1-2	2-3	1-2	2-3	1-2	2-3	1-2	2-3
Exercise																
CPress	0.98	0.98	5.1	4.3	0.87 – 1.14	0.90- 1.13	3.1	3.2	0.93	0.96	5.3	4.4	0.86 – 1.18	0.88 – 1.13	4.4	3.4
SRow	0.97	0.97	5.1	5.5	0.87 – 1.15	0.87 – 1.17	3.4	3.6	0.89	0.90	6.0	6.2	0.86 – 1.19	0.85 – 1.18	4.9	5.2
LPress	0.94	0.94	5.6	6.2	0.87 – 1.18	0.86 – 1.19	9.2	9.4	0.91	0.93	6.0	5.6	0.87 – 1.19	0.88 – 1.18	10.3	9.4