

Relph, Nicola ORCID: <https://orcid.org/0000-0003-3323-3444> and Herrington, Lee (2015) Criterion-related validity of knee joint-position-sense measurement using image capture and isokinetic dynamometry. *Journal of Sport Rehabilitation* .

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/1721/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
 - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

Criterion-Related Validity of Knee Joint-Position-Sense Measurement Using Image Capture and Isokinetic Dynamometry

Nicola Relph and Lee Herrington

Context: Clinicians require portable, valid, and cost-effective methods to monitor knee joint-position-sense (JPS) ability. **Objective:** To examine the criterion-related validity of image-capture JPS measures against an isokinetic-dynamometer (IKD) procedure. **Design:** Random crossover design providing a comparison of knee JPS measures from image capture and IKD procedures. **Participants:** 10 healthy participants, 5 female, age 28.0 ± 13.29 y, mass 60.3 ± 9.02 kg, height 1.65 ± 0.07 m, and 5 male, 29.6 ± 10.74 y, mass 73.6 ± 5.86 kg, height 1.75 ± 0.07 m. **Main Outcome Measures:** The dependent variables were absolute error scores (AES) provided by 2 knee directions (flexion and extension). The independent variables were the method (image capture and IKD). **Results:** There was no significant difference between clinical and IKD AES into knee-flexion data ($P = .263$, $r = 0.55$). There was a significant difference between clinical and IKD AES into knee-extension data ($P = .016$, $r = .70$). **Conclusions:** Analysis of photographic images to assess JPS measurements using knee flexion is valid against an IKD positioning method, but JPS measurements using knee extension may not be valid against IKD techniques. However, photo-analysis measurements provided a lower error score using knee-extension data and thus may provide an optimal environment to produce maximal knee JPS acuity. Therefore, clinicians do not need expensive equipment to collect representative JPS ability.

Keywords: proprioception, isokinetic dynamometer

Clinicians use knee joint-position-sense (JPS) measurements to assess static knee proprioception ability.¹ This is an important measurement, as it can either identify patients with a JPS deficiency that may lead to an increased risk of knee injury or progress along a proprioceptive-based rehabilitation program. The traditional clinical JPS measurement technique involves passive knee movement by the clinician to a specific target angle, then active reproduction of this angle by the patient.¹ Image capture can be used to collect knee position and hence knee JPS information. However, as the clinician is part of this data-collection process, measurement bias² may be introduced to the data. Therefore, an isokinetic dynamometer (IKD) provides an alternate means to position the knee target angle, removing researcher bias. Kiran et al³ reported high correlations between concurrent measurement of JPS using an IKD, photo analysis, and electrogoniometry. However, all target knee positions were completed by the IKD arm and therefore did not replicate a typical clinical setting. Grob et al⁴ did consider the correlation between a

self-built low-speed motor and passive researcher positioning techniques on different occasions. Results indicated a poor correlation between the 2 measurements ($r = -.2$), suggesting that the methods should not be used interchangeably. It is notable that when the target angle was positioned by the researcher rather than a pulley system, participants produced better JPS acuity results. However, the matching method was produced using a visual analog scale, which has limited ecological validity.⁵

Smith et al⁵ produced a systematic review on the reliability of JPS measurement techniques. Their findings suggested that intrarater reliability depended on data-acquisition techniques; image capture produced greater reliability than electrogoniometry and dynamometry. However, no study has considered the concurrent validity of assessment methods using the same participants.⁵ An analysis of the validity of JPS techniques is difficult, as there is no universally accepted "gold standard" method of collecting JPS data. However, the use of an IKD to position a limb at a defined angle is accepted. Therefore, criterion-related, specifically concurrent validity was investigated in this study by comparing a clinical JPS measurement technique with an IKD JPS protocol. Concurrent validity is defined as a comparison between 1 previously validated protocol and a new or previously unvalidated procedure.⁶ Clinicians use JPS to measure the effectiveness of a rehabilitation program, so it is

Commented [JG1]: Please provide 1 or 2 more keywords/phrases for indexing purposes that are not included in the title.

Relph is with the Dept of Medical and Sport Sciences, University of Cumbria, Carlisle, UK. Herrington is with the School of Health Science, University of Salford, Salford, UK. Address author correspondence to Nicola Relph at Nicola.Relph@Cumbria.ac.uk.

knee-flexion data ($P = .263$, $r = .55$). There was a significant difference between image-capture AES ($2.5^\circ \pm 0.7^\circ$) and IKD AES ($4.3^\circ \pm 1.9^\circ$) knee-extension data ($P = .016$, $r = .70$).

Discussion

Clinicians use JPS to measure the effectiveness of a rehabilitation program and identify patients who may be more at risk for knee injury, so it is imperative that the measurements be valid. Criterion-related validity was confirmed for knee-flexion JPS; there were no differences between JPS in a clinical and IKD setting ($P = .263$, $r = .55$). However, knee-extension JPS using an image-capture technique was different than an IKD-based technique ($P = .263$, $r = .7$). The IKD data provided significantly greater error scores than the image-capture data for knee extension. This supports previous evidence that JPS measurement techniques should not be used interchangeably; however, passive positioning by a researcher may provide a more optimal environment for maximal JPS performance.⁴ It is possible in the IKD setting that participants had to adapt to the addition of the lever arm increasing the mass of the leg and the torque required to extend the knee; hence, effort was not as natural when compared with the image-capture setting and ecological validity was reduced. This may not have the same effect on knee flexion, as the torque required in this direction would be assisted by gravity. Another feasible explanation was the seating in both tests. In the image-capture test condition participants were seated on the edge of a plinth and hence were not conscious of a back rest and could use pelvis rotation to assist knee extension and the associated hamstring lengthening. Previous research suggests heightened afferent information when muscles are lengthened.¹⁰ In the IKD setting participants were seated on the edge of the seat and not supported by the back rest but may have been less likely to use pelvis rotation to assist knee extension and hence perhaps use a less natural (more resistance to) knee-extension movement. Therefore, a clinical setting may provide a more “optimal” environment for knee-extension JPS measurement, as ecological validity is increased.

Results of this validity study have important implications for clinicians. The image-capture measurement of knee JPS with passive positioning of target angles produced similar (knee flexion) and improved (knee extension) AES compared with the IKD

setting. This suggests that a clinical measurement technique provides a more optimal environment and “best scores” for JPS than an IKD setting. Therefore, knee JPS can be measured in a clinical setting using cheap and easily accessible equipment; expensive IKD equipment is not necessary.

References

1. Beynon BD, Renstrom PA, Konradsen L, Elmqvist LG, Gottlieb D, Dirks M. Validation of techniques to measure knee proprioception. In: Lephart SM, Fu FH, eds. *Proprioception and Neuromuscular Control in Joint Stability*. Champaign, IL: Human Kinetics; 2000:127–138.
2. Sica GT. Bias in research studies. *Radiology*. 2006;238(3):780–789. [PubMed](#) [doi:10.1148/radiol.2383041109](https://doi.org/10.1148/radiol.2383041109)
3. Kiran D, Carlson M, Medrano D, Smith DR. Correlation of three different knee joint position sense measures. *Phys Ther Sport*. 2010;11:81–85. [PubMed](#) [doi:10.1016/j.ptsp.2010.06.002](https://doi.org/10.1016/j.ptsp.2010.06.002)
4. Grob KR, Kuster MS, Higgins SA, Lloyd DG, Yata H. Lack of correlation between different measurements of proprioception in the knee. *J Bone Joint Surg Br*. 2002;84(4):614–618. [PubMed](#) [doi:10.1302/0301-620X.84B4.11241](https://doi.org/10.1302/0301-620X.84B4.11241)
5. Smith TO, Davies L, Hing CB. A systematic review to determine the reliability of knee joint position sense assessment measures. *Knee*. 2013;20(3):162–169. [PubMed](#) [doi:10.1016/j.knee.2012.06.010](https://doi.org/10.1016/j.knee.2012.06.010)
6. Sim J, Arnell P. Measurement validity in physical therapy research. *Phys Ther*. 1993;73(2):102–110. [PubMed](#)
7. Payton CJ. Motion analysis using video. In: Payton, CJ, Bartlett, RM, eds. *Biomechanical Evaluation of Movement in Sport and Exercise: The British Association of Sport and Exercise Sciences Guidelines*. London: Routledge; 2008:8–32.
8. Andersen SB, Terwilliger DM, Denegar CR. [Comparison of open versus closed kinetic chain test positions for measuring joint position sense](#). *J Sport Rehabil*. 1995;4(3):165–171.
9. Field A. *Discovering Statistics Using SPSS*. London, UK: Sage; 2005.
10. Ribot-Ciscar E, Roll JP. Ago-antagonist muscle spindle inputs contribute together to joint movement coding in man. *Brain Res*. 1998;791(1-2):167–176. [PubMed](#) [doi:10.1016/S0006-8993\(98\)00092-4](https://doi.org/10.1016/S0006-8993(98)00092-4)

Appendix: Questionnaires Used to Define Participants' Knee-Function Score (KOOS and Lysholm) and Activity Level (GPPAQ and Tegner)



General Practice Physical Activity Questionnaire

Date.....

Name.....

1. Please tell us the type and amount of physical activity involved in your work.

		Please mark one box only
a	I am not in employment (e.g. retired, retired for health reasons, unemployed, full-time carer etc.)	
b	I spend most of my time at work sitting (such as in an office)	
c	I spend most of my time at work standing or walking. However, my work does not require much intense physical effort (e.g. shop assistant, hairdresser, security guard, childminder, etc.)	
d	My work involves definite physical effort including handling of heavy objects and use of tools (e.g. plumber, electrician, carpenter, cleaner, hospital nurse, gardener, postal delivery workers etc.)	
e	My work involves vigorous physical activity including handling of very heavy objects (e.g. scaffolder, construction worker, refuse collector, etc.)	

2. During the last week, how many hours did you spend on each of the following activities?
Please answer whether you are in employment or not

Please mark one box only on each row

	None	Some but less than 1 hour	1 hour but less than 3 hours	3 hours or more
a	Physical exercise such as swimming, jogging, aerobics, football, tennis, gym workout etc.			
b	Cycling, including cycling to work and during leisure time			
c	Walking, including walking to work, shopping, for pleasure etc.			
d	Housework/Childcare			
e	Gardening/DIY			

3. How would you describe your usual walking pace? Please mark one box only.

Slow pace (i.e. less than 3 mph)	<input type="checkbox"/>	Steady average pace	<input type="checkbox"/>
Brisk pace	<input type="checkbox"/>	Fast pace (i.e. over 4mph)	<input type="checkbox"/>

Appendix Figure 1 — GPPAQ.

Commented [JG2]: As I indicated in an earlier e-mail, we cannot use these images without permission from the copyright holder, if indeed they are copyrighted. You will need to determine this and obtain permission if necessary before we can publish these.

Knee injury and Osteoarthritis Outcome Score (KOOS), English version LK1.0

KOOS KNEE SURVEY

Today's date: ___/___/___ Date of birth: ___/___/___

Name: _____

INSTRUCTIONS: This survey asks for your view about your knee. This information will help us keep track of how you feel about your knee and how well you are able to perform your usual activities. Answer every question by ticking the appropriate box, only one box for each question. If you are unsure about how to answer a question, please give the best answer you can.

Symptoms

These questions should be answered thinking of your knee symptoms during the **last week**.

S1. Do you have swelling in your knee?
Never Rarely Sometimes Often Always

S2. Do you feel grinding, hear clicking or any other type of noise when your knee moves?
Never Rarely Sometimes Often Always

S3. Does your knee catch or hang up when moving?
Never Rarely Sometimes Often Always

S4. Can you straighten your knee fully?
Always Often Sometimes Rarely Never

S5. Can you bend your knee fully?
Always Often Sometimes Rarely Never

Stiffness

The following questions concern the amount of joint stiffness you have experienced during the **last week** in your knee. Stiffness is a sensation of restriction or slowness in the ease with which you move your knee joint.

S6. How severe is your knee joint stiffness after first wakening in the morning?
None Mild Moderate Severe Extreme

S7. How severe is your knee stiffness after sitting, lying or resting **later in the day**?
None Mild Moderate Severe Extreme

Appendix Figure 2(a) — KOOS.

Knee injury and Osteoarthritis Outcome Score (KOOS), English version LK1.0

Pain

P1. How often do you experience knee pain?

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| Never | Monthly | Weekly | Daily | Always |
| <input type="checkbox"/> |

What amount of knee pain have you experienced the **last week** during the following activities?

P2. Twisting/pivoting on your knee

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

P3. Straightening knee fully

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

P4. Bending knee fully

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

P5. Walking on flat surface

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

P6. Going up or down stairs

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

P7. At night while in bed

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

P8. Sitting or lying

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

P9. Standing upright

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

Function, daily living

The following questions concern your physical function. By this we mean your ability to move around and to look after yourself. For each of the following activities please indicate the degree of difficulty you have experienced in the **last week** due to your knee.

A1. Descending stairs

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

A2. Ascending stairs

- | | | | | |
|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| None | Mild | Moderate | Severe | Extreme |
| <input type="checkbox"/> |

Appendix Figure 2(b) — KOOS.

Knee injury and Osteoarthritis Outcome Score (KOOS), English version LK1.0

For each of the following activities please indicate the degree of difficulty you have experienced in the **last week** due to your knee.

A3. Rising from sitting	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A4. Standing	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A5. Bending to floor/pick up an object	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A6. Walking on flat surface	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A7. Getting in/out of car	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A8. Going shopping	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A9. Putting on socks/stockings	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A10. Rising from bed	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A11. Taking off socks/stockings	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A12. Lying in bed (turning over, maintaining knee position)	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A13. Getting in/out of bath	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A14. Sitting	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				
A15. Getting on/off toilet	None	Mild	Moderate	Severe	Extreme
	<input type="checkbox"/>				

Appendix Figure 2(c) — KOOS.

Knee injury and Osteoarthritis Outcome Score (KOOS), English version LK1.0

For each of the following activities please indicate the degree of difficulty you have experienced in the **last week** due to your knee.

A16. Heavy domestic duties (moving heavy boxes, scrubbing floors, etc)

None Mild Moderate Severe Extreme

A17. Light domestic duties (cooking, dusting, etc)

None Mild Moderate Severe Extreme

Function, sports and recreational activities

The following questions concern your physical function when being active on a higher level. The questions should be answered thinking of what degree of difficulty you have experienced during the **last week** due to your knee.

SP1. Squatting

None Mild Moderate Severe Extreme

SP2. Running

None Mild Moderate Severe Extreme

SP3. Jumping

None Mild Moderate Severe Extreme

SP4. Twisting/pivoting on your injured knee

None Mild Moderate Severe Extreme

SP5. Kneeling

None Mild Moderate Severe Extreme

Quality of Life

Q1. How often are you aware of your knee problem?

Never Monthly Weekly Daily Constantly

Q2. Have you modified your life style to avoid potentially damaging activities to your knee?

Not at all Mildly Moderately Severely Totally

Q3. How much are you troubled with lack of confidence in your knee?

Not at all Mildly Moderately Severely Extremely

Q4. In general, how much difficulty do you have with your knee?

None Mild Moderate Severe Extreme

Thank you very much for completing all the questions in this questionnaire.

Appendix Figure 2(d) — KOOS.

Lysholm Knee Questionnaire / Tegner Activity Scale

Name:
First Last

Date: 09 01 14

Physician:

1. Limp:

- a) None
- b) Slight or periodical
- c) Severe and constant

2. Support:

- a) None
- b) Stick or crutch
- c) Weight-bearing impossible

3. Locking:

- a) No locking and no catching sensations
- b) Catching sensation but no locking
- c) Locking occasionally
- d) Locking frequently
- e) Locked joint on examination

4. Instability:

- a) Never giving way
- b) Rarely during athletics or other severe exertion
- c) Frequently during athletics or other severe exertion (or incapable of participation)
- d) Occasionally in daily activities
- e) Often in daily activities
- f) Every step

5. Pain:

- a) None
- b) Inconstant and slight during severe exertion
- c) Marked during severe exertion
- d) Marked on or after walking more than 2 km
- e) Marked on or after walking less than 2 km
- f) Constant

6. Swelling:

- a) None
- b) On severe exertion
- c) On ordinary exertion
- d) Constant

7. Stair-climbing:

- a) No problems
- b) Slightly impaired
- c) One step at a time
- d) Impossible

8. Squatting:

- a) No problems
- b) Slightly impaired
- c) Not beyond 90°
- d) Impossible

Activity Level Before Injury	Current Activity Level	Activity Level Following Surgery if applicable	
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Competitive sports Soccer - national and international elite
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Competitive sports Soccer, lower divisions Ice hockey Wrestling Gymnastics
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Competitive sports Bandy Squash or badminton Athletics (jumping, etc.) Downhill skiing
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Competitive sports Tennis Athletics (running) Motorcross, speedway Handball Basketball Recreational sports Soccer Bandy and ice hockey Squash Athletics (jumping) Cross-country track findings both recreational and competitive
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Recreational sports Tennis and badminton Handball Basketball Downhill skiing Jogging, at least five times per week
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Work Heavy labor (e.g., building, forestry) Competitive sports Cycling Cross-country skiing Recreational sports Jogging on uneven ground at least twice weekly
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Work Moderately heavy labor (e.g., truck driving, heavy domestic work) Recreational sports Cycling Cross-country skiing Jogging on even ground at least twice weekly
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Work Light labor (e.g., nursing) Competitive and recreational sports Swimming Walking in forest possible
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Work Light labor Walking on uneven ground possible but impossible to walk in forest
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Work Sedentary work Walking on even ground possible
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	Sick leave or disability pension because of knee problems

Tegner:

Lysholm Score:

Appendix Figure 3(b) — Lysholm/Tegner.




Figure 1 — Typical setup for image-capture knee-joint position-sense measurements.

Table 1 Participant Characteristics (Mean ± SD)

	Age (y)	Mass (kg)	Height (m)	BMI	GPPAQ range	KOOS	Lysholm	Tegner
Females (n = 5)	28.0 ± 13.29	60.3 ± 9.02	1.65 ± 0.07	22.1 ± 1.80	Inactive to active	98.6 ± 3.18	98.8 ± 2.68	5.0 ± 1.22
Males (n = 5)	29.6 ± 10.74	73.6 ± 5.86	1.75 ± 0.07	24.1 ± 1.97	Active	92.5 ± 10.87	87.6 ± 17.5	7.8 ± 1.30

Abbreviations: BMI, body-mass index; GPPAQ, General Practitioner Physical Activity Questionnaire; KOOS, Knee injury and Osteoarthritis Outcome Score (the closer the score to 100, the better the knee condition); Lysholm, Lysholm Knee Score (the closer the score to 100, the better the knee condition); Tegner, Tegner Activity Scale (the closer the score to 10, the more physically active) (see Appendix for more details).

Table 2 Isokinetic Dynamometer Protocol

Action	Angle (°)	Hold time (s)
From 0° (full extension) into knee flexion		
passive	90/80/70/90/75	5
passive	0	2
active	Replication	5
passive	0	Back to step 1
From 90° into knee extension (0°)		
passive	30/45/60/45/45	5
passive	90	2
active	Replication	5
passive	90	Back to step 1

Note: Passive action defines isokinetic dynamometer lever movement. Active motion defines participant muscle contraction.