

Sellami, Maha, Wissem, Dhab, Hayes, Lawrence ORCID: <https://orcid.org/0000-0002-6654-0072> , Padulo, Johnny, Rhibi, Fatma, Djemail, Hanen and Chaouachi, Anis (2017) Combined sprint and resistance training abrogates age differences in somatotrophic hormones. PLoS ONE, 12 (8). e0183184.

Downloaded from: <https://insight.cumbria.ac.uk/id/eprint/3180/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
- a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

RESEARCH ARTICLE

Combined sprint and resistance training abrogates age differences in somatotrophic hormones

Maha Sellami¹, Wissem Dhahbi¹, Lawrence D. Hayes², Johnny Padulo^{1,3,4*}, Fatma Rhibi⁵, Hanen Djemail⁶, Anis Chaouachi¹

1 Tunisian Research Laboratory "Sport Performance Optimization" National Center of Medicine and Science in Sports, Tunis, Tunisia, **2** Active Ageing Research Group, Department of Medical and Sport Sciences, University of Cumbria, Bowerham Road, Lancaster, United Kingdom, **3** University eCampus, Novedrate, Italy, **4** Faculty of Kinesiology, University of Split, Split, Croatia, **5** Movement, Sport, Health and Sciences Laboratory (M2S), University of Rennes 2, Rennes, France, **6** Military Hospital of Instruction of Tunis, Department of Endocrinology, Tunis, Tunisia

* sportcnetic@gmail.com



Abstract

The aim of this investigation was to compare serum growth hormone (GH), insulin-like growth factor-1 (IGF-1) and insulin-like growth factor-binding protein-3 (IGFBP-3) in response to a combined sprint and resistance training (CSRT) program in young and middle-aged men. Thirty-eight healthy, moderately trained men participated in this study. Young and middle-aged men were randomly assigned to, a young training group (YT = 10, 21.4 ± 1.2 yrs) or a young control group (YC = 9, 21.6 ± 1.8 yrs), a middle-aged training group (MAT = 10, 40.4 ± 2.1 yrs) or a middle-aged control group (MAC = 9, 40.5 ± 1.8 yrs). Participants performed the Wingate Anaerobic Test (WAnT) before and after a 13-week CSRT program (three sessions per week). Blood samples were collected at rest, after warm-up, immediately post-WAnT, and 10 min post-WAnT. CSRT induced increases in GH at rest and in response to the WAnT in YT and MAT ($P < 0.05$). CSRT-induced increases were observed for IGF-1 and IGFBP-3 at rest in MAT only ($P < 0.05$). Pre-training, GH, IGF-1 and IGFBP-3 were significantly higher at rest and in response to the WAnT in young participants as compared to their middle-aged counterparts ($P < 0.05$). Post-training, YT and MAT had comparable basal GH ($P > 0.05$). In response to the WAnT, amelioration of the age-effect was observed between YT and MAT for IGF-1 and IGF-1/IGFBP-3 ratio following CSRT ($P > 0.05$). These data suggest that CSRT increases the activity of the GH/IGF-1 axis at rest and in response to the WAnT in young and middle-aged men. In addition, CSRT reduces the normal age-related decline of somatotrophic hormones in middle-age men.

OPEN ACCESS

Citation: Sellami M, Dhahbi W, Hayes LD, Padulo J, Rhibi F, Djemail H, et al. (2017) Combined sprint and resistance training abrogates age differences in somatotrophic hormones. PLoS ONE 12(8): e0183184. <https://doi.org/10.1371/journal.pone.0183184>

Editor: Alejandro Lucía, Universidad Europea de Madrid, SPAIN

Received: March 1, 2017

Accepted: July 29, 2017

Published: August 11, 2017

Copyright: © 2017 Sellami et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

The aging process is associated with a precipitous decline in skeletal muscle mass and strength, estimated as 35–40% between 20 and 80 yrs[1], with an accelerated decline after 50 yrs[2].

Moreover, cross-sectional studies have observed that the ability to develop muscle strength and power declines from 40 to 80 yrs [3, 4]. It has been suggested that reduced muscle function may result from neural degeneration combined with muscle atrophy. Muscle atrophy occurs when muscle protein degradation exceeds muscle protein synthesis. Said muscle atrophy may be partly attributable to a reduction in anabolic hormone production [5], whilst age-associated physical inactivity may exacerbate muscle loss. Moreover, the reduction in systemic anabolic hormones may be further exacerbated by physical inactivity in older adults [6].

The systemic reduction in insulin-like growth factor-1 (IGF-1) has been attributed, in part, to decreased secretion of growth hormone (GH), the main secretagogue of IGF-1. GH secretion reduces ~14% per decade after the second decade [7], and reaches, by the age of ~60 yrs, half of the GH secretion of younger counterparts (20–30 yrs [8]). IGF-1, the main stimulated protein downstream of GH, concomitantly decreases with age (~10% per decade) and is associated with cell proliferation, cell differentiation, energy metabolism and prevention of apoptosis [9]. Most IGF-1 circulates in the blood bound to IGF-binding proteins (IGFBPs) which mediate bioavailability into tissue [10]. The most abundant is the insulin-like growth factor-binding protein 3 (IGFBP-3), which carries 90–95% of IGF-1 in circulation [11]. Hence, it is necessary to consider IGFBP-3 when total serum IGF-1 is determined to provide a measure of bioavailability. Serum IGFBP-3 is regulated by GH signaling [12, 13] and reduced with advanced age [14]. The main function of IGFBP-3 is to permit IGF-1 transport and to regulate anti-proliferative and apoptotic effects through cell surface receptor by opposing IGF-1 action [15, 16]. Moreover, IGFBP-3 is a valuable tool for diagnosis of GH perturbation during the aging process [17].

Because GH naturally declines with age, there has been increasing interest in the role of GH as an anti-aging factor, and some evidence suggests GH administration increases fatty acid oxidation, protein synthesis and, consequently, lean body mass [18]. To date, however, there have been no approved clinical studies to assess the effects of GH administration on muscle mass loss with aging. In fact, GH administration has been shown to have no effect on muscle function in healthy aged men and women [19]. As GH administration is widely used as a muscle mass and performance enhancer [20], whether exercise could be a nonpharmacological intervention to enhance GH and subsequently muscle mass and muscle strength, requires further investigation.

Whilst acute exercise-induced elevations in GH and IGF-1 are consistently reported [20–26], the effect of long-term exercise training on basal GH and IGF-1 secretion is more ambiguous. It is well known that GH response exercise training depends on several factors including age, diet, stress, or training intensity [20, 27]. Metabolic (i.e. glucose) and hormonal (i.e. catecholamines, cortisol, testosterone) factors also influence GH [28–30] which in turn are also dependent upon exercise intensity and training type [31, 32]. In this context, increase of these hormones is greater following anaerobic training in young and middle-aged trained men [6, 31–33]. Moreover, Nevill et al. [34] suggested that serum GH response to treadmill running was greater in sprinters compared to endurance athletes. In addition, endurance training (running for eight weeks) resulted in increased systemic IGF-1 (+15%) in subjects aged ~66 yrs [35], yet Vitiello, Wilkinson [36] observed no perturbation in IGF-1 amongst moderately and well-trained endurance athletes aged ~69 yrs. As such, intensive training using sprint or resistance training appears necessary to induce changes to serum GH. Moreover, numerous studies suggest sprint training may elicit greater increase in muscle strength than the endurance training [31, 34]. Borst et al. [37] suggested that 25 weeks' resistance training in middle-aged men (~37 yrs) resulted in increased GH and IGF-1 with increase in strength performances. However, Adams et al. [38] suggested it necessary to add different types of exercise to resistance exercise in order to obtain higher muscle power during sprint exercise. In fact, some

authors suggest combined sprint and resistance training is more efficient than sprint training only [39] or resistance training only [40].

Most existing GH and IGF-1 exercise studies have focused on young (~20 yrs) and elderly (>65 yrs) subjects. Therefore, there is a paucity of data concerning the effect of concurrent exercise training (sprint interval and resistance exercises) on somatotrophic hormones in adults ~40 yrs. Thus, we examined the effect of 13 weeks' concurrent training on GH, IGF-1, and IGFBP-3 in young (~20 yrs) and middle-aged (~40 yrs) men before and after supramaximal exercise. We hypothesized *a priori* that younger participants would have greater GH, IGF-1, and IGFBP-3. Moreover, we hypothesized that training would increase somatotrophic hormone concentrations in both age groups.

Materials and methods

Participants

Military Participants reviewed and signed consent forms specifically approved by the "Department of staff and training committee" (Bouchoucha, Tunisia). The "Department of staff and training committee" approved the entire study design which has been conducted according to the principles expressed in the Declaration of Helsinki.

During the design of the study, statistical power analysis was carried out to calculate sample size. This procedure showed that nine subjects for each of four groups were needed to achieve a statistical power of 80% to detect a small effect ($d = 0.29$) when assessed by four-factor mixed analysis of variance (ANOVA) with a level of significance of 5%. Therefore, thirty-eight healthy, moderately trained men (military participants) were recruited for participation in the present investigation.

To assess the physical condition of military participants, we used an adapted version of the Baecke questionnaire [41]. Before the study, subjects performed 1 h of running and leisure-time physical activity at least three times per week ($180 \text{ min} \cdot \text{week}^{-1}$). Inclusion criteria included the absence of the following: contraindications to maximal exercise testing (e.g., cardiovascular or pulmonary disease); endocrine disorders; metabolic syndrome symptoms (e.g., hypertension, impaired fasting glucose). Following recruitment and familiarization, participants completed medical history and dietary questionnaires. Thereafter, young and middle-aged participants were randomized to receive 13 weeks' combined sprint and resistance training (CSRT), or control. Therefore, four groups existed: a young training group (YT = 10; age 21.4 ± 1.2 yrs, body height 178.3 ± 3.2 cm, body mass 74.4 ± 5.4 kg), a young control group (YC = 9; age 21.6 ± 1.8 yrs, body height 179.7 ± 6.4 cm, body mass 69.5 ± 7.3 kg), a middle-aged training group (MAT = 10; age 40.4 ± 2.1 yrs, body height 175.8 ± 5.2 cm, body mass 78.4 ± 5.2 kg) and a middle-aged control group (MAC = 9; 40.5 ± 1.8 yrs, body height 177.3 ± 4.4 cm, body mass 76.6 ± 3.9 kg).

A conventional dietary survey was conducted by a sports nutritionist of the Department of Physical Education and Military Sport to monitor individual participants' diets over the 13 weeks. Participants were asked to abstain from high glycemic loads, saturated and trans-fatty acids, caffeine, alcohol, drugs, vitamins or supplements, and low-fiber diets for the duration of the study.

Before training (during the medical examination), subjects were familiarized with testing procedures to negate learning effect. Participants avoided physical activity for 48 h preceding each test. The testing period was divided into two phases: before (P1), and after (P2), training. Each period lasted seven days and included anthropometric measurements and two consecutive laboratory visits separated by 48 h. The second phase (P2) began 48 h after training cessation and finished seven days later. All tests were performed in the morning 2h postprandial

(standard breakfast: 10 kcal·kg⁻¹, 55% carbohydrate, 33% lipids and 12% protein). Anthropometric parameters were measured on the morning of the first day. Measurement of body mass (kg) and height (cm) were taken from all participants. Body mass was measured to the nearest 0.1 kg, with subjects in light clothing and without shoes, using electronic scales (Kern, MFB 150K100). Height was determined to the nearest 0.5cm with a measuring tape fixed to the wall. Subsequently, skin-folds were measured using Harpenden calipers (Harpenden skinfold calipers, Sweden). Percentage body fat was determined by the four skin-folds method [42].

Exercise tests

During the medical exam, participants performed the Astrand-Ryhming test [43] on a cycle ergometer (Monark Ergoline: ER900, Ergoline, Jaeger, Würzburg, Germany) to estimate maximal oxygen uptake (VO_{2max}). Heart rate (S810, Polar Instruments Inc., Oulu, Finland) and rating of perceived exertion (RPE; Borg 1973) was recorded at the end of each stage. A maximal test was confirmed when participants achieved a minimum of any three of the following criteria; volitional exhaustion, peak heart rate within 10 beats of age predicted maximum, blood lactate above 8 mmol·L⁻¹, final RPE > 18 on Borg scale.

48 hours later (day 2), subjects performed a force-velocity (F/V) test [44] on a cycle ergometer (Monark Ergomedic 894E Peak Bike, Monark, Varberg, Sweden). The test began 5 min after warm-up (15 min at a power output corresponding to 50% estimated VO_{2max}). This test comprised five short trials (6 s) against increasing resistance (2 kg each sprint) until the velocity began to decrease during the 6 s trials. Recovery time between each trial was 5 min. The highest pedaling cadence recorded after each trial was collected from a photoelectric cell fixed on the wheel of the cycle ergometer and connected to a computer. The load that permitted the highest peak power output was used for the Wingate Anaerobic Test (WAnT).

48 hours later (day 3), subjects performed the WAnT on a mechanically-braked Monark cycle ergometer (Monark Ergomedic 894E Peak Bike, Monark, Varberg, Sweden). The test commenced 5 min after warm-up. Subjects were asked to cycle maximally for 30s. Maximal power during the trial was considered as the highest value (W_{peak}), while average power during the WAnT was considered as mean power (W_{mean}).

Before the F/V test and the WAnT, a heart rate monitor (S810, Polar Instruments Inc., Oulu, Finland) was used to control exercise and warm-up intensity. Warm-up intensity was calculated using the re-engineered equation of Swain et al. [45] to determine the maximum heart rate percentage (%MHR) using %VO_{2max}:

$$\%MHR = 0.6463 \times \%VO_{2max} + 37.182$$

Where %MHR is the percentage of maximal heart rate and %VO_{2max} is the percentage of estimated maximal oxygen uptake.

A pilot study was carried out to ensure the reproducibility and sensitivity of W_{peak} and W_{mean} indices, using two measurements of 10 subjects in a single day. Both indices showed excellent intraclass correlation coefficients (ICC = 0.91–0.94), small standard error of measurements (SEM; 3.41–4.56%, <5%) and a small coefficient of variation (CV; <5%).

Similar testing procedures have been used in numerous studies involving normal weight adolescent [46], young adults [47–49], and middle-aged men [32,50–53]. The standard WAnT procedure has limitations because factors such as i) active muscle mass volume and ii) exercise bioenergetics tend to alter maximal power output (P_{max}) (for review see Driss and Vanderwalle [54]. Standardized WAnT intensity has previously failed to utilize an optimal load due to heterogeneous F/V profiles of individuals [44, 55–58]. According to Driss and Vanderwalle [54], the load during a standardized WAnT underestimates P_{max} in normal weight and powerful

adults. Moreover, the load used by researchers at the Wingate Institute was modified several times: $75\text{g}\cdot\text{kg}^{-1}$ body weight[59], then $67\text{g}\cdot\text{kg}^{-1}$ [60], then $75\text{g}\cdot\text{kg}^{-1}$ [61].

Exercise training program

Trained subjects (YT and MAT) underwent 13 weeks'CSRT as previously described [53]. Briefly, CSRT involved three consecutive sessions separated by 48 h: sprint running sessions (13 sessions), resistance training sessions (13 sessions) and sprint cycling sessions (13 sessions). Sessions were performed during the morning and lasted no longer than 70 min, inclusive of 15 min warm-up (jogging and stretching) and 15 min cool-down (jogging and stretching).

During the first training session, YT and MAT performed a sprint running session, which included three to five sets of three to five short bouts at maximum velocity. A passive recovery of 2–3 min was permitted between each set.

Forty-eight hours later, YT and MAT performed the resistance training session, which included five to six exercises targeting all major muscle groups (squat with Smith machine, machine leg extension, machine leg curl, calf raises over a step, triceps pushdown with cable machine, bicep preacher curl, and bench press. The load used during these exercises (% of one-repetition maximum [1-RM]) was progressively increased from 40 to 65% of 1-RM, and increased by 5% of 1-RM per week[62]. To produce maximal power output (i.e., velocity \times load), the positive phase [63] of each exercise was performed as fast as possible [64]. The number of repetitions was maintained as 10–15 per set, and the number of sets increased from three to four over the training period. Therefore, training volume increased progressively during the CSRT program. Rest period between sets were 3–5 min for upper body muscles[65] and at least 1 min for lower limbs to allow for tolerance to increased repetitions. To adjust loads during resistance training sessions, we determined muscle strength using a 1-RM for the six resistance exercises, before CSRT, during the sixth week and post-CSRT. All subjects were familiarized with the test procedures. Ten-minute warm-up (stretching and cycling at 50% $\text{VO}_{2\text{max}}$) preceded the test. After 5 min rest, subjects performed 5 repetitions at approximately 50% of the estimated 1-RM followed by another set of 3 repetitions at 70% of the estimated 1-RM. Subjects then performed one repetition of progressively heavier loads until failure. Maximum strength was determined as the maximum load that could be lifted once with proper technique.

During the third training session, subjects performed a sprint cycling session. Each series comprised three to five repetitions of 10–30 s. The 10–30 s trials were performed maximally. Subjects recovered actively (at a power output corresponding to 50% $\text{VO}_{2\text{max}}$) for 3–5 min between each sprint.

Blood sampling and analysis

Upon the participants' arrival, a heparinized catheter (Insyte-W, 1.1 mm o.d. \times 30 mm) was inserted into an antecubital vein, following sitting for 20 min. Blood was obtained between 08:00 and 09:00 h. to control for diurnal variation on visit two of exercise testing. Venous blood samples were drawn at four times: at rest (t_0 [after 20 min sitting on the bike]), after warm-up (t_w), immediately post-WAnT (t_{end}) and 10 min post-WAnT (t_{10}). For each sample, 10 mL of blood was collected in tubes containing ethylenediaminetetraacetic acid (EDTA) to determine concentrations of serum GH, IGF-1, and IGFBP-3. Samples were centrifuged immediately (at 3000 rpm for 15 min at 4°C), before being divided into appropriate aliquots and stored at -80°C for later analysis.

Blood sample was collected from the finger (20 μL) at the third minute post-WAnT and placed in Eppendorfs for measurement of peak blood lactate concentration ($[\text{La}]_{\text{peak}}$). Blood

lactate concentration was determined using an enzymatic lactate analyzer (Microzym, Cetrax, France).

GH was assayed by chemiluminescence (Immulite, Diagnostic Products Corp., Los Angeles, CA, USA). The GH assay sensitivity limit was $0.1 \text{ ng}\cdot\text{ml}^{-1}$ and inter- and intra-assay CV was 5.7–10% and 4.9–8.3% respectively. IGF-1 was measured using the non-extraction IGF-1 immunoradiometric assay (IRMA) kit (Diagnostic Systems Laboratories, Webster, TX, USA). The theoretical sensitivity, or minimum detection limit, as calculated by interpolation of the mean plus two standard deviations (SD) of 20 standard replicas $0 \text{ ng}\cdot\text{ml}^{-1}$ was $2 \text{ ng}\cdot\text{ml}^{-1}$. The inter-assay CV was 7.4% and 4.2% at concentrations of $35.5 \text{ ng}\cdot\text{ml}^{-1}$ and $383.9 \text{ ng}\cdot\text{ml}^{-1}$ respectively. The intra-assay CV was 6.8% and 6.3% for mean concentrations of $34.03 \text{ ng}\cdot\text{ml}^{-1}$ and $373.86 \text{ ng}\cdot\text{ml}^{-1}$ respectively. IGFBP-3 concentrations were measured by using the non-extraction IRMA kit (Diagnostic Systems Laboratories, Webster, TX, USA). The lower threshold of detection, calculated as the mean two SD of 22 standard replica $0 \text{ ng}\cdot\text{ml}^{-1}$ of IGFBP-3, was $\sim 0.5 \text{ ng}\cdot\text{ml}^{-1}$. The intra-assay CV was 1.8% and 3.9% for mean concentrations of $82.7 \text{ ng}\cdot\text{ml}^{-1}$ and $7.4 \text{ ng}\cdot\text{ml}^{-1}$ respectively. The inter-assay CV was 1.9% and 0.6% for mean concentrations of $76.9 \text{ ng}\cdot\text{ml}^{-1}$ and $8.0 \text{ ng}\cdot\text{ml}^{-1}$ respectively. All biochemical assays were run in duplicate.

Statistical analysis

Data were analyzed using SPSS version 23.0 for Windows (SPSS, Inc. Chicago, IL, USA). Means and standard deviations (SD) were calculated after verifying normality of distributions using the Kolmogorov-Smirnov procedure. For anthropometric, physiological and physical performance, along with area under the curve (AUC), data were analyzed using a multifactorial three-way (time [P1, P2] \times age [young, middle-aged] \times group [trained, control]) ANOVA.

Hormone responses were analyzed using a four-factor ANOVA (time [P1, P2] \times Wingate time [warm-up, immediately post-WAnT and 10 min post-WAnT] \times age [young, middle-aged] \times group [trained, control]).

Total AUCs for GH, IGF-1, and IGFBP-3 were calculated to determine the total hormone exposure over the measured period. It simplified the statistical analyses by rendering the multivariate results into univariate result [66]. The AUC is determined using the linear trapezoidal method which uses linear interpolation between data points.

Greenhouse-Geisser corrections were used when the assumption of sphericity (Mauchly's test) was violated. To help protect against type II errors, an estimate of power (ω) and effect size (η^2_p) were calculated. Bonferroni-adjusted pairwise post hoc comparisons were performed where appropriate. Pearson's product-moment correlation coefficients were calculated to assess relationships between variables. Significance level was set *a priori* at $P < 0.05$.

Results

Anthropometric data

At P1, there was a significant main effect of age for body mass ($F = 4.99$, $P = 0.03$, $\eta^2_p = 0.13$, $\omega = 0.58$), whereby YT and YC were significantly lighter than MAT and MAC ($P < 0.05$). At P2, both training groups experienced a decrease in body mass from P1 (72.4 ± 5.2 and $76.2 \pm 5.4 \text{ kg}$ for YT, and MAT respectively [$P < 0.05$]), whereas the control groups' body mass was not significantly different from P1 ($P > 0.05$). The main effect of time (from P1 to P2) was significant ($F = 8.85$, $P < 0.01$, $\eta^2_p = 0.21$, $\omega = 0.82$) after CSRT.

At P1, there was no main effect of age for body fat percentage ($11.7 \pm 1.4\%$, $11.3 \pm 1.8\%$, $12.9 \pm 1.2\%$, and $12.4 \pm 2.2\%$ for YT, YC, MAT, and MAC respectively; $F = 2.45$, $P = 0.13$, $\eta^2_p = 0.07$, $\omega = 0.33$). At P2, both training groups experienced a decrease in body fat from P1 ($10.2 \pm 0.8\%$ and $11.2 \pm 1.4\%$ for YT, and MAT respectively; $F = 14.52$, $P < 0.01$, $\eta^2_p = 0.30$, $\omega = 0.96$), while

control groups' body fat percentages were not significantly different from P1 ($F = 2.16$, $P > 0.05$). We also observed significant interaction between Time \times Group ($F = 5.82$, $P = 0.02$, $\eta^2_p = 0.15$, $\omega = 0.65$).

At P1, no significant main effect of age for fat-free mass (FFM) was observed (65.2 ± 5.5 kg, 64.1 ± 4.5 kg, 62.1 ± 5.2 kg, and 60.4 ± 3.2 kg for YT, YC, MAT, and MAC respectively with $F = 2.14$, $P = 0.15$, $\eta^2_p = 0.06$, $\omega = 0.30$). At P2, FFM was 66.1 ± 5.1 kg, 64.6 ± 5.7 kg, 63.8 ± 5.4 kg, and 61.4 ± 4.4 kg for YT, YC, MAT, and MAC respectively. No group experienced a significant change in FFM between P1 and P2 ($P > 0.05$) and there was no main effect of time ($F = 0.68$, $P = 0.41$, $\eta^2_p = 0.02$, $\omega = 0.13$).

Physical performance and physiological response

The results of the WAnT are displayed in Table 1. There was a significant main effect of age in W_{peak} ($F = 5.98$, $P = 0.02$, $\eta^2_p = 0.15$, $\omega = 0.66$). At P1, W_{peak} was significantly higher in young groups compared to middle-aged groups ($P < 0.05$). However, this main effect of age for W_{peak} at P1 ($P < 0.001$) was ameliorated at P2 ($P > 0.05$).

In addition, W_{peak} increased significantly after training in both YT and MAT ($P < 0.05$). At P2, trained groups exhibited significantly ($P < 0.05$) higher W_{peak} compared to control groups. Significant interaction between Time \times Group was registered in W_{peak} ($F = 4.92$, $P = 0.03$, $\eta^2_p = 0.13$, $\omega = 0.58$). No main effect of age or time ($P > 0.05$) were observed for W_{mean} .

$[La]_{peak}$ increased significantly ($P < 0.05$) in trained groups (YT and MAT) after CSRT (Table 1). A significant main effect of time was observed ($F = 20.13$, $P < 0.01$, $\eta^2_p = 0.37$, $\omega = 0.99$) for $[La]_{peak}$, whilst there was no main effect of age ($F = 1.38$, $P = 0.25$, $\eta^2_p = 0.04$, $\omega = 0.21$) between groups at P2.

For VO_{2max} , we observed a significant main effect of time ($F = 18.35$, $P < 0.01$, $\eta^2_p = 0.35$, $\omega = 0.99$). Estimated VO_{2max} increased significantly after CSRT in both trained groups ($P < 0.001$), but not in control groups ($P > 0.05$). There was no significant main effect of age ($F = 2.69$, $P = 0.11$, $\eta^2_p = 0.07$, $\omega = 0.36$) in VO_{2max} between groups at P2.

Growth hormone response

There was a significant main effect of age for GH ($F = 17.47$, $P < 0.01$, $\eta^2_p = 0.34$, $\omega = 0.98$) and GH AUC ($F = 18.26$, $P < 0.01$, $\eta^2_p = 0.35$, $\omega = 1.00$) (Table 2).

Table 1. Wingate outcomes and physiological parameters determined before (P1) and after (P2) training.

		YT (n = 10)	YC (n = 9)	MAT (n = 10)	MAC (n = 9)
W_{peak} (W)	P1	1016 \pm 126 ^{a,d}	1000 \pm 312 ^e	885 \pm 155 ^a	887 \pm 102
	P2	1050 \pm 123 ^c	944 \pm 246 ^e	997 \pm 145 ^b	824 \pm 113
W_{mean} (W)	P1	584 \pm 58	500 \pm 93	434 \pm 86	445 \pm 37
	P2	598 \pm 71	473 \pm 80.6	563 \pm 67	402 \pm 80
VO_{2max} (mL \cdot min ⁻¹ \cdot kg ⁻¹)	P1	42.2 \pm 6.1 ^a	43.8 \pm 5.1	39.8 \pm 9.5 ^a	38.5 \pm 3.2
	P2	45.5 \pm 5.7 ^c	42.1 \pm 3.2	45.6 \pm 11.2 ^b	40.1 \pm 3.8
La_{peak} (mmol \cdot l ⁻¹)	P1	14.6 \pm 2.2 ^a	13.9 \pm 3.4	13.4 \pm 2.7 ^a	13.2 \pm 3.2
	P2	16.1 \pm 2.3 ^c	14.3 \pm 3.3	15.1 \pm 2.6 ^b	13.3 \pm 3.1

Data are means \pm SD; maximal Power (W_{peak}); mean power (W_{mean}) in absolute values (W); Peak lactate concentration (La_{peak}); Maximal oxygen uptake (VO_{2max}); young trained (YT); young control (YC); middle-aged trained (MAT); middle-aged control (MAC); before training (P1); after training (P2).

^a significant differences from before and after training, ^a: $P < .05$.

^b significant differences between MAT and MAC, ^b: $P < .05$.

^c Significant differences between YT and YC, ^c: $P < .05$.

^d significant differences between YT and MAT, ^d: $P < .05$.

^e significant differences between YC and MAC, ^e: $P < .05$.

<https://doi.org/10.1371/journal.pone.0183184.t001>

Significantly higher GH was observed at rest, after warm-up, immediately post- and 10 min post-WAnT in the young groups as compared to the middle-aged groups at P1 ($P < 0.05$). This main effect of age was not present in GH_0 or GH_w between MAT and YT at P2 ($P > 0.05$).

There was a main effect of time for GH ($F = 17.83$, $P < 0.01$, $\eta^2_p = 0.34$, $\omega = 0.98$) and GH AUC ($F = 10.83$, $P < 0.01$, $\eta^2_p = 0.34$, $\omega = 0.93$). For YT, GH_0 , GH_{end} , and GH_{10} were significantly higher at P2 compared to P1 ($P < 0.05$). For MAT, GH_0 , GH_{end} , GH_w , and GH_{10} were significantly greater at P2 as compared to P1 ($P < 0.05$). We observed no significant changes from P1 to P2 for GH in control groups ($P > 0.05$).

We observed significant interaction between the Time \times Group ($F = 7.55$, $P = 0.01$, $\eta^2_p = 0.18$, $\omega = 0.76$). GH_0 and GH_{10} (both $P < 0.01$) were significantly higher in YT compared to YC at P2. GH_0 , GH_{end} , and GH_{10} were significantly higher in MAT compared to MAC at P2 ($P < 0.05$).

IGF-1 response

A significant main effect of Wingate-time ($F = 203.19$, $P < 0.01$, $\eta^2_p = 0.86$, $\omega = 1.00$) was observed in all groups (Table 3).

At P1 and P2, IGF-1 increased throughout the progression of the WAnT in all groups ($P < 0.05$), i.e., from IGF-1₀ to IGF-1₁₀. In addition, significant interaction between Wingate-Time \times Age ($F = 16.25$, $P < 0.001$, $\eta^2_p = 0.32$, $\omega = 1.00$) Wingate-Time \times Group ($F = 3.57$, $P = 0.02$, $\eta^2_p = 0.10$, $\omega = 0.78$), Wingate-Time \times Time ($F = 12.82$, $P < 0.001$, $\eta^2_p = 0.27$, $\omega = 1.00$), Wingate-Time \times Time \times Group ($F = 5.00$, $P < 0.001$, $\eta^2_p = 0.13$, $\omega = 0.91$), and Wingate-Time \times Time \times Group \times Age ($F = 3.71$, $P = 0.01$, $\eta^2_p = 0.10$, $\omega = 0.79$) were observed in IGF-1.

We observed a significant main effect of age for IGF-1 ($F = 42.08$, $P < 0.01$, $\eta^2_p = 0.55$, $\omega = 1.00$) and IGF-1 AUC ($F = 41.02$, $P < 0.01$, $\eta^2_p = 0.50$, $\omega = 1.00$). IGF-1 was significantly higher at all phases of the WAnT in young as compared to middle-aged groups at P1 ($P < 0.05$). However, this age effect was not present at P2 ($P > 0.05$ between YT and MAT), whilst, for the control groups, the main effect of age remained statistically significant at P2 ($P < 0.01$). A main effect of time was observed in IGF-1 AUC ($F = 30.42$, $P = 0.01$, $\eta^2_p = 0.17$, $\omega = 1.00$). Significantly greater IGF-1_w and IGF-1_{end} in YT, and IGF-1₁₀, IGF-1_w, and IGF-1_{end} was observed at P2 compared

Table 2. GH concentration (ng·ml⁻¹) determined before (P1) and after (P2) training.

		GH ₀	GH _w	GH _{end}	GH ₁₀	GH AUC
YT (n = 10)	P1	0.33±0.21 ^{a,d}	8.65±2.26 ^d	12.49±4.57 ^{a,d}	12.51±2.55 ^d	256.90±39.70 ^{a,d}
	P2	0.71±0.39 ^c	9.42±4.11	14.46±5.45 ^d	16.74±5.36 ^{a,c,d}	309.91±75.31
YC (n = 9)	P1	0.32±0.30 ^e	8.60±1.44 ^e	12.66±3.79 ^e	13.31±5.22 ^e	262.60±39.22 ^e
	P2	0.45±0.03 ^e	8.83±3.93 ^e	13.10±2.17 ^e	14.33±8.89 ^e	267.20±70.04 ^e
MAT (n = 10)	P1	0.24±0.31 ^a	5.78±3.10 ^a	8.15±4.23 ^a	9.06±3.33 ^a	187.33±36.60 ^a
	P2	0.85±0.94 ^b	8.75±2.97	10.78±1.73 ^b	12.97±2.26 ^b	262.91±46.72 ^b
MAC (n = 9)	P1	0.13±0.01	5.99±3.96	9.02±3.30	9.64±2.52	191.71±65.62
	P2	0.14±0.03	8.15±2.40	10.60±1.75	11.39±1.66	208.50±17.73

Data are means ±SD; plasma GH concentration at rest (GH₀); after warm-up (GH_w); at the end of exercise (GH_{end}); during recovery (GH₁₀); GH AUC: Area under the curve (ng·ml⁻¹ per min), young trained (YT); young control (YC); middle-aged trained (MAT); middle-aged control (MAC); before training (P1); after training (P2) and training

^a significant differences from before and after training, ^a: $P < .05$.

^b significant differences between MAT and MAC, ^b: $P < .05$.

^c Significant differences between YT and YC, ^c: $P < .05$.

^d significant differences between YT and MAT, ^d: $P < .05$.

^e significant differences between YC and MAC, ^e: $P < .05$.

<https://doi.org/10.1371/journal.pone.0183184.t002>

Table 3. IGF-1 concentrations (ng·ml⁻¹) determined before (P1) and after (P2) training.

		IGF-1 ₀	IGF-1 _w	IGF-1 _{end}	IGF-1 ₁₀	IGF-1 AUC
YT (n = 10)	P1	327.60±137.54 ^d	383.80±169.02 ^{a,d}	417.91±172.67 ^{a,d}	678.81±117.33 ^d	13767.70±4425.0 ^{a,d}
	P2	340.90±130.97 ^d	436.79±176.66 ^d	556.63±136.98 ^c	749.87±113.10 ^{c,d}	15886.71±4034.13
YC (n = 9)	P1	340.22±154.45 ^e	323.65±135.19 ^e	433.73±113.92 ^e	465.40±58.12 ^e	13846.01±2541.70 ^e
	P2	361.78±139.05 ^e	353.18±146.30 ^e	443.82±100.05 ^e	528.12±118.73 ^e	12994.21±3763.7 ^e
MAT (n = 10)	P1	157.10±54.0 ^a	223.78±53.25 ^a	292.55±61.62 ^a	300.44±58.11	8379.40±1172.31 ^a
	P2	191.00±66.42	268.11±70.56	458.39±71.78 ^b	392.03±84.64	10403.32±1032.53 ^b
MAC (n = 9)	P1	150.89±87.83	257.58±46.14	358.89±75.70	401.58±78.95	9077.52±1363.22
	P2	153.88±59.25	210.43±44.97	238.29±38.88	331.73±40.52	7951.21±1192.63

Data are means ±SD; plasma IGF-1 concentration at rest (IGF-1₀); after warm-up (IGF-1_w); at the end of exercise (IGF-1_{end}); during recovery (IGF-1₁₀), IGF-1 AUC: Area under the curve (ng·ml⁻¹ per min), young trained (YT); young control (YC); middle-aged trained (MAT); middle-aged control (MAC); before training (P1); after training (P2).

^a significant differences from before and after training, ^a: $P < .05$.

^b significant differences between MAT and MAC, ^b: $P < .05$.

^c Significant differences between YT and YC, ^c: $P < .05$.

^d significant differences between YT and MAT, ^d: $P < .05$.

^e significant differences between YC and MAC, ^e: $P < .05$.

<https://doi.org/10.1371/journal.pone.0183184.t003>

to P1 in MAT ($P < 0.05$). When compared to control groups, significantly greater IGF-1_{end} was observed in MAT compared to MAC at P2 ($P < 0.05$). This last result is confirmed by a significant interaction between Time×Group ($F = 60.90$, $p = 0.00$, $\eta^2_p = 0.64$, $\omega = 1.00$).

IGFBP-3 response

There was a significant main effect of Wingate-Time ($F = 13.50$, $P < 0.01$, $\eta^2_p = 0.28$, $\omega = 1.00$). At P1 and P2, IGFBP-3 increased throughout the progression of the WAnT in all groups ($P < 0.05$), i.e., from IGFBP-3₀ to IGFBP-3₁₀ (Table 4).

Table 4. IGFBP-3 concentrations (ng·ml⁻¹) determined before (P1) and after (P2) training.

		IGFBP-3 ₀	IGFBP-3 _w	IGFBP-3 _{end}	IGFBP-3 ₁₀	IGFBP-3AUC
YT (n = 10)	P1	4072±389 ^d	4019±422	4220±333 ^d	4100±449 ^d	132162±12088
	P2	4163±328 ^c	4211±405	4330±350 ^c	4349±398 ^{c, d}	13174±8902 ^c
YC (n = 9)	P1	3749±403 ^e	3886±358 ^e	4012±384 ^e	3677±373	123202±12087
	P2	3788±349 ^e	3826±305 ^e	4022±335 ^e	3632±348	122455±5192 ^e
MAT (n = 10)	P1	3060±389 ^a	3743±290 ^a	3596±295	3759±372	112767±9871
	P2	4130±416	4010±294	3951±238 ^b	3936±391 ^b	124857±6672 ^b
MAC (n = 9)	P1	3373±403	3613±332	3789±366	3775±383	111005±7022
	P2	3332±379	3648±328	3763±357	3731±375	111860±9854

Data are means ±SD; plasma IGFBP-3 concentration at rest (IGFBP-3₀); after warm-up (IGFBP-3_w); at the end of exercise (IGFBP-3_{end}); during recovery (IGFBP-3₁₀), young trained (YT); IGFBP-3 AUC: Area under the curve (ng·ml⁻¹ per min), young control (YC); middle-aged trained (MAT); middle-aged control (MAC); before training (P1); after training (P2)

^a significant differences from before and after training, ^a: $P < .05$.

^b significant differences between MAT and MAC, ^b: $P < .05$.

^c Significant differences between YT and YC, ^c: $P < .05$.

^d significant differences between YT and MAT, ^d: $P < .05$.

^e significant differences between YC and MAC, ^e: $P < .05$.

<https://doi.org/10.1371/journal.pone.0183184.t004>

There was a significant main effect of age for IGFBP-3 ($F = 25.57$, $P < 0.01$, $\eta^2_p = 0.43$, $\omega = 1.00$). Significantly higher basal IGFBP-3 was observed for YT and YC compared to MAT and MAC at P1 ($P = 0.001$ and $P = 0.002$ respectively). However, this main effect of age was not observed at P2 (i.e., between YT and MAT at P2; $P > 0.05$).

Significant increases in IGFBP-3 were observed in MAT at rest ($P = 0.001$) and during warm-up ($P = 0.003$) at P2 compared to P1. When compared to control groups, IGFBP-3₀, IGFBP-3_{end}, and IGFBP-3₁₀ were significantly higher in trained groups compared to control groups at P2 ($P < 0.05$). In fact, a main effect of group was also observed ($F = 14.12$, $P < 0.01$, $\eta^2_p = 0.29$, $\omega = 0.95$).

IGF-1/IGFBP-3 ratio

The IGF-1/IGFBP-3 ratio was affected by Wingate-time ($F = 130.27$, $P < 0.01$, $\eta^2_p = 0.79$, $\omega = 1.00$). Post-hoc Bonferroni adjustments indicated the ratio increased significantly ($P < 0.05$) from rest to the end of exercise in all groups at P1 and P2. i.e. from IGF-1/IGFBP-3₀ to IGF-1/IGFBP-3₁₀ (Fig 1).

A main effect of age ($F = 25.57$, $P < 0.01$, $\eta^2_p = 0.43$, $\omega = 1.00$) was observed for IGF-1/IGFBP-3 at P1 and was significantly higher at all phases of the WAnT in young compared to middle-aged groups ($P < 0.05$). However, this age effect was not present following CSRT ($P > 0.05$ between YT and MAT), whilst, for control groups, the main effect of age remained statistically significant at P2 ($P < 0.01$). Significant interactions were observed between Age \times Group ($F = 14.29$, $P < 0.01$, $\eta^2_p = 0.30$, $\omega = 1.00$). CSRT induced significant increases in IGF-1/IGFBP-3_w and IGF-1/IGFBP-3_{end} in YT, and increases in IGF-1/IGFBP-3₁₀, IGF-1/IGFBP-3_w and IGF-1/IGFBP-3_{end} in MAT ($P < 0.05$). When compared to control groups, significantly higher IGF-1/IGFBP-3_{end} were observed in MAT as compared to MAC at P2 ($P < 0.05$) with significant interactions observed between Time \times Group ($F = 32.82$, $P < 0.01$, $\eta^2_p = 0.49$ and $\omega = 1.00$).

Correlative analysis

Before training, a significant positive correlation existed between GH₀ and W_{peak} for YT ($r = 0.563$, $P = 0.003$). GH_{end} was also positively correlated with W_{peak} at P1 ($r = 0.792$, $P = 0.001$) and P2 ($r = 0.710$, $P < 0.001$) in MAT, but only at P2 in YT ($r = 0.268$, $P = 0.002$).

Discussion

The primary finding of the present investigation is that 13 weeks' CSRT induced a significant increase in W_{peak} in both YT and MAT which ameliorated the effect of age between these groups. Increased sprint performance was associated with increased GH at rest, in response to the WAnT, and during recovery in YT and MAT, whilst IGF-1 and IGFBP-3 increased at rest in MAT only. Prior to CSRT, an effect of age was observed in GH, IGF-1, IGFBP-3, and IGF-1/IGFBP-3 basally and in response to exercise. However, CSRT ameliorated this age effect as both training groups (MAT and YT) had comparable basal GH and GH AUC, IGF-1_{end} and IGF-1 AUC, and IGF-1/IGFBP-3.

The results of the AUC analysis for GH and IGF-1 suggest CSRT produce prolonged elevations in these hormones in response to the WAnT. Moreover, the increased IGF-1/IGFBP-3 ratio in response to exercise in trained groups suggests that there may be more unbound IGF-1 available for hormone-receptor interactions.

GH increases are associated with improvements in lean body mass and reduced fat mass [67] which was supported by the present investigation. In the current study, trained groups exhibited a trend for greater FFM at P2 (~0.9–1.7kg increases) and significantly less fat mass




Fig 1. [IGF-1]/[IGFBP-3] ratio changes determined before (P1; above) and after (P2; below) training. Plasma [IGF-1]/[IGFBP-3] ratio at rest (₀), after warm-up (_w), at the end of exercise (_{end}), and during recovery (₁₀) in young trained (YT), young control (YC), middle-aged trained (MAT), and middle-aged control (MAC) groups. Data are presented as mean \pm standard error for clarity.

<https://doi.org/10.1371/journal.pone.0183184.g001>

which may increase oxidative capacity as more active tissue would permit a greater oxygen uptake [68]. Our observation of increased estimated $\text{VO}_{2\text{max}}$ of ~16% following CSRT is in line with previous investigators who report increased cardiorespiratory fitness following high intensity training [69, 70]. In their recent review, Sloth et al. [71] suggested short term sprint interval training performed alone induces increases of 4.2–13.4% $\text{VO}_{2\text{max}}$ in young adult males. Therefore, the combination of sprint interval training with strength exercise appears a pragmatic approach to improve aerobic capacity in young and middle-aged subjects. These adaptations may be explained by increased enzymatic activity and mitochondrial mass [71–73]. However, a mechanistic explanation is outside the scope of this investigation.

As increased aerobic capacity is associated with increased blood lactate during intense efforts [54, 74], the higher blood lactate observed in trained groups may be due to increased ATP hydrolysis leading to greater lactate efflux in the plasma compartment as supported previously [53, 75].

CSRT produced a significant increase in W_{peak} in trained groups, which supports several studies that report anaerobic training leading to enhanced power production in young [76] and older [77, 78] subjects. Interestingly, we found a positive correlation between serum growth factors and physical performance alterations, providing further associative evidence linking growth factors to improved muscle power [79].

In the present study, training-induced increases in basal GH occurred between trained groups, and ameliorated the age effect at P2. Such findings are in accordance with recent investigations demonstrating that intense training increases GH in males [80, 81]. However, other studies have not observed perturbations to basal GH after sprint [34, 82, 83] or resistance [83] training performed alone. The relatively short half-life of GH and the “snapshot” nature of basal hormone sampling may explain discrepancies in results as GH during rest loses half of its physiologic activity in 10–30 minutes following secretion [84]. This brief period of hormonal activity depends on several factors such as diet, sleep duration, blood sampling (stress response from the vascular puncture) and time of the day [20, 84] which contributes to fluctuations in basal GH. The most reproducible pulse of GH secretion however, occurs in response to intense exercise, when rate of secretion exceeds rate of degradation [85]. For example, Nevill et al. [34] reported GH to be tenfold higher than basal levels in sprinters following exercise and were greater than those measured in endurance athletes. GH increases may also be exercise volume dependent as 30 s all-out sprints resulted in greater GH response than 6 s sprints and remained high for >90 minutes compared to 60 min following 6 s sprint [86]. An alternative explanation for discrepancies with previous investigations is the utilisation of concurrent training in the present study, rather than only resistance or only sprint training previously investigated.

GH anabolic actions are mostly mediated through IGF-1 and include regulation of whole body protein synthesis and breakdown [20]. Interestingly, the increased GH AUC following CSRT was associated with increase in IGF-1 AUC in YT and MAT suggesting that CSRT is a potent stimulus of the GH/IGF-1 axis. Data from the present investigation are in agreement with some [87] but not all [88] previous investigations in reporting increased basal IGF-1 following exercise training. For example, Cooper, Taaffe (76) reported comparable IGF-1 in masters runners and minimally exercising men aged 60–70 yrs ($129 \pm 10 \text{ ng} \cdot \text{ml}^{-1}$ and $124 \pm 11 \text{ ng} \cdot \text{ml}^{-1}$, respectively). Differences in exercise modality and participant characteristics may explain discrepancies in findings, as the present investigation studied the influence of CSRT in young and middle-aged, rather than older, men as in the aforementioned investigation. Whilst increased systemic IGF-1 is generally considered to confer an anabolic advantage, a recent investigation [89] reported decreased systemic IGF-1 following 12 weeks’ resistance exercise in older adults (74 ± 6 yrs), despite increased lean mass. These authors concluded that, during periods of active muscle building, IGF-1 is redistributed from circulation into tissue. Moreover, increased IGF-1 was observed without significant alteration in IGFBP-3 AUC in MAT following CSRT. IGFBP-

3 is known to enhance IGF-1 retention in the circulation and block its access to the cognate receptor (IGF-1R)[15, 16]. The effect of intense training on IGFBP3 in elderly is still conflicting [90], and further investigation in large population is needed to better understand its variability over time in response to intense training.

There is sufficient evidence that anabolism in the elderly are intimately tied to alterations in GH/IGF-1 axis and declines in total and free IGF-1 response [91]. Nevertheless, CSRT appears to reduce this age-related effect on IGF-1/IGFBP-3 (a surrogate for IGF-1 bioavailability) between trained groups at rest, at the end of WAnT, and during recovery with important increases at all phases following the WAnT. IGF-1 bioavailability appears a strong predictor on the effectiveness of the IGF-1 action [92] and its increase confirms improvements of GH/IGF-1 axis in young and middle-aged trained men with training.

Conclusion

In conclusion, 13 weeks' CSRT improved peak power output and body composition in young and middle-aged military men. This training intervention increases circulating GH in trained groups at rest and in response to exercise. Training-induced increases in basal GH occurred with amelioration of the age effect between trained groups. GH improvements in middle-aged men were associated with increased IGF-1 levels following CSRT. These were associated with increased IGF-1 bioavailability index in young and middle-aged trained groups following training. As such, CSRT appears to counteract the aging effect of somatotrophic hormones.

Acknowledgments

The authors thank all the experts, doctors and officers of the Military Hospital of Instruction of Tunis and the General Direction of the Military Health for their participation in this experimental work.

Author Contributions

Conceptualization: MS.

Data curation: MS HD AC.

Formal analysis: WD FR.

Funding acquisition: MS.

Investigation: MS HD AC.

Methodology: MS AC.

Project administration: MS AC.

Resources: MS HD.

Software: MS WD HD.

Supervision: MS AC.

Validation: MS WD LDH JP FR HD AC.

Visualization: MS WD LDH JP FR HD AC.

Writing – original draft: MS LDH.

Writing – review & editing: MS WD LDH JP FR HD AC.

References

1. Frontera WR, Hughes VA, Lutz KJ, Evans WJ. A cross-sectional study of muscle strength and mass in 45-to 78-yr-old men and women. *Journal of applied physiology*. 1991; 71(2):644–50. PMID: [1938738](#)
2. Fleg JL, Lakatta EG. Role of muscle loss in the age-associated reduction in VO2 max. *Journal of applied physiology*. 1988; 65(3):1147–51. PMID: [3182484](#)
3. Korhonen MT, Mero A, Suominen H. Age-related differences in 100-m sprint performance in male and female master runners. *Medicine and science in sports and exercise*. 2003; 35(8):1419–28. <https://doi.org/10.1249/01.MSS.0000079080.15333.CA> PMID: [12900699](#)
4. Gent DN, Norton K. Aging has greater impact on anaerobic versus aerobic power in trained masters athletes. *Journal of sports sciences*. 2013; 31(1):97–103. <https://doi.org/10.1080/02640414.2012.721561> PMID: [22974317](#)
5. Sipilä S, Narici M, Kjaer M, Pollanen E, Atkinson RA, Hansen M, et al. Sex hormones and skeletal muscle weakness. *Biogerontology*. 2013; 14(3):231–45. Epub 2013/05/03. <https://doi.org/10.1007/s10522-013-9425-8> PMID: [23636830](#)
6. Hayes LD, Sculthorpe N, Herbert P, Baker JS, Spagna R, Grace FM. Six weeks of conditioning exercise increases total, but not free testosterone in lifelong sedentary aging men. *The Aging Male*. 2015; 18(3):195–200. <https://doi.org/10.3109/13685538.2015.1046123> PMID: [26030347](#)
7. Hermann M, Berger P. Hormonal changes in aging men: a therapeutic indication? *Experimental gerontology*. 2001; 36(7):1075–82. PMID: [11404052](#)
8. Corpas E, Harman SM, Blackman MR. Human growth hormone and human aging. *Endocrine reviews*. 1993; 14(1):20–39. <https://doi.org/10.1210/edrv-14-1-20> PMID: [8491152](#)
9. Kim T, Chang JS, Kim H, Lee KH, Kong ID. Intense Walking Exercise Affects Serum IGF-1 and IGFBP3. *Journal of lifestyle medicine*. 2015; 5(1):21. <https://doi.org/10.15280/jlm.2015.5.1.21> PMID: [26528426](#)
10. RUTANEN E-M, PEKONEN F, MÄKINEN T. Soluble 34K binding protein inhibits the binding of insulin-like growth factor I to its cell receptors in human secretory phase endometrium: evidence for autocrine/paracrine regulation of growth factor action. *The Journal of Clinical Endocrinology & Metabolism*. 1988; 66(1):173–80.
11. Giovannini S, Cesari M, Marzetti E, Leeuwenburgh C, Maggio M, Pahor M. Effects of ACE-inhibition on IGF-1 and IGFBP-3 concentrations in older adults with high cardiovascular risk profile. *The journal of nutrition, health & aging*. 2010; 14(6):457–60.
12. Baxter RC, Martin JL. Radioimmunoassay of growth hormone-dependent insulinlike growth factor binding protein in human plasma. *Journal of Clinical Investigation*. 1986; 78(6):1504. <https://doi.org/10.1172/JCI112742> PMID: [2431001](#)
13. Olivecrona H, Hilding A, Ekström C, Barle H, Nyberg Br, Möller C, et al. Acute and Short-Term Effects of Growth Hormone on Insulin-Like Growth Factors and Their Binding Proteins: Serum Levels and Hepatic Messenger Ribonucleic Acid Responses in Humans 1. *The Journal of Clinical Endocrinology & Metabolism*. 1999; 84(2):553–60.
14. Ho K, Hoffman D. Aging and growth hormone. *Hormone Research in Paediatrics*. 1993; 40(1–3):80–6.
15. Hwa V, Oh Y, Rosenfeld RG. The insulin-like growth factor-binding protein (IGFBP) superfamily 1. *Endocrine reviews*. 1999; 20(6):761–87. <https://doi.org/10.1210/edrv.20.6.0382> PMID: [10605625](#)
16. Naspi A, Panasiti V, Abbate F, Roberti V, Devirgiliis V, Curzio M, et al. Insulin-like-growth-factor-binding-protein-3 (IGFBP-3) contrasts melanoma progression in vitro and in vivo. *PloS one*. 2014; 9(6):e98641. <https://doi.org/10.1371/journal.pone.0098641> PMID: [24905466](#)
17. Codner E, Mericq V, Ugarte F, Iniguez G, Espinoza M, Avila A, et al. Usefulness of the measurement of insulin-like growth factor (IGF-I) and IGF-1 binding protein-3 (IGFBP-3) for the diagnosis of growth hormone (GH) deficiency in children. *Revista medica de Chile*. 1999; 127(7):807–13. PMID: [10668288](#)
18. Birzniece V, Nelson AE, Ho KK. Growth hormone and physical performance. *Trends in Endocrinology & Metabolism*. 2011; 22(5):171–8.
19. Blackman MR, Sorkin JD, Münzer T, Bellantoni MF, Busby-Whitehead J, Stevens TE, et al. Growth hormone and sex steroid administration in healthy aged women and men: a randomized controlled trial. *Jama*. 2002; 288(18):2282–92. PMID: [12425705](#)
20. Velloso C. Regulation of muscle mass by growth hormone and IGF-I. *British journal of pharmacology*. 2008; 154(3):557–68. <https://doi.org/10.1038/bjpp.2008.153> PMID: [18500379](#)
21. Cappon J, Brasel J, Mohan S, Cooper D. Effect of brief exercise on circulating insulin-like growth factor I. *Journal of applied physiology*. 1994; 76(6):2490–6. PMID: [7928875](#)
22. Manetta J, Brun JF, Maimoun L, Callis A, Prefaut C, Mercier J. Effect of training on the GH/IGF-I axis during exercise in middle-aged men: relationship to glucose homeostasis. *American Journal of*

- Physiology-Endocrinology and Metabolism. 2002; 283(5):E929–E36. <https://doi.org/10.1152/ajpendo.00539.2001> PMID: 12376319
23. Wideman L, Weltman JY, Hartman ML, Veldhuis JD, Weltman A. Growth hormone release during acute and chronic aerobic and resistance exercise. *Sports medicine*. 2002; 32(15):987–1004. PMID: 12457419
24. Godfrey RJ, Madgwick Z, Whyte GP. The exercise-induced growth hormone response in athletes. *Sports medicine*. 2003; 33(8):599–613. PMID: 12797841
25. Elio F, Antonelli G, Gatti R, Chiappin S, Spinella P, Cappellin E. Effects of two different types of exercise on GH/IGF axis in athletes. Is the free/total IGF-I ratio a new investigative approach? *Clinica Chimica Acta*. 2008; 387(1):71–4.
26. Frystyk J. Exercise and the growth hormone-insulin-like growth factor axis. *Medicine and science in sports and exercise*. 2010; 42(1):58–66. <https://doi.org/10.1249/MSS.0b013e3181b07d2d> PMID: 20010129
27. Salvadori A, Fanari P, Marzullo P, Codecasa F, Tovaglieri I, Cornacchia M, et al. Dynamics of GH secretion during incremental exercise in obesity, before and after a short period of training at different work-loads. *Clinical endocrinology*. 2010; 73(4):491–6. <https://doi.org/10.1111/j.1365-2265.2010.03837.x> PMID: 20550528
28. Allen DB. Growth suppression by glucocorticoid therapy. *Endocrinology and metabolism clinics of North America*. 1996; 25(3):699–717. PMID: 8879994
29. Meinhardt UJ, Ho KK. Modulation of growth hormone action by sex steroids. *Clinical endocrinology*. 2006; 65(4):413–22. <https://doi.org/10.1111/j.1365-2265.2006.02676.x> PMID: 16984231
30. Mazziotti G, Giustina A. Glucocorticoids and the regulation of growth hormone secretion. *Nature Reviews Endocrinology*. 2013; 9(5):265–76. <https://doi.org/10.1038/nrendo.2013.5> PMID: 23381030
31. Zouhal H, Jacob C, Delamarche P, Gratas-Delamarche A. Catecholamines and the effects of exercise, training and gender. *Sports medicine*. 2008; 38(5):401–23. PMID: 18416594
32. Zouhal H, Vincent S, Moussa E, Botcazou M, Delamarche P, Gratas-Delamarche A. Early advancing age alters plasma glucose and glucoregulatory hormones in response to supramaximal exercise. *Journal of Science and Medicine in Sport*. 2009; 12(6):652–6. <https://doi.org/10.1016/j.jsams.2008.03.003> PMID: 18547864
33. Sellami M, Ben Abderrahman A, Kebisi W, De Sousa MV, Zouhal H. Effect of sprint and strength training on glucoregulatory hormones: Effect of advanced age. *Experimental Biology and Medicine*. 2017; 242(1):113–23. <https://doi.org/10.1177/1535370216662711> PMID: 27470931
34. Nevill ME, Holmyard DJ, Hall GM, Allsop P, van Oosterhout A, Burrin JM, et al. Growth hormone responses to treadmill sprinting in sprint- and endurance-trained athletes. *European journal of applied physiology and occupational physiology*. 1996; 72(5–6):460–7. PMID: 8925817.
35. Poehlman ET, Rosen CJ, Copeland KC. The influence of endurance training on insulin-like growth factor-1 in older individuals. *Metabolism*. 1994; 43(11):1401–5. PMID: 7526125
36. Vitiello MV, Wilkinson CW, Merriam GR, Moe KE, Prinz PN, Ralph DD, et al. Successful 6-month endurance training does not alter insulin-like growth factor-I in healthy older men and women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 1997; 52(3):M149–M54.
37. Borst SE, De Hoyos DV, Garzarella L, Vincent K, Pollock BH, Lowenthal DT, et al. Effects of resistance training on insulin-like growth factor-I and IGF binding proteins. *Medicine and science in sports and exercise*. 2001; 33(4):648–53. PMID: 11283443
38. Adams K, O'shea JP, O'shea KL, Climstein M. The effect of six weeks of squat, plyometric and squat-plyometric training on power production. *The Journal of strength & conditioning research*. 1992; 6(1):36–41.
39. Reaburn P, Logan P, Mackinnon L. Serum Testosterone Response to High-Intensity Resistance Training in Male Veteran Sprint Runners. *The Journal of Strength & Conditioning Research*. 1997; 11(4):256–60.
40. Kotzamanidis C, Chatzopoulos D, Michailidis C, Papaiakevou G, Patikas D. The effect of a combined high-intensity strength and speed training program on the running and jumping ability of soccer players. *The Journal of Strength & Conditioning Research*. 2005; 19(2):369–75.
41. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *The American journal of clinical nutrition*. 1982; 36(5):936–42. PMID: 7137077.
42. Durnin JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *British journal of nutrition*. 1974; 32(01):77–97.

43. Åstrand P-O, Ryhming I. A nomogram for calculation of aerobic capacity (physical fitness) from pulse rate during submaximal work. *Journal of applied physiology*. 1954; 7(2):218–21. PMID: [13211501](#)
44. Vandewalle H, Peres G, Heller J, Monod H. All out anaerobic capacity tests on cycle ergometers. *European journal of applied physiology and occupational physiology*. 1985; 54(2):222–9. PMID: [4043052](#)
45. Swain DP, Abernathy KS, Smith CS, Lee SJ, Bunn SA. Target heart rates for the development of cardio-respiratory fitness. *Medicine and science in sports and exercise*. 1994; 26(1):112–6. PMID: [8133731](#)
46. Botcazou M, Jacob C, Gratas-Delamarche A, Vincent S, Bentué-Ferrer D, Delamarche P, et al. Sex effect on catecholamine responses to sprint exercise in adolescents and adults. *Pediatric exercise science*. 2007; 19(2):132–44. PMID: [17603137](#)
47. Moussa E, Zouhal H, Vincent S, Prioux J. Effect of sprint duration (6 s or 30 s) on plasma glucose regulation in untrained male subjects. *Journal of sports medicine and physical fitness*. 2003; 43(4):546. PMID: [14767419](#)
48. Jacob C, Zouhal H, Prioux J, Gratas-Delamarche A, Bentue-Ferrer D, Delamarche P. Effect of the intensity of training on catecholamine responses to supramaximal exercise in endurance-trained men. *European journal of applied physiology*. 2004; 91(1):35–40. <https://doi.org/10.1007/s00421-003-1002-4> PMID: [14614627](#)
49. Derbré F, Vincent S, Maitel B, Jacob C, Delamarche P, Delamarche A, et al. Androgen responses to sprint exercise in young men. *International journal of sports medicine*. 2010; 31(05):291–7.
50. Zouhal H, Gratas-Delamarche A, Bentué-Ferrer D, Rannou F, Granier P, Delamarche P. Plasma catecholamine responses during a supramaximal exercise in endurance trained. Sex differences? *Science & Sports*. 1998; 3(13):112–8.
51. Zouhal H, Gratas-Delamarche A, Rannou F, Granier P, Bentue-Ferrer D, Delamarche P. Between 21 and 34 years of age, aging alters the catecholamine responses to supramaximal exercise in endurance trained athletes. *International journal of sports medicine*. 1999; 20(06):343–8.
52. Zouhal H, Jacob C, Rannou F, Gratas-Delamarche A. Effect of training status on the sympathoadrenal activity during a supramaximal exercise in human. *Journal of sports medicine and physical fitness*. 2001; 41(3):330. PMID: [11533563](#)
53. Sellami M, Abderrahman AB, Casazza GA, Kebisi W, Lemoine-Morel S, Bouguerra L, et al. Effect of age and combined sprint and strength training on plasma catecholamine responses to a Wingate-test. *European journal of applied physiology*. 2014; 114(5):969–82. <https://doi.org/10.1007/s00421-014-2828-7> PMID: [24500570](#).
54. Driss T, Vandewalle H. The measurement of maximal (anaerobic) power output on a cycle ergometer: a critical review. *BioMed research international*. 2013; 2013.
55. Dotan R, Bar-Or O. Load optimization for the Wingate anaerobic test. *European journal of applied physiology and occupational physiology*. 1983; 51(3):409–17. PMID: [6685039](#)
56. Patton JF, Murphy MM, Frederick F. Maximal power outputs during the Wingate anaerobic test. *International journal of sports medicine*. 1985; 6(02):82–5.
57. Winter EM, Jones AM, Davison RR, Bromley PD, Mercer TH. *Sport and Exercise Physiology Testing Guidelines: Volume I—Sport Testing: The British Association of Sport and Exercise Sciences Guide*: Routledge; 2006.
58. Jaafar H, Rouis M, Attiogbé E, Vandewalle H, Driss T. A Comparative Study between the Wingate and Force–Velocity Anaerobic Cycling Tests: Effect of Physical Fitness. *International journal of sports physiology and performance*. 2016; 11(1):48–54. <https://doi.org/10.1123/ijsp.2015-0063> PMID: [25849068](#)
59. Ayalon A, Inbar O, Bar-Or O. Relationships among measurements of explosive strength and anaerobic power. *Biomechanics IV*: Springer; 1974. p. 572–7.
60. Dotan R, Bar-Or O. Climatic heat stress and performance in the Wingate Anaerobic Test. *European journal of applied physiology and occupational physiology*. 1980; 44(3):237–43. PMID: [7190918](#)
61. Inbar O, Bar-Or O. Anaerobic characteristics in male children and adolescents. *Medicine and science in sports and exercise*. 1986; 18(3):264–9. PMID: [3523102](#)
62. Padulo J, Laffaye G, Chaouachi A, Chamari K. Bench press exercise: the key points. *The Journal of sports medicine and physical fitness*. 2015; 55(6):604–8. PMID: [24823345](#).
63. Padulo J, Laffaye G, Ardigò LP, Chamari K. Concentric and eccentric: muscle contraction or exercise? *Journal of human kinetics*. 2013; 37(1):5–6.
64. Padulo J, Mignogna P, Mignardi S, Tonni F, D'ottavio S. Effect of different pushing speeds on bench press. *International journal of sports medicine*. 2012; 33(05):376–80.
65. de Salles BF, Simao R, Miranda F, Novaes Jda S, Lemos A, Willardson JM. Rest interval between sets in strength training. *Sports medicine*. 2009; 39(9):765–77. <https://doi.org/10.2165/11315230-000000000-00000> PMID: [19691365](#).

66. Fekedulegn DB, Andrew ME, Burchfiel CM, Violanti JM, Hartley TA, Charles LE, et al. Area under the curve and other summary indicators of repeated waking cortisol measurements. *Psychosomatic medicine*. 2007; 69(7):651–9. <https://doi.org/10.1097/PSY.0b013e31814c405c> PMID: 17766693
67. Cuneo RC, Salomon F, Wiles CM, Hesp R, Sonksen PH. Growth hormone treatment in growth hormone-deficient adults. II. Effects on exercise performance. *Journal of applied physiology*. 1991; 70(2):695–700. PMID: 2022561.
68. Jurimae J, Tillmann V, Purge P, Jurimae T. Body composition, maximal aerobic performance and inflammatory biomarkers in endurance-trained athletes. *Clinical physiology and functional imaging*. 2015. <https://doi.org/10.1111/cpf.12299> PMID: 26373614.
69. Gibala MJ, Little JP, Van Essen M, Wilkin GP, Burgomaster KA, Safdar A, et al. Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *The Journal of physiology*. 2006; 575(3):901–11.
70. Ozaki H, Loenneke JP, Thiebaud RS, Abe T. Resistance training induced increase in VO₂ max in young and older subjects. *European Review of Aging and Physical Activity*. 2013; 10(2):107.
71. Sloth M, Sloth D, Overgaard K, Dalgas U. Effects of sprint interval training on VO₂max and aerobic exercise performance: a systematic review and meta-analysis. *Scandinavian journal of medicine & science in sports*. 2013; 23(6):e341–e52.
72. Burgomaster KA, Heigenhauser GJ, Gibala MJ. Effect of short-term sprint interval training on human skeletal muscle carbohydrate metabolism during exercise and time-trial performance. *Journal of applied physiology*. 2006; 100(6):2041–7. <https://doi.org/10.1152/jappphysiol.01220.2005> PMID: 16469933
73. Burgomaster KA, Hughes SC, Heigenhauser GJ, Bradwell SN, Gibala MJ. Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans. *Journal of applied physiology*. 2005; 98(6):1985–90. <https://doi.org/10.1152/jappphysiol.01095.2004> PMID: 15705728
74. Sellami M, Chamari K, Zagatto AM, Keksi W, Chaouachi A, Zouhal H. Racial differences in hemoglobin and plasma volume variation: implications for muscle performance and recovery. *Ethnicity & Health*. 2017:1–12.
75. Maharam LG, Bauman PA, Kalman D, Skolnik H, Perle SM. Masters Athletes. *Sports medicine*. 1999; 28(4):273–85. PMID: 10565553
76. Cooper CS, Taaffe DR, Guido D, Packer E, Holloway L, Marcus R. Relationship of chronic endurance exercise to the somatotrophic and sex hormone status of older men. *European journal of endocrinology / European Federation of Endocrine Societies*. 1998; 138(5):517–23. PMID: 9625362.
77. Sculthorpe N, Herbert P, Grace FM. Low-Frequency High-Intensity Interval Training is an Effective Method to Improve Muscle Power in Lifelong Sedentary Aging Men: A Randomized Controlled Trial. *Journal of the American Geriatrics Society*. 2015; 63(11):2412–3. Epub 2015/11/26. <https://doi.org/10.1111/jgs.13863> PMID: 26603065.
78. Knowles AM, Herbert P, Easton C, Sculthorpe N, Grace FM. Impact of low-volume, high-intensity interval training on maximal aerobic capacity, health-related quality of life and motivation to exercise in ageing men. *Age (Dordrecht, Netherlands)*. 2015; 37(2):25. Epub 2015/03/17. <https://doi.org/10.1007/s11357-015-9763-3> PMID: 25773069; PubMed Central PMCID: PMC4359174.
79. Chikani V, Cuneo RC, Hickman I, Ho KK. Growth hormone (GH) enhances anaerobic capacity: impact on physical function and quality of life in adults with GH deficiency. *Clinical endocrinology*. 2016. <https://doi.org/10.1111/cen.13147> PMID: 27346880.
80. Gonzalez AM, Hoffman JR, Townsend JR, Jajtner AR, Boone CH, Beyer KS, et al. Intramuscular anabolic signaling and endocrine response following high volume and high intensity resistance exercise protocols in trained men. *Physiological reports*. 2015; 3(7). Epub 2015/07/23. <https://doi.org/10.14814/phy2.12466> PMID: 26197935; PubMed Central PMCID: PMC4552541.
81. Eliakim A, Brasel J, Mohan S, Barstow T, Berman N, Cooper D. Physical fitness, endurance training, and the growth hormone-insulin-like growth factor I system in adolescent females. *The Journal of Clinical Endocrinology & Metabolism*. 1996; 81(11):3986–92.
82. Stokes KA, Nevill ME, Cherry PW, Lakomy HK, Hall GM. Effect of 6 weeks of sprint training on growth hormone responses to sprinting. *European journal of applied physiology*. 2004; 92(1–2):26–32. <https://doi.org/10.1007/s00421-003-1038-5> PMID: 14985991.
83. McCall GE, Byrnes WC, Fleck SJ, Dickinson A, Kraemer WJ. Acute and chronic hormonal responses to resistance training designed to promote muscle hypertrophy. *Canadian journal of applied physiology = Revue canadienne de physiologie appliquee*. 1999; 24(1):96–107. PMID: 9916184.
84. Brandt T. *Neurological disorders: course and treatment*: Gulf Professional Publishing; 2003.
85. Harvey S, Scanes CG, Daughaday WH. *Growth hormone*: CRC press; 1994.

86. Stokes K, Nevill M, Hall G, Lakomy H. The time course of the human growth hormone response to a 6 s and a 30 s cycle ergometer sprint. *Journal of sports sciences*. 2002; 20(6):487–94. <https://doi.org/10.1080/02640410252925152> PMID: 12137178
87. Tsai C-L, Wang C-H, Pan C-Y, Chen F-C. The effects of long-term resistance exercise on the relationship between neurocognitive performance and GH, IGF-1, and homocysteine levels in the elderly. *Frontiers in behavioral neuroscience*. 2015;9.
88. Wieczorek-Baranowska A, Nowak A, Michalak E, Karolkiewicz J, Pospieszna B, Rutkowski R, et al. Effect of aerobic exercise on insulin, insulin-like growth factor-1 and insulin-like growth factor binding protein-3 in overweight and obese postmenopausal women. *The Journal of sports medicine and physical fitness*. 2011; 51(3):525–32. Epub 2011/09/10. PMID: 21904293.
89. Arnarson A, Gudny Geirsdottir O, Ramel A, Jonsson PV, Thorsdottir I. Insulin-Like Growth Factor-1 and Resistance Exercise in Community Dwelling Old Adults. *The journal of nutrition, health & aging*. 2015; 19(8):856–60. Epub 2015/09/29. <https://doi.org/10.1007/s12603-015-0547-3> PMID: 26412290.
90. HELLÉNIUS MLB, BRISMAR KE, BERGLUND BH, FAIRE UH. Effects on glucose tolerance, insulin secretion, insulin-like growth factor 1 and its binding protein, IGFBP-1, in a randomized controlled diet and exercise study in healthy, middle-aged men. *Journal of internal medicine*. 1995; 238(2):121–30. PMID: 7543131
91. Eliakim A, Nemet D, Cooper DM. Exercise, training, and the GH-IGF-I axis. *The Endocrine System in Sports and Exercise* Malden (MA): Blackwell Publishing. 2005:165–79.
92. Kucera R, Topolcan O, Pecen L, Kinkorova J, Svobodova S, Windrichova J, et al. Reference values of IGF1, IGFBP3 and IGF1/IGFBP3 ratio in adult population in the Czech Republic. *Clinica Chimica Acta*. 2015; 444:271–7.