

Cammack, Nicholas and Howe, Louis ORCID: https://orcid.org/0000-0003-2001-2802 (2019) Ankle dorsiflexion range of motion impacts sagittal but not frontalplane landing mechanics. In: British Association of Sport and Exercise Sciences (BASES) Student Conference 2019: Sport and Exercise Science - Transforming Lives, 17-18 April 2019, University of Dundee and Abertay University, UK. (Unpublished)

Downloaded from: https://insight.cumbria.ac.uk/id/eprint/4803/

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available <u>here</u>) for educational and not-for-profit activities

### provided that

• the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form

• a hyperlink/URL to the original Insight record of that item is included in any citations of the work

• the content is not changed in any way

• all files required for usage of the item are kept together with the main item file.

### You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found <u>here</u>. Alternatively contact the University of Cumbria Repository Editor by emailing <u>insight@cumbria.ac.uk</u>.

# Ankle dorsiflexion range of motion impacts sagittal-plane, but not frontal-plane landing mechanics

Cammack, N. A.<sup>1</sup> & Howe, L. P.<sup>1</sup>

### Introduction

Poor landing mechanics has been highlighted as injury risk factor for the development of both acute and chronic injuries (Shimokochi et al., 2009). Deficits in ankle dorsiflexion range of motion (DF ROM) have been shown to lead to altered landing mechanics in the lower extremities. However, at present there is limited evidence for restrictions in ankle DF ROM influencing hip or trunk kinematics during landing tasks.

### Aims

The aims of study investigated the relationship between ankle DF ROM and the aberrant strategies in lower extremity and trunk kinematics during a single leg drop vertical jump (SLDVJ).

# Methods

Thirty-one participants volunteered for this investigation (12) women and 19 men, mean ± SD; age 21 ± 1.7 years, height 173.1 ± 8.2 cm, mass 72.3 ± 13.7). Ankle DF ROM was measured for the right limb using the weight bearing lunge test (WBLT). With institutional ethical approval, a single-leg drop vertical jump was performed off a 10 cm box on the right limb for three repetitions. Using 2d motion analysis, sagittal-plane initial contact and peak flexion angles were calculated for the hip, knee and ankle joints. Frontal-plane projection angles (FPPA) for the knee and the lateral trunk angle were also calculated at the moment of peak knee flexion. Pearson correlation tests were used to identify if a relationship existed between ankle DF ROM and the landing mechanics in both planes.

References: Dowling, B., Mcpherson, A.L. and Paci, J.M. (2018) 'Weightbearing ankle dorsiflexion range of motion and sagittal plane kinematics during single leg drop jump landing in healthy male athletes.', The Journal of sports medicine and physical fitness, 58(6), pp. 867-874. Shinokochi, Y., Yong Lee, S., Shultz, R.J. (2009) 'The relationships among sagittal-plane lower extremity moments: implications for landing strategy in anterior cruciate ligament injury prevention', Journal of Athletic Training, 44(1), pp. 33-38.



17-18 April University of Dundee and Abertay University



**Figure 1**: Peak flexion frontal-plane projection angles during SLDVJ.



Figure 2: Peak flexion in sagittal-plane during SLDVJ



www.cumbria.ac.uk s1603947@cumbria.ac.uk



Figure 3: Initial contact angle in sagittal-plane during SLDVJ

### Results

A significant negative relationship was found between ankle DF ROM and peak flexion at the knee (R = -0.53, P < 0.002) and ankle (R = -0.49, P < 0.006)joints. No relationship was found between ankle DF ROM and other measures of landing performance at the initial contact or the moment of peak flexion.

### Variables

Lateral trunk lean Knee valgus ICA – hip ICA – knee ICA – ankle Peak hip flexion Peak knee flexion Peak ankle flexion

**Table 1**: Correlation between the WBLT and landing
 mechanics in the frontal and sagittal-plane

## Summary and conclusions

These results show that reduced ankle ROM may result in reduced knee flexion and ankle dorsiflexion during single leg landings. However, in single leg landings restrictions in ankle DF ROM do not increase FPPAs.

### **Key Points:**

- risk factor.
- lateral trunk angle

P value	R value
0.277	0.201
0.665	-0.081
0.117	-0.288
0.055	-0.348
0.055	-0.349
0.0.82	-0.317
0.002*	-0.526*
0.006*	-0.486*

Restrictions in ankle DF ROM results in decreased knee flexion and ankle DF, thus increasing injury

Restricted ankle DF ROM does not effect FPPA or Further research into what sufficient levels of ankle DF ROM are required to perform jump landing tasks without compensation.

