

Donovan, Tim (2020) Anatomical Clutter in Medical Image Perception. In: Anatomical Society Winter Meeting 2019, 18th-20th December 2019, Lancaster University. (Unpublished)

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/5409/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
- a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

Anatomical Clutter in Medical Image Perception

Medical image perception is a difficult task, and despite advances in imaging technology error rates of up to 30% still persist. The medical image interpretation task requires cognitive processes such as perception, comprehension, reasoning, decision-making and problem solving, all of which are affected by knowledge, memory and reasoning strategies as well as their biases. Understanding the processes that underlie expert performance in the detection of pathology has not yet been fully achieved.

A conventional medical image, such as a chest x-ray, is a 2D representation of the 3D anatomy. Fully understanding the image means having a 3D mental representation of the anatomy or template to which the image can be compared. One also needs accurate anatomic-pathologic schemas leading to perceptual differentiation of abnormalities. Experts generally detect pathology within 1 or 2 seconds of viewing an image, as they are sensitive to pathology and can attenuate or nullify all the anatomical detail in an image. This is analogous to mentally applying a pre-whitening filter to the image data, and removing anatomical clutter. The greatest motivation for the development of 3D radiographic imaging is that it removes anatomical background clutter that may decrease the conspicuity of pathology.

Currently most radiology trainees are instructed on how to classify abnormalities, whereas it would probably be better to initially instruct them on how to gather the holistic impression of the image such as normal anatomy (and variants) so that they can then fine-tune the perception of pathology. There is no short cut; this requires a great deal of practice and feedback.