

Hayes, Lawrence ORCID: <https://orcid.org/0000-0002-6654-0072> , Sculthorpe, Nicholas and Grace, Fergal M. (2015) Re: Emotions, immunity and sport: winner and loser athlete's profile of fighting sport. *Brain, Behavior, and Immunity*, 47 . p. 238.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/1834/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
 - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

Letter to the Editor

Re: Emotions, immunity and sport: Winner and loser athlete's profile of fighting sport

Brain, Behavior, and Immunity 47, 2015, p.238

Lawrence D. Hayes, Nicholas Sculthorpe, Fergal M. Grace

To the Editor,

We read with interest the recent work of Pesce et al. (2015) who investigated levels of salivary testosterone (sal-T), cortisol (sal-C) and their ratios (T/C) alongside anger (RS Score) and anxiety states (AS score) during the approach to seasonal competition in kick-boxers. The authors make the general assumption that there are associations between sal-T and sal-C with RS and AS scores and the proinflammatory salivary cytokine IL-1 β .

Whilst we found the rationale and study design to be of interest to the behavioural and immunological community, we highlight new limitations with the authors' chosen instrument (saliva hormone measurement) which requires a cautionary approach to the interpretation of their findings. The authors state that sal-T is a reliable indicator of serum or plasma concentrations, however, recent investigations have demonstrated this is not always the case. Furthermore, the reported changes in sal-T, sal-C and sal-T:C are all within the critical difference for measurement, that would confirm that a biologically meaningful change had occurred. In this respect, we draw the authors' attention to the original work of Fraser and Fogarty (1989). Indeed, we have shown the critical difference for sal-T, sal-C and T:C measurement to be 148%, 90% and 207%, respectively. Further to this, our other recent work has indicated that the timing of salivary sampling appears to be at least as important as the type of exercise, such that the true baseline measures should be obtained 24 h or more prior to exercise (Hayes et al., 2015).

Moreover, while the authors correctly refer to the lack of reliability between saliva and plasma cytokines, they fail to repeat the same for sal-T and sal-C. We further note that contrary to their concluding statement, their cited manuscript in support of measuring IL-1 β , in fact, concludes that saliva levels are not directly associated with plasma levels and only becomes significantly associated when adjusted for smoking, age and adiponectin levels in adolescent girls.

Additionally, the authors' supposition that increases in T levels are evident following resistance exercise and that these increases in anabolic hormones are 'essential for muscular adaptation' overlooks some recent compelling evidence to the contrary (Mitchell et al., 2013). In conclusion, we argue the high variability of sal-T and sal-C in controlled laboratory conditions (Hayes et al., 2014)

will be greater in an applied setting, such as the present study and recommend that conclusions based on these measures should be made both with caution and with reference to the critical difference in order to contextualize any observations made.

References

C.G. Fraser, Y. Fogarty, Interpreting laboratory results, *BMJ*, 298 (6689) (1989), pp. 1659–1660

L.D. Hayes, F.M. Grace, J.S. Baker, N. Sculthorpe, Exercise induced responses in salivary testosterone, cortisol, and their ratios in men: a meta-analysis, *Sports Med.* (2015)

L.D. Hayes, N. Sculthorpe, J.D. Young, J.S. Baker, F.M. Grace, Critical difference applied to exercise-induced salivary testosterone and cortisol using enzyme-linked immunosorbent assay: distinguishing biological from statistical change, *J. Physiol. Biochem.*, 70 (4) (2014), pp. 991–996

C.J. Mitchell, T.A. Churchward-Venne, L. Bellamy, G. Parise, S.K. Baker, S.M. Phillips, Muscular and systemic correlates of resistance training-induced muscle hypertrophy, *PLoS One*, 8 (10) (2013), p. e78636

M. Pesce, I. Fratta, V. Ialenti, A. Patruno, A. Ferrone, S. Francheschelli, A. Rizzuto, R. Tatangelo, G. Campagna, L. Speranza, M. Felaco, A. Grilli, Emotions, immunity and sport: winner and loser athlete's profile of fighting sport, *Brain Behav. Immun.* (2015)