

Tyler, Jacqueline ORCID: <https://orcid.org/0009-0002-5396-3380> , Miller, Paul K. ORCID: <https://orcid.org/0000-0002-5611-1354> and Donovan, Tim ORCID: <https://orcid.org/0000-0003-4112-861X> (2023) Can miscarriage be accurately predicted in early pregnancy using transvaginal ultrasound parameters? In: UK Imaging and Oncology Congress (UKIO 2023): Synergy and Symbiosis, Breaking Down Barriers in Healthcare, 5-7 June 2023, Liverpool, UK. (Unpublished)

Downloaded from: <https://insight.cumbria.ac.uk/id/eprint/7080/>

***Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.***

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

**provided that**

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
- a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

**You may not**

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing [insight@cumbria.ac.uk](mailto:insight@cumbria.ac.uk).

# ABSTRACT

**CAN MISCARRIAGE BE ACCURATELY PREDICTED IN EARLY PREGNANCY USING TRANSVAGINAL ULTRASOUND PARAMETERS?**

## INTRODUCTION:

ULTRASOUND IS OFTEN THE FIRST LINE OF INVESTIGATION WHEN EVALUATING EARLY PREGNANCY. CURRENT NICE GUIDELINES REGARDING CONFIRMATION OF MISCARRIAGE ARE RESTRICTIVE AND CAN OFTEN MEAN THAT THE FIRST ULTRASOUND SCAN IS INCONCLUSIVE LEADING TO INCREASED STRESS FOR PATIENTS AND FURTHER SCANS AND INVESTIGATIONS. THERE HAVE BEEN SEVERAL STUDIES UNDERTAKEN TO EVALUATE THE ULTRASOUND PARAMETERS THAT CAN PREDICT MISCARRIAGE THAT HAVE THE POTENTIAL TO BE USED IN CLINICAL PRACTICE ALLOWING BETTER COUNSELLING OF PATIENTS REGARDING THEIR RISK OF PREGNANCY LOSS.

## AIM:

TO UNDERTAKE A LITERATURE REVIEW TO COLLATE THE EVIDENCE AND EVALUATE THE ULTRASOUND PARAMETERS THAT MAY PREDICT EARLY PREGNANCY LOSS.

## METHODOLOGY:

A SYSTEMATIC LITERATURE SEARCH WAS UNDERTAKEN USING PUBMED AND SCIENCE DIRECT INCLUDING LITERATURE PUBLISHED FROM 2012 TO 2022. INCLUSION AND EXCLUSION CRITERIA WERE APPLIED AND ANY LITERATURE SUITABLE FOR INCLUSION WAS CRITICALLY APPRAISED USING THE CASP FRAMEWORK.

## RESULTS:

MEAN SAC DIAMETER, CROWN-RUMP LENGTH, YOLK SAC DIAMETER AND EMBRYONIC/FETAL HEART RATE CAN BE USED TO PREDICT PREGNANCY LOSS. MULTIVARIATE PREDICTIVE MODELS COMBINING THESE PARAMETERS DEMONSTRATED A SENSITIVITY OF 72.5% AND SPECIFICITY OF 98.4% SUGGESTING THAT THEY MAY PROVE USEFUL IN CLINICAL PRACTICE HOWEVER; THESE WOULD NEED TO BE EVALUATED ON A LARGER SCALE IN MULTICENTRE TRIALS TO IMPROVE VALIDITY.

## CONCLUSION

MULTIVARIATE PREDICTIVE MODELS USING ULTRASOUND PARAMETERS CAN PREDICT MISCARRIAGE WITH A GOOD DEGREE OF ACCURACY AND COULD BE USED IN CLINICAL PRACTICE TO FACILITATE BETTER COUNSELLING OF PATIENTS.