

TITLE PAGE

Postural stability of older female Scottish country dancers in comparison to physically active controls.

Acknowledgements

The authors would like to thank Dr Norah Nelson, Dr Paul K. Dougall and Miss Shanita Padaruth for the contribution to this study.

Abstract

Physical activity assists older individuals' functional ability and postural stability. Recently, Scottish country dance (SCD) was reported as being a beneficial form of physical activity for functional ability in older females. This study aims to examine the effect of SCD on postural stability. Scottish country dancers ($n=20$) were compared to physically active controls ($n=33$) for static postural sway measured on a force platform. The Romberg and Tandem stances were used, under 'eyes open' and 'eyes closed' conditions. 95% ellipse area and sway velocity were calculated from the center of pressure displacement. 95% ellipse area was the same for both groups in all tests. The control group had greater sway velocity for all tests ($p<0.01$) except Tandem eyes closed. SCD participation resulted in similar postural sway as participation in other physical activities, however non-dancers may need a greater amount of regulatory activity to maintain balance.

Keywords: aging, balance in elderly, dance, functional ability

Postural stability of older female Scottish country dancers in comparison to physically active controls.

Introduction

Recent reports and initiatives have highlighted the importance of physical activity in older adults, not only for the prevention of disease, but more specifically on postural stability, in a bid to reduce the incidence of falls in this population (American College of Sports Medicine et al., 2009; Howe, Rochester, Neil, Skelton, & Ballinger, 2011). Physically active older participants have been shown to have better static and dynamic postural control than sedentary age-matched individuals (Bulbulian & Hargan, 2000; Perrin, Gauchard, Perrot, & Jeandel, 1999). Indeed, various intervention studies including specific balance training (Clemson et al., 2010), strength training (Henwood & Taaffe, 2006), walking (Brooke-Wavell, Athersmith, Jones, & Masud, 1998), vibration platform training (Cheung et al., 2007), square stepping (Shigematsu et al., 2002), and Tai Chi (Taylor-Piliae et al., 2010) have all been found to improve static postural stability in healthy older adults. Notwithstanding the success of these interventions, the most recent Cochrane review on exercise for improving postural stability in older people highlighted the need to examine further the effects of different forms of physical activity on postural stability in an attempt to strengthen the currently weak evidence (Howe et al., 2011).

Group exercise is one of the most popular physical activities among middle-aged and older adults (Hunt, Ford, & Mutrie, 2001). In particular, social dancing is an activity that is practiced by middle-aged and older adults all over the world. The cultural resonance and social benefits that come with group dance make this an appealing activity for the older age group (Cooper & Thomas, 2002; Lima & Vieira, 2007; Wikstrom, 2004). Varying forms of dance interventions including Traditional Greek, Caribbean, Argentinian tango, Traditional Korean,

1 Turkish Forlore and aerobic dance have shown to benefit both static and dynamic postural
2 stability in older participants (Eyigor, Karapolat, Durmaz, Ibisoglu, & Cakir, 2009; Federici,
3 Bellagamba, & Rocchi, 2005; Hackney, Kantorovich, & Earhart, 2007; Hopkins, Murrah,
4 Hoeger, & Rhodes, 1990; Jeon, Choe, M,A, & Chae, 2000; Sofianidis, Hatzitaki, Douka, &
5 Grouios, 2009). Less research is available on habituated older exercisers, however it has been
6 reported that habituated older folk dancers and gymnasts have better dynamic postural stability
7 than aged matched controls (Uusi-Rasi et al., 1999) likely due to the stresses placed on the
8 neuromotor system.

9 Scottish country dance is a hugely popular activity in older individuals, not only in
10 Scotland, but also in North America, Australia and New Zealand. Scottish country dance, like
11 other traditional dance forms such as square dancing and folk dancing, articulates choreographed
12 multidirectional movements incorporating many elements of neuromuscular control. It consists
13 of a number of set dances in pairs and groups, with varying degrees of intensity. Participants
14 perform various vertical and lateral movements, turning and spinning around in different
15 directions to the rhythms of both slow and faster tempos, therefore continuously altering and
16 stressing their postural stability.

17 Scottish country dance has recently been shown to be beneficial to functional ability of
18 older individuals (Dewhurst, Nelson, Dougall, & Bampouras, in press). The aim of the present
19 study was to compare the postural stability in Scottish country dancers and age-matched
20 physically active controls. It is hypothesized that due to the challenging postural demands of
21 Scottish country dance, the dance group will show better postural stability than the age-matched
22 physically active group, while preventing an increased reliance on the visual system which is
23 normally reported in older adults (Choy, Brauer, & Nitz, 2003; Dewhurst, Riches, & De Vito,

2007). Women are the focus of the present study as it has been reported that women tend to fall more often (Schultz, Ashton-Miller, & Alexander, 1997) and have twice the rate of hip fracture than men (Kannus et al., 1999).

Methods

Participants

Fifty three healthy older females participated in the study after providing written informed consent. The sample consisted of Scottish country dancers ($n = 20$, age 67.9 ± 5.9 (range 60 - 79) years, , body mass 67.2 ± 9.1 kg, stature 1.61 ± 0.06 m) and controls ($n = 33$, age 71.1 ± 5.7 (range 62 - 79) years, body mass 65.0 ± 10.6 kg, stature 1.57 ± 0.07 m).

Participants had no known neuromuscular disorders and had to be considered as medically stable, according to the criteria described by Greig et al (1994). All participants were physically active, engaging in some form of physical activity at least three times a week. The dancers had a minimum of 10 years participatory involvement in Scottish country dancing. Ethical approval was obtained from the University of Strathclyde and all procedures followed were in accordance with the Helsinki Declaration of 1975.

Physical activity assessment

Current levels of physical activity were measured using the Rapid Assessment of Physical Activity (RAPA) scale which is an easy-to-use, valid measure of physical activity for use with older adults (Topolski et al., 2006). This scale consists of 9 tick box responses relating to level and intensity of usual physical activity, as well as additional strength and flexibility activities. RAPA 1 measures cardio-respiratory based physical activity (scoring 1-7 points, with a maximum score of 7), RAPA 2a and 2b measures strength- and flexibility-based physical activity, respectively (yes/ no answer).

1 **Static postural stability assessment**

2 Participants performed the Romberg (standing feet together with arms resting by the side
3 of the body) and the Tandem (feet are aligned heel to toe with the dominant foot at the back)
4 tests standing barefoot and centrally aligned on a force platform (AMTI OR6-6; Advanced
5 Mechanical Technology, Inc., Watertown, Massachusetts). Trials were performed under two
6 visual conditions, eyes open (EO) and eyes closed (EC), administered in a randomised order, and
7 each position was maintained for 35 s with the first 5 s discarded. Two minutes rest was given
8 between trials. If participants lost their balance during a trial, they were allowed to repeat. A
9 successful trial was one in which the subject did not move her feet or, in the case of the EC trials,
10 open their eyes while standing on the platform during the assessment period. A maximum of
11 three attempts were given per trial. During the EO trials, subjects had to focus on a visual target
12 placed two meters in front of the subjects at eye level; participants were allowed to wear
13 spectacles, if required.

14 Data were obtained from the force platform with a sampling frequency of 100 Hz
15 (NetForce, Advanced Mechanical Technology, Inc., Watertown, Massachusetts) and centre of
16 pressure (CoP) based time domain parameters were calculated (BioAnalysis, Advanced
17 Mechanical Technology, Inc., Watertown, Massachusetts). 95% ellipse area (sway area)
18 indicates the amount of CoP movement and is a method used to estimate the confidence area of
19 the CoP path where approximately 95% of the points on the COP path are enclosed in. Sway
20 velocity indicates the speed at which CoP adjustments are made. Both measurements have been
21 used extensively to assess postural balance in older individuals (Brooke-Wavell et al., 1998;
22 Dewhurst et al., 2007; Prieto, Myklebust, Hoffmann, Lovett, & Myklebust, 1996). Finally, in
23 order to assess the influence of vision on the CoP parameters, the Romberg quotient and Tandem

quotient were calculated as the ratio of the eyes closed value to the eyes open value for the sway area.

Statistical analysis

Current levels of physical activity, strength and flexibility exercise participation scores taken from the RAPA scale were analysed using chi-square. Normality of data of the CoP parameters was examined using the Kolmogorov-Smirnov test. As normality was not confirmed for all variables, non-parametric tests were utilized for the analysis. Subsequently, Spearman's correlation was used to examine for relationships between age and sway area or sway velocity and the Mann-Whitney U test was used to examine for differences between dancers and controls. When differences between dancers and controls were revealed, non-parametric effect sizes (ES) were calculated with 0.1 denoting a small effect, 0.3 a medium effect and 0.5 a large effect (Grissom and Kim, 2011). Significance was set at 0.05. For all statistical analyses, IBM SPSS v 19.0 (SPSS, Chicago, ILL) was used.

Results

Physical activity assessment

The quantification of physical activity showed the median score for the cardio-respiratory based physical activity for both groups being 6 on the RAPA scale: "I do 30 minutes or more a day of moderate physical activities, 5 or more days a week". The two groups were not significantly different in participating in strength training ($\chi^2 < 0.05$, 15% for both groups) but the controls participated in flexibility training more than the dancers ($\chi^2 < 0.05$, 61% and 43%, respectively).

Static stability postural assessment

Completion rates per group for each test can be found in Table 1. The age of the groups was not significantly different ($U = 229, p = 0.63$). No significant correlation was found for age with Romberg EC and Tandem EC sway area. Further no significant correlation was found for age with Tandem EO sway velocity. Finally, although significant correlations were found for age with Romberg EO and Tandem EO sway area as well as Romberg EO, Romberg EC and Tandem EC sway velocity they were all low (Spearman's ρ range: 0.317 – 0.392).

Sway area was the same for dancers and controls in all of the postural assessments (Figure 1A and B). Sway velocity was greater in controls than dancers in the Romberg EO ($U = 63, p = 0.001, ES = 0.11$), Romberg EC ($U = 113, p = 0.001, ES = 0.21$) and Tandem EO ($U = 146, p = 0.003, ES = 0.25$) but not Tandem EC. Finally, neither Romberg quotient nor Tandem quotient were affected by the participation in dance (Figure 2).

INSERT FIGURE 1 HERE

INSERT FIGURE 2 HERE

Discussion

The main finding of the present study was that although Scottish country dancers have similar sway area to physically active age-matched controls, an increased amount of regulatory activity may have been needed to maintain it compared to their physically active counterparts, as suggested by the difference in sway velocity.

Scottish country dance involves a combination of proprioceptively challenging moves including unipedal stance and hopping, while maintaining a straight posture with a constant shifting of the center of gravity. Additionally, the vestibular system is challenged during dances as multiple spinning actions are involved. Both the proprioceptive and vestibular systems are

1 responsible for maintaining an upright posture (Dewhurst et al., 2007). It was hypothesized that
2 the stimulation provided by Scottish country dance to the sensory systems and consequently the
3 motor control systems will result in better postural stability when compared to non-dance
4 physical activity participation. While no additional benefits of participation in Scottish country
5 dance were shown for sway area when compared to physically active controls, it appears that
6 mechanisms to achieving this sway area differ between the groups.

7 Sway velocity is related to the amount of activity needed in order to maintain a given
8 level of postural stability (Prieto et al., 1996) and it is an important factor in maintaining postural
9 control in quiet standing (Masani, Popovic, Nakazawa, Kouzaki, & Nozaki, 2003). Increased
10 sway velocity has been linked with reduced balance ability in older individuals (Brooke-Wavell
11 et al., 1998; Dewhurst et al., 2007; Prieto et al., 1996) and increased incidence of falls (Ferne,
12 Gryfe, Holliday, & Llewellyn, 1982). The greater sway velocity of the control group in
13 comparison to the dancers may suggest that more regulatory activity was needed to achieve the
14 same level of sway area, suggesting poorer postural control.

15 The maintenance of a quiet upright stance involves a complex integration of the sensory
16 systems (somatosensory, visual and vestibular), the motor system and a central integrating
17 control system (Horak & MacPherson, 1996). The lower sway velocity in the dancers could be a
18 consequence of a more sensitive sensory input and/or a superior motor response system. As
19 mentioned previously, the vestibular and proprioceptive senses would be continually stressed
20 during the Scottish country dance routines; this in turn may have delayed or reduced the age
21 associated decline in the sensitivity of the mechanoreceptors within the vestibular apparatus and
22 proprioceptors within the joints, tendons and muscles to a greater extent than less dynamic
23 physical activities. In addition, Scottish country dance has been shown to benefit functional

1 motor tasks such as walking speed and 8 foot up and go (Dewhurst et al. in press). This
2 beneficial effect on the motor system may translate to a better regulative motor response. As
3 regular physical activity in older individuals has been found to develop or maintain the efficiency
4 of the reflexes involved in postural control (Gauchard, Gangloff, Jeandel, & Perrin 2003), we
5 propose that it is possible that the nature of Scottish country dance further develops or maintains
6 these reflexes.

7 The ratio between eyes closed to eyes open sway area assesses reliance on the visual
8 system, with an increased ratio suggesting larger reliance on vision (Choy et al., 2003; Dewhurst
9 et al., 2007). When comparing physically active older individuals to sedentary controls, Perrin et
10 al (1999) reported that active individuals relied less on vision during both static and dynamic
11 postural stability tasks. Further, dance has been shown to strengthen the accuracy of
12 proprioceptive inputs and shift sensorimotor dominance from vision to proprioception during a
13 dynamic postural task (Golomer, Cremieux, Dupui, Isableu, & Ohlmann, 1999) however, in the
14 present study the eyes open to eyes closed ratio did not differ between the groups for either the
15 Romberg or the Tandem tests.

16 The lack of a strong relationship between age and postural stability measures reported
17 here may appear contradictory to literature, as static postural stability has been extensively
18 reported to reduce with increased age (e.g. Browne, O'Hare, Finn, & Colin, 2002; Dewhurst et
19 al., 2007; Prieto et al., 1996;). However, although studies have used older subjects of similar
20 characteristics to the present study, the comparisons made are with young counterparts, rather
21 than within the older individuals' age range. This only provides a limited picture as to how
22 balance degrades over years. Baloh et al (1998) reported that in a 3-year follow up of 79-91 year
23 olds, dynamic sway velocity increased annually. However, the age of the participants is outside

1 the age range used in the present study. A comparison of balance in 60-70 and 70-80 year old
2 participants showed a trend for decreased balance with increasing age (Steffen, Hacker, &
3 Mollinger, 2002), however this contradicted a previous study which, similar to our findings,
4 reported little relationship of age to balance (Bogle Thorbahn & Newton, 1996). Given the
5 importance of balance in this specific population, we concur with the call by Steffen et al (2002)
6 for more work in the area with appropriate sample sizes that will allow norms to be established.

7 Static postural stability has been extensively used as a measure of balance in older
8 individuals (Amiridis, Arabatzi, Violaris, Stavropoulos & Hatzitaki, 2005; Dewhurst et al., 2007;
9 Prieto et al., 1996). However, dynamic postural assessment has also been shown to be a sensitive
10 measure in detecting the effect of aging on postural stability (Baloh, Fife, Zwerling, Socotch,
11 Jacobson et al., 1994). Given the dynamic nature of Scottish country dance, dynamic postural
12 assessment may have been more suited to detecting differences between Scottish country dancers
13 and other physically active individuals. Dynamic postural control should be assessed in future
14 works along with fall frequency in dancers compared to other physically active individuals.

15 Bulbulian and Hargan (2000) and Perrin et al. (1999) highlighted the importance of
16 physical activity for postural stability, reporting that physically active older individuals had
17 better postural stability than inactive controls, with suggestions of a dose response both with the
18 length of engagement in physical activity (Bulbulian & Hargan, 2000; Perrin et al., 1999) as well
19 as the quantity of physical activity (Brooke-Wavell et al., 1998). The present study posits that it
20 is possible to maintain postural stability in older adults not only through general physical activity
21 but also through more specific activities such as Scottish country dance.

References

- American College of Sports Medicine, Chodzko-Zajko, W. J., Proctor, D. N., Fiatarone Singh, M. A., Minson, C. T., Nigg, C. R., et al. (2009). American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Medicine and Science in Sports and Exercise*, 41(7), 1510-1530.
- Amiridis I.G., Arabatzi. F., Violaris. P., Stavropoulos. E., & Hatzitaki. V. (2005). Static balance improvement in elderly after dorsiflexors electrostimulation training. *European Journal of Applied Physiology*, 94(4), 424-433.
- Baloh. R.W., Fife. T.D., Zwerling. L., Socotch. T., Jacobson. K., Bell. T., Beykirch. K. (1994) Comparison of static and dynamic posturography in young and older normal people. *Journal of the American Geriatric Society*, 42(4), 405–412.
- Bisson, E. J., McEwen, D., Lajoie, Y., & Bilodeau, M. (2011). Effects of ankle and hip muscle fatigue on postural sway and attentional demands during unipedal stance. *Gait & Posture*, 33(1), 83-87.
- Bogle Thorbahn, L. D., & Newton, R. A. (1996). Use of the Berg balance test to predict falls in elderly persons. *Physical Therapy*, 76(6), 576-83; discussion 584-585.
- Brooke-Wavell, K., Athersmith, L. E., Jones, P. R., & Masud, T. (1998). Brisk walking and postural stability: A cross-sectional study in postmenopausal women. *Gerontology*, 44(5), 288-292.
- Bulbulian, R., & Hargan, M. L. (2000). The effect of activity history and current activity on static and dynamic postural balance in older adults. *Physiology & Behavior*, 70(3-4), 319-325.

- 1 Browne, J., O'Hare, G., Finn, A., Colin, J. (2002). Clinical Assessment of the Quantitative
2 Posturography System (QPS). *Physiotherapy*, 88(4), 217-233.
- 3 Cheung, W. H., Mok, H. W., Qin, L., Sze, P. C., Lee, K. M., & Leung, K. S. (2007). High-
4 frequency whole-body vibration improves balancing ability in elderly women. *Archives of*
5 *Physical Medicine and Rehabilitation*, 88(7), 852-857.
- 6 Choy, N. L., Brauer, S., & Nitz, J. (2003). Changes in postural stability in women aged 20 to 80
7 years. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*,
8 58(6), 525-530.
- 9 Clemson, L., Singh, M. F., Bundy, A., Cumming, R. G., Weissel, E., Munro, J., et al. (2010).
10 LiFE pilot study: A randomised trial of balance and strength training embedded in daily life
11 activity to reduce falls in older adults. *Australian Occupational Therapy Journal*, 57(1), 42-
12 50. Cooper, L., & Thomas, H. (2002). Growing old gracefully - social dance in the third age.
13 *Ageing and Society*, 22(6), 689-708.
- 14 Dewhurst, S., Nelson, N., Dougall, P. M., Bampouras, T.M. (In press). Scottish Country Dance
15 Can Benefit Functional Ability in Older Females. *Journal of Aging and Physical Activity*.
- 16 Dewhurst, S., Riches, P. E., & De Vito, G. (2007). Moderate alterations in lower limbs muscle
17 temperature do not affect postural stability during quiet standing in both young and older
18 women. *Journal of Electromyography and Kinesiology*, 17(3), 292-298.
- 19 Eyigor, S., Karapolat, H., Durmaz, B., Ibisoglu, U., & Cakir, S. (2009). A randomized controlled
20 trial of Turkish folklore dance on the physical performance, balance, depression and quality
21 of life in older women. *Archives of Gerontology and Geriatrics*, 48(1), 84-88.

- 1 Federici, A., Bellagamba, S., & Rocchi, M. B. (2005). Does dance-based training improve
2 balance in adult and young old subjects? A pilot randomized controlled trial. *Aging Clinical*
3 *and Experimental Research*, 17(5), 385-389.
- 4 Fernie, G.R., Gryfe, C.I., Holliday, P.J., & Llewellyn, A. (1982). The relationship of postural
5 sway in standing to the incidence of falls in geriatric subjects. *Age and Ageing*, 11(1), 11-16.
- 6 Gauchard, G. C., Gangloff, P., Jeandel, C., & Perrin, P. P. (2003). Physical activity improves
7 gaze and posture control in the elderly. *Neuroscience Research*, 45(4), 409-417.
- 8 Golomer, E., Cremieux, J., Dupui, P., Isableu, B., & Ohlmann, T. (1999). Visual contribution to
9 self-induced body sway frequencies and visual perception of male professional dancers.
10 *Neuroscience Letters*, 267(3), 189-192.
- 11 Greig, C. A., Young, A., Skelton, D. A., Pippet, E., Butler, F. M., & Mahmud, S. M. (1994).
12 Exercise studies with elderly volunteers. *Age and Ageing*, 23(3), 185-189.
- 13 Grissom, R.J., & Kim, J.J. (2011). Effect sizes for research: univariate and multivariate
14 applications (2nd ed). East Sussex, UK: Routledge
- 15 Hackney, M. E., Kantorovich, S., & Earhart, G. M. (2007). A study on the effects of Argentine
16 tango as a form of partnered dance for those with Parkinson disease and the healthy elderly.
17 *American Journal of Dance Therapy*, 29(2), 109-127.
- 18 Henwood, T. R., & Taaffe, D. R. (2006). Short-term resistance training and the older adult: The
19 effect of varied programmes for the enhancement of muscle strength and functional
20 performance. *Clinical Physiology and Functional Imaging*, 26(5), 305-313.
- 21 Hopkins, D. R., Murrah, B., Hoeger, W. W., & Rhodes, R. C. (1990). Effect of low-impact
22 aerobic dance on the functional fitness of elderly women. *The Gerontologist*, 30(2), 189-
23 192.

- 1 Horak, F. B. & MacPherson, J. M. (1996). Postural orientation and equilibrium. In J. Shepard &
2 L. Rowell (Eds.), *Handbook of Physiology: Section 12, Exercise: Regulation and*
3 *Integration of Multiple Systems* (255-292). New York: Oxford University Press.
- 4 Howe, T. E., Rochester, L., Neil, F., Skelton, D. A., & Ballinger, C. (2011). Exercise for
5 improving balance in older people. *Cochrane Database of Systematic Reviews*, 11,
6 CD004963.
- 7 Hui, E., Chui, B. T., & Woo, J. (2009). Effects of dance on physical and psychological well-
8 being in older persons. *Archives of Gerontology and Geriatrics*, 49(1), e45-50.
- 9 Hunt, K., Ford, G., & Mutrie, N. (2001). Is sport for all? Exercise and physical activity
10 patterns in early and late middle age in the west of Scotland. *Health Education*, 101(4), 151-
11 158.
- 12 Jeon, M. Y., Choe, M.A, & Chae, Y. R. (2000). Effect of Korean traditional dance movement
13 training on balance, gait and leg strength in home bound elderly women. *Journal of Korean*
14 *Academy of Nursing*, 30(3), 647-658.
- 15 Kannus, P., Parkkari, J., Koskinen, S., Niemi, S., Palvanen, M., Jarvinen, M., et al. (1999). Fall-
16 induced injuries and deaths among older adults. *Journal of the American Medical*
17 *Association*, 281(20), 1895-1899.
- 18 Kreutz, G. (2008). Does partnered dance promote health? The case of tango Argentino. *The*
19 *Journal of the Royal Society for the Promotion of Health*, 128(2), 79-84.
- 20 Lima, M. M. S., & Vieira, A. P. (2007). Ballroom dance as therapy for the elderly in Brazil.
21 *American Journal of Dance Therapy*, 29(2), 129-142.

- 1 Masani, K., Popovic, M. R., Nakazawa, K., Kouzaki, M., & Nozaki, D. (2003). Importance of
2 body sway velocity information in controlling ankle extensor activities during quiet stance.
3 *Journal of Neurophysiology*, 90(6), 3774-3782.
- 4 Perrin, P. P., Gauchard, G. C., Perrot, C., & Jeandel, C. (1999). Effects of physical and sporting
5 activities on balance control in elderly people. *British Journal of Sports Medicine*, 33(2),
6 121-126.
- 7 Prieto, T. E., Myklebust, J. B., Hoffmann, R. G., Lovett, E. G., & Myklebust, B. M. (1996).
8 Measures of postural steadiness: Differences between healthy young and elderly adults.
9 *IEEE Transactions on Bio-Medical Engineering*, 43(9), 956-966.
- 10 Schultz, A. B., Ashton-Miller, J. A., & Alexander, N. B. (1997). What leads to age and gender
11 differences in balance maintenance and recovery? *Muscle & Nerve, Supplement*, 5, S60-64.
- 12 Shigematsu, R., Chang, M., Yabushita, N., Sakai, T., Nakagaichi, M., Nho, H., et al. (2002).
13 Dance-based aerobic exercise may improve indices of falling risk in older women. *Age and*
14 *Ageing*, 31(4), 261-266.
- 15 Sofianidis, G., Hatzitaki, V., Douka, S., & Grouios, G. (2009). Effect of a 10-week traditional
16 dance program on static and dynamic balance control in elderly adults. *Journal of Aging and*
17 *Physical Activity*, 17(2), 167-180.
- 18 Steffen, T. M., Hacker, T. A., & Mollinger, L. (2002). Age- and gender-related test performance
19 in community-dwelling elderly people: Six-minute walk test, Berg balance scale, timed up
20 & go test, and gait speeds. *Physical Therapy*, 82(2), 128-137.
- 21 Taylor-Piliae, R. E., Newell, K. A., Cherin, R., Lee, M. J., King, A. C., & Haskell, W. L. (2010).
22 Effects of Tai Chi and Western exercise on physical and cognitive functioning in healthy
23 community-dwelling older adults. *Journal of Aging and Physical Activity*, 18(3), 261-279.

- 1 Topolski, T. D., LoGerfo, J., Patrick, D. L., Williams, B., Walwick, J., & Patrick, M. B. (2006).
2 The rapid assessment of physical activity (RAPA) among older adults. *Preventing Chronic*
3 *Disease*, 3(4), A118.
- 4 Uusi-Rasi, K., Sievanen, H., Vuori, I., Heinonen, A., Kannus, P., Pasanen, M., et al. (1999).
5 Long-term recreational gymnastics, estrogen use, and selected risk factors for osteoporotic
6 fractures. *Journal of Bone and Mineral Research*, 14(7), 1231-1238.
- 7 Wikstrom, B. M. (2004). Older adults and the arts: The importance of aesthetic forms of
8 expression in later life. *Journal of Gerontological Nursing*, 30(9), 30-36.

9
10

1 **Table legend**

2 Table 1. Completion rates (in percentages) per group for each static postural assessment. EO,

3 eyes open; EC, eyes closed.

4

1 Table

2

	Sway area				Sway velocity			
	Romberg	Romberg	Tandem	Tandem	Romberg	Romberg	Tandem	Tandem
	EO	EC	EO	EC	EO	EC	EO	EC
Controls	94	97	94	82	91	97	94	76
Dancers	95	100	90	70	95	85	95	70

3

Figure Legends


Figure 1. Posturography parameters during the Romberg and Tandem assessment: 95% Ellipse Area Romberg (A), 95% Ellipse Area Tandem (B), Sway Velocity Romberg (C), Sway Velocity Tandem (D) for the eyes open (Romberg (REO) and Tandem (TEO)) and eyes closed (Romberg (REC) and Tandem (TEC)) trials. Both dancers (open bars) and control groups (filled bars) are shown.

* denotes significant difference between the dancers and control groups ($p < 0.01$). Data are presented as mean \pm standard error.


Figure 2. 95% Ellipse Area Quotient (eyes closed/ eyes open). Romberg quotient (RQ) and Tandem quotient (TQ) are displayed for the dancers (open bars) and control groups (filled bars). Data are presented as mean \pm standard error.

Figures

Figure 1.



1 Figure 2



2
3
4
5
6