

Relph, Nicola ORCID: <https://orcid.org/0000-0003-3323-3444> , Small, Katie ORCID: <https://orcid.org/0000-0001-7908-7441> and Smith, Adam (2017) The effect of running nine marathons in nine days on proprioception and balance. *British Journal of Sports Medicine*, 51 (4). p. 378.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/2706/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
 - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

The effect of running nine marathons in nine days on proprioception and balance

Nicola Relph, Katie Small, Adam Smith

Monaco abstracts 239, British Journal of Sports Medicine 2017; 51 (4):378, Published online 1 February 2017. <http://dx.doi.org/10.1136/bjsports-2016-097372.239>

Abstract

Background Multi-day running events are becoming increasingly popular with nonprofessional athletes. However, it is unknown how and when this type of event affects injury risk factors such as proprioception and balance.

Objective To evaluate the effect of running one, five consecutive and nine consecutive marathons on joint position sense (JPS) and dynamic balance.

Design Cross-sectional, observational study.

Setting Sports-injury clinic.

Participants 21 athletes completing the event (age 44.8 ± 8.09 yrs, mass 74.4 ± 13.14 kg, years running 10.4 ± 7.39 yrs, average weekly mileage 44.8 ± 12.09 miles) entered the study. Ten athletes completed the JPS testing and five athletes completed the dynamic balance testing.

Assessment of Risk Factors Risk factors were assessed the day before the event (day zero) and then following completion of one (day one), five (day five) and nine (day nine) consecutive marathons. The independent variable was time.

Main Outcome Measurements JPS error into flexion and extension, star-excursion balance test (SEBT) and number of injuries per runner.

Results Friedman's ANOVAs indicated no effects of time on JPS into extension ($p=0.94$), dominant-leg anterior ($p=0.52$) or posterior-lateral ($p=0.65$) SEBT reach. JPS flexion error significantly decreased by 1.3° from day one to five ($p=0.032$). Non-dominant leg anterior, posterior-medial and posterior-lateral SEBT reaches significantly decreased from day zero to nine by 0.09% ($p=0.043$), 0.13% ($p=0.043$) and 0.17% ($p=0.043$) of leg length respectively. Dominant leg posterior-medial SEBT reach also significantly decreased by 0.12% of leg length ($p=0.043$) from day zero to nine. There were 4.2 injuries per runner.

Conclusions The results suggest multi-day running events can cause over four lower limb injuries per runner and reduce dynamic balance ability. Nonprofessional athletes completing these events should be aware of this high injury occurrence and prepare appropriately. Injury prevention programmes incorporating dynamic balance may be recommended. However future studies with additional risk factors and larger sample sizes are needed to substantiate these findings.