

Leslie, Andrew ORCID: <https://orcid.org/0000-0001-6327-1711> , Mencuccini, Maurizio and Perks, Mike P. (2017) A resource capture efficiency index to compare differences in early growth of four tree species in northern England. *iForest: Biogeosciences and Forestry*, 10 (2). pp. 397-405.

Downloaded from: <https://insight.cumbria.ac.uk/id/eprint/2790/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
- a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

A resource capture efficiency index to compare differences in early growth of four tree species in northern England.

Leslie, A.D¹.; Mencuccini, M². and Perks, M.P³.

Key words:

Eucalyptus gunnii, Fraxinus excelsior, Acer pseudoplatanus, Alnus glutinosa, resource capture efficiency

¹Senior Lecturer, National School of Forestry, University Of Cumbria, Ambleside Campus, Rydal Road, Ambleside, LA22 9BB. Andrew.leslie@cumbria.ac.uk (corresponding author)

²Professor, School of Geosciences, University of Edinburgh, Crew Building, The King's Buildings, West Mains Road, Edinburgh EH9 3JN. M.Mencuccini@ed.ac.uk

³Project Leader, Centre for Forestry and Climate Change, Forest Research, Northern Research Station, Roslin, Midlothian EH25 9SY. mike.perks@forestry.gov.uk

Abstract

At a trial established in Cumbria, northern England, significant differences in growth rate between tree species were apparent, with cider gum (*Eucalyptus gunnii*) and alder (*Alnus glutinosa*) exhibiting most rapid volume and biomass accumulation. Estimations were made of leaf area, specific leaf area, leaf area ratio (based on stem mass not whole tree mass) and length of growing season. These measurements were undertaken to explain tree growth difference and developing a growth potential index based on growing season length and leaf area. The high leaf area of cider gum and alder explained some of their superior growth, while alder also had the longest period in leaf, compared with ash (*Fraxinus excelsior*) and sycamore (*Acer pseudoplatanus*). The slow growth of ash can be explained by the short period in leaf and also the relatively low leaf area ratio.

Leaf area to stem weight also differed between species with that of ash being relatively low. Specific leaf area was also low for ash, a trait shared with cider gum, which suggests that these species invest highly in each unit of leaf area. Of the tree species assessed, the length of the growing season was longest for alder, enabling it to maintain growth for a longer period. By multiplying growing season by leaf area a resource capture index was calculated and this explained 56% of the variation in stem dry weight between trees. The potential and limitations for using this index are discussed.

Introduction

This paper compares the early growth at a trial in northern England of four broadleaved trees species identified as having potential for short rotation forestry (Hardcastle 2006). Short rotation forestry for bioenergy was identified as a cost-effective and rapid means of reducing greenhouse gases in Matthews and Broadmeadow (2009), yet there is very limited experience of such systems in the UK. While quantifying yields from tree species in short rotation forestry systems is crucial, understanding the underlying factors contributing to their productivity is also important. Yield varies considerably between tree species and on a specific site is dependent on their resource capture efficiency and resource use efficiency (Stape et al. 2004). Resource capture efficiency represents how efficiently a tree is able to allocate its resources to harvesting light for photosynthesis. There are two components; leaf mass fraction (LMF) and specific leaf area (SLA) (Poorter et al. 2012). LMF (the ratio of leaf dry mass to total plant dry mass) reflects the priority of allocation of photosynthate to the leaves. Differences in LMF exist among tree species, with higher LMF being an attribute of faster

growing trees (Poorter et al. 2012) and also shade bearing trees (Valladares & Niinemets 2008). LMF is higher in evergreen than deciduous trees but this is partly because evergreen trees retain leaves for two or more years (Poorter et al. 2012). This higher LMF in evergreens may therefore be largely because of the lifespan of the leaf rather than higher partitioning of annual photosynthate to leaves (Poorter et al. 2012).

SLA reflects the amount of resources allocated to each unit of leaf area and is the ratio of leaf area to leaf dry mass. SLA is generally higher in conifers, compared with broadleaves and higher in fast growing trees than those that are slow growing. A meta-analysis undertaken by Poorter et al. (2009 in Poorter et al. 2012) found differences in SLA between types of trees were greater than for LMF. High SLA is a characteristic of plants that have a high relative growth rate (RGR), small seed mass and both RGR and SLA were good predictors of a plant's potential invasiveness. Leaf area ratio (LAR) is a variable that combines LMF and SLA, being the ratio between leaf area and total tree weight and strongly influences RGR, particularly on nutrient rich sites (Poorter & Remkes 1990).

Growth of a tree is also influenced by the duration of its growing season. The length of growing season has a strong influence on a tree's productivity and differences exist among species, origins and individual trees in terms of their period of dormancy. The dormancy in pioneer trees is largely determined by temperature, rather than photoperiod. In contrast, late successional trees follow a more conservative approach, requiring a longer period of chilling and are highly sensitive to photoperiod (Basler & Korner 2012). Nutrition has also an influence on period of dormancy; tree species with enhanced access to nutrients, including nitrogen fixing trees like alder adopt a higher risk approach to the development of their foliage, in a similar way to pioneer trees (Tateno 2003).

At this trial, growth was measured but also attributes important in influencing this growth were quantified, and these were compared between the tree species. A resource capture efficiency index was devised (leaf area x period in leaf) and its relationship to stem volume was investigated. Results are compared with those of other researchers and differences observed between the tree species explained by their ecology.

The hypotheses tested were that:

- there were differences in volume and biomass growth, SLA and LAR between the tree species at the trial;
- growth of stem volume was positively related to leaf area and growing season;
- the differences in growth between individuals and species can be explained by their resource capture efficiency and other aspects of their ecology.

Materials and methods

The Experiment

A one hectare trial was established at Newton Rigg, Cumbria, England (55°40'56"N, 2°47'22"W) at an elevation of 160 m above sea level. Ash (*Fraxinus excelsior*), sycamore (*Acer pseudoplatanus*) and alder (*Alnus glutinosa*) were planted in winter 2008 and cedar gum (*Eucalyptus gunnii*) and shining gum (*Eucalyptus nitens*) in April 2009. Excluding the eucalypts, species selected for the trial were all classified by EMIS (Perks, Harrison & Bathgate 2006) as being "suitable", rather than "very suitable" or "unsuitable" for the site. The transplants were cell grown 20-40 cm plants raised by Alba Trees in the Scottish borders and origins are described in Table 1, along with recommended origins for the species.

<Table 1 here>

Aspects of the climate at the trial were obtained from the weather station at the Newton Rigg campus and from The Establishment Management Information System (EMIS) (Perks et al. 2006) and are shown in Table 2.

<Table 2 here>

The trial site was originally grass pasture, the soil being a slightly acid (pH6) clay loam brown earth. Soil bulk density was 0.76 at 0 cm to 15 cm depth and 1.07 at 15 cm to 30 cm depth, while the soil nitrogen concentration was 0.36g cm⁻³ and 0.29 g cm⁻³ at 0 cm to 15 cm and 15 cm to 30 cm depth respectively (Centre for Ecology & Hydrology 2013). Soil depth was not measured but was sufficiently deep to not be a limitation to early tree growth.

The site use was previously improved pasture, which was sprayed before planting with Propazymide (3.75 litres ha⁻¹) to kill the sward. The transplants were planted manually using a "T" notch and protected in 60 cm tubes. Planting was at 2.5m spacing between rows and 1.5 m spacing within rows giving a stocking rate of 2,667 stems ha⁻¹. Routine spraying of weeds was undertaken twice during each growing season using glyphosate (5 litres ha⁻¹). The design adopted for the trial was a randomised complete block design, which is the most commonly used design in forest experiments (Wright and Andrew 1976) and large 10 x 8 tree plots were adopted.

Measurements and overview of analysis for each study

Measurements taken in the studies are described in the four sections below. For the measurements two trees of each of the four species were selected randomly from each of the six replicates in the trial. Shining gum was killed in its first winter and is not included as part of this study.

Stem volume and biomass

Height and root collar diameter were measured for the twelve trees of each species in November 2010 (after two growing seasons) and November 2011 (after three growing seasons). Between November 2010 and 2011 all the cider gum had died and so the later measurements were not possible for this species. Height was measured using a height rod, while the root collar diameter was measured to the nearest 0.1 mm using a digital vernier gauge and taking the mean of two measurements taken at 90° to one another. Stem volumes were calculated using height (h) and diameter (d) and assuming that the tree stems were conical in shape. To enable stem weights for the trees to be estimated, wood samples were taken from a different sample of five trees of each species and sections were cut from the base, middle and top of their stems. Volumes of these stem samples were measured using a water displacement method using OHAUS analytical standard scales to measure weight which was converted to a volume using a water density of 1 g cm⁻³. Stem samples were oven dried at a temperature of 80°C for 3 days, until no further loss in weight was observed and then weighed again to obtain dry weight. Specific gravity (SG) was then calculated for the wood samples and SG (g cm⁻³) and V (cm³) were multiplied together to estimate whole-stem dry weight of the study trees.

Leaf Area

In September 2010 after two growing seasons, the crowns of the twelve alder, ash and sycamore trees measured for stem volume were wrapped in plastic bird netting to trap leaves as they fell. For sycamore the collections of leaves in late October was straightforward as most of the leaves had already been shed but for alder, many leaves had to be carefully removed from the crowns of the sample trees. For trees with less than fifty leaves all leaves were measured and for those with more than 50 leaves, all leaves were counted and a systematic sample of 50 was taken. For each leaf, length (L) along the mid rib and width (W) at the widest point of the lamina and petiole length (P) was

160 measured to the nearest millimetre. The use of netting to capture leaves proved to be unsuitable for
161 trapping leaves of ash as their compound leaves disintegrated and some of the small leaflets fell out
162 of the bird netting. As such, the leaf length and width could not be measured but the leaf stalk (S)
163 without the leaflets, which remained trapped in the netting was measured for each leaf. For cider
164 gum, the evergreen species, the method of trapping fallen autumn leaves was not appropriate. For
165 each of the twelve trees, all the leaves were counted, classified as mature or juvenile and 50 leaves
166 were removed from the trees in a systematic way from the bottom to top of the trees to ensure a
167 good spread. Measurements of L, W and P were taken for each type of leaf.

168
169 From the leaves collected, a sample of forty was taken for each tree species across the range of
170 sizes. L, W and P was measured and also S for ash and the leaf area (LA) was then determined
171 using Compu Eye software and an Epson Perfection 1240 flatbed scanner. For cider gum forty
172 juvenile leaves and forty mature leaves were measured. For all species, leaves were then dried for
173 48 hours at a temperature of 70°C and weighed to obtain an oven-dried weight (M) using OHAUS
174 analytical standard scales and following the approach adopted by Verwijst and Wen (1996). As the
175 original ash leaves had disintegrated new ash leaves were collected at the end of the summer of the
176 following year for leaf area and weight determination purposes.

177
178 The total leaf area for the twelve sample trees of the four species was calculated using allometric
179 methods, similar to the approach adopted in other studies (Wargo, 1978; Verwijst & Wen 1996,
180 Ugeese et al. 2008, Serdar & Demirsoy 2006). This involved the determination of relationships
181 between measurements of L and W (and S for ash) to leaf area and leaf weight using least squares
182 regression. Best fit functions were selected based on R^2 and standard error statistics. Best fit
183 relationships were used to estimate the leaf area of each leaf sampled from the twelve trees of each
184 species. For each tree, a mean leaf area was calculated and this was multiplied by the total number
185 of leaves present to obtain an estimate of total leaf area per tree. For the twelve trees of each
186 species, the results from the leaf area measurements and of the stem weights were used to SLA (leaf
187 area/ leaf dry mass) and LAR (leaf area/stem dry weight) parameters. LAR was calculated based on
188 stem dry weight rather than the whole tree weight as below ground biomass was not assessed.

189 **Growing season**

190
191 The same twelve trees of each species used in the leaf area study, were assessed during 2011 to
192 determine the length of the third growing season of the tree species at the trial. The method adopted
193 elements from a study of leaf development in rowan (*Sorbus aucuparia*) (Forest Research no date)
194 and one investigating leaf senescence in birch (*Betula pendula*) (Worrell 2006). The development
195 of the bud was scored on a 0-5 scale with 0 for a dormant bud and 5 for full leaf expansion (the scale
196 was 1 to 6 in original study from Forest Research, no date). The stages in the bud burst scoring
197 were as follows:

- 198
199 0. Bud is closed and in a fully dormant winter state
200
201 1. Bud is swollen and the bud scales just started to open, however the bud is still vertical
202
203 2. Bud scales have separated and the tightly furled leaves are visible. The bud is bent sideways and
204 can appear “hooded”
205
206 3. Bud scales are completely separated, leaves are starting to unfurl and separate but the leaflets
207 (pinnae) on each leaf remain still furled. The leaves appear brownish in colour since the
208 underside is predominantly visible.
209
210 4. The leaves are elongated and leaflets have started to separate as well. The appearance is now
much more green since the topside of the leaves is now visible
211
212 5. All leaflets have separated on the lowest two leaves and the shoot is expanding.

213 The end of the growing season was assessed through a five stage leaf retention score based on a
214 four stage scoring system originally developed by Worrell (2006) (a zero was added for no leaves).

As the trees were still relatively small, the assessment was made by estimating the percentage of the combined leaf area of the tree crown which was still green, not yellow or brown or had lost leaves. This was scored in the following categories

0. No leaves present;
1. One leaf present to 20% of crown green;
2. 21-40% of crown green;
3. 41-60% of crown green;
4. 61-80% of crown green;
5. 81-100% of crown green.

For ash, sycamore and alder the growing season length was calculated by multiplying the bud development score or the leaf retention score by number of days. This gave a relative measure of photosynthetic duration. The growing season for cider gum, an evergreen tree could not directly be measured in the same way.

Resource capture efficiency

For each of the twelve ash, alder and sycamore trees the influence of growing season and leaf area on growth was examined using a resource capture efficiency index, calculated by multiplying tree growing season (collected in 2011) by leaf area (collected in 2010). The formula and units of measurement are described below:

Resource capture index = leaf area (m²) x growing season (leaf retention score x days)

The relationship between stem dry weight in 2010 and resource capture efficiency was investigated.

Statistical analysis

Statistical tests were conducted using IBM SPSS Statistics v19. The approaches used to test variables for significance of differences between species is described in Figure 1.

<Figure 1 here>

Regression was used to characterise relationships between L, W and LA and between L, W and LDW. IN SPSS there were eleven functions available for regression and the best-fit equation was selected by a combination of the smallest standard error of the estimate (SEE) and the highest R². The same approach was used to define a relationship between stem dry weight and the resource capture efficiency.

Results

Where non parametric approaches to analysis have been applied the median is shown as a measure of centrality, otherwise the mean for the data is presented.

Stem volume and biomass

The stem volumes of the twelve trees of the four tree species for were assessed in late 2010 after two growing seasons and for the surviving three species in late 2011 after three growing seasons. The data for height, basal stem diameter and calculated volume are shown in Table 2. For the two and three growing seasons height, diameter and stem volume were compared between species and significant differences were found by species (Table 3).

263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312

<Table 3 here>

Table 3 shows the stem volume, specific gravity and stem dry weights after two growing seasons. Kruskal Wallis and Mann Whitney tests were applied to these data and significant differences were found and are described in Table 4. Medians for these data are shown.

<Table 4 here>

Leaf Area

To determine leaf area, relationships between L, W and LA and between L, W and LDW were investigated. The results from best-fit regressions are described in Table 4 and Table 5. LA (Table 5) and LDW (Table 6) were estimated for the twelve trees of each species by applying the regression models to the L x W measurements for all but ash, where they were estimated from leaf stalk length. Kruskal Wallis test and Mann Whitney tests showed highly significant differences in leaf number, LA and LDW between species. The LA of alder and sycamore were not significantly different, but all others were significantly different. The LDW of cider gum was different from all others, but differences between the other species were not significant. In terms of leaf number, differences between ash and sycamore were not significantly but were significantly different from other species as were alder and cider gum.

<Table 5 here>

<Table 6 here>

LAR and SLA was calculated for the four tree species and the median values are shown in Figure 2. Statistically significant differences in LAR and SLA were found between species.

<Figure 2 here>

Kruskal Wallis and Mann Whitney tests were applied to the LAR data and ash LAR was found to be significantly different from the other three species. Kruskal Wallis and Mann Whitney tests demonstrated that the SLA of all species were significantly different from each another, except for ash and cider gum. P values were significant for SLA and very highly significant for LAR.

Growing season

The period of bud burst and senescence for ash, sycamore and alder for 2011 are shown in Figure 3. Alder had a longer growing season than the other two species, with an earlier and more rapid bud burst and a later and longer period leading up to complete leaf drop. Ash and sycamore showed a similar response, with sycamore having more rapid bud burst and being slower to drop its leaves.

<Figure 3 here>

Cumulative growing season index data was not normally distributed and a Kruskal Wallis test showed that differences in growing season between species were highly significant, while Mann Whitney tests showed that there were very highly significant differences between sycamore and alder, and ash and alder.

Characterising resource capture efficiency.

313 A correlation was performed of resource capture index against stem dry weight, which gave a R of
314 0.729 and was very highly significant ($p < 0.0001$). A regression was conducted and the best fitting
315 function was quadratic (Figure 4, $R^2 = 0.557$, $SEE = 1.233$, $y = -0.17x^2 + 0.683x + 2.267$).

316
317 <Figure 4 here>

318
319 As a comparison correlations were performed for the two components (growing season index and
320 LA) separately of the resource capture index against stem dry weight. For growing season index the
321 correlation with stem dry weight was relatively weak with a R of 0.444 and a high significance
322 ($p < 0.01$). In contrast the correlation between dry weight and LA was stronger with a R of 0.733 and
323 was very highly significant ($p < 0.0001$). A regression showed the best fitting function to be cubic
324 (Figure 5, $R^2 = 0.542$, $SEE = 1.256$, $y = -1.977x^3 + 2.132x^2 + 5.093x + 2.391$).

325
326 <Figure 5 here>

327 328 Discussion

329
330 The winter of 2009-2010 was the coldest in the UK since 1978-1979 and the UK experienced the
331 coldest December in 100 years (Prior & Kendon 2011). This was followed by another severe winter,
332 which apart from that of 2009-2010, was the coldest since the winter of 1985-1986 (Anon 2011). The
333 native and naturalised broadleaves were able to cope with these conditions, but the eucalypts fared
334 badly, particularly the less cold-tolerant shining gum which exhibited complete mortality over each of
335 the two severe winters. An analysis of the cold damage at the trial is described in Leslie, Mencuccini
336 and Perks (2014). Eucalypts have been planted on a limited extent in the UK; between 2011 and
337 2016 nurseries sold 220,000 transplants (Purse and Leslie 2016b) and this scale represented trial
338 rather than commercial planting. However the limited information on suitable origins for cider gum
339 (Cope, Leslie and Weatherall 2008) and shining gum (Evans 1986) suggests that the origins used in
340 this trial were not particularly well adapted. Furthermore it is accepted that shining gum is one of the
341 least hardy eucalypts planted in the UK and it is recommended that it be planted only within 20km of
342 the coast in southern England and closer to the coast elsewhere in the UK (Purse and Leslie 2016a).
343 The severely cold conditions and the origins planted explain some of the eucalypts poor
344 performance.

345
346 Of the tree species that survived the first winter, the largest stem volumes were achieved by cider
347 gum and alder, with the eucalypt producing nearly twice the volume of alder (Table 3). Cider gum
348 had accumulated the largest leaf area, which would partly explain the fast growth of this species
349 (Table 5). The median leaf area of trees of alder (0.1919 m^2) was significantly different from ash
350 (0.0627 m^2) and cider gum (0.4999 m^2), while that of sycamore (0.1856 m^2) was significantly different
351 from cider gum. While ash had the lowest leaf area, it attained nearly the same stem volume as
352 sycamore suggesting that it exhibits greater photosynthetic efficiency. Also, the leaf area of alder
353 was not significantly different from sycamore yet it attained a significantly larger stem volume. These
354 observations suggest sycamore has low photosynthetic efficiency. By the end of the third growing
355 season cider gum had been killed in the winter of 2010-11 and the relative ranking of the remaining
356 species in terms of stem volume remained the same.

357
358 The leaf area of the trees was measured at the end of the growing season and this may not have
359 fully captured the leaf area over the whole season, as it does not incorporate leaf turnover. There
360 are considerable differences in leaf longevity between temperate tree species; mean leaf lifespan of
361 alder is 90 days and in maples and oaks can be as long as 180 days. (Kikuzawa 1995). Leaf
362 longevity may explain some of the differences found between species in SLA. The SLA of alder and
363 sycamore was relatively high and cider gum and ash was relatively low (with no significant difference
364 between the two). This suggests a greater allocation of resources per unit leaf area in cider gum and
365 ash and less resources per unit leaf area in sycamore and alder. Generally there is a positive
366 relationship between leaf mass: leaf area and the longevity of the leaves (Wright & Westoby 2002).
367 Thus some trees invest relatively little in each metre of leaf area, allowing rapid build up of canopy,

fast cycling of leaves and high initial growth. In contrast other trees invest more heavily in each square metre of leaf area but retain these leaves for longer, resulting in a longer period of return from those leaves (Wright & Westoby 2002).

In terms of SLA, this would suggest that trees which retain their leaves for longer periods will have a lower SLA and those with short leaf longevity have a high SLA. Alder leaves are retained by the tree for a relatively short period (Kikuzawa 1995) and so, as found in this study (Figure 2) exhibit a relatively high SLA of ($8.8 \text{ m}^2 \text{ kg}^{-1}$) which would support such a strategy, each leaf being allocated a relatively low investment of resources. There are no studies of the leaf longevity of cider gum, but Whitehead and Beadle (2004) note that in general eucalypt leaves are thick, tough and long-lived, a reflection of their evergreen habit and their association with sites of low soil nutrients and mild winters. A study in Australia found *Eucalyptus paniculata* leaf lifespan to be 1.09 years and that of *Eucalyptus umbra* to be 2.06 years (Wright & Westoby 2002), but Laclau et al. (2009) studying *Eucalyptus grandis* in Brazil found unfertilised trees in plantation retained their leaves for 111 days. The relatively low SLA ($4.9 \text{ m}^2 \text{ kg}^{-1}$) of cider gum (Figure 2) suggests a relatively long leaf lifespan. Ash also exhibited a low SLA and a study by Alberti et al. (2005) of older trees also found a low SLA for ash, compared with Wych elm (*Ulmus glabra*). Another characteristic of trees with high SLA, such as the alder and sycamore in this study, is that they tend to exhibit high photosynthetic nitrogen use efficiencies, whereas trees with a low SLA adopt a different strategy; absorbing a greater proportion of the light available through a higher chlorophyll content in the leaves (Poorter & Evans 1998).

Statistically significant differences in specific gravity were found between species, with alder having a particularly low density (Table 4), also being lower than the 0.540 Mg m^{-3} cited by Claessens (2005 in Claessens et al. 2010), perhaps due to the young age of the trees. The specific gravity of ash was similar (0.550 as opposed to 0.560 Mg m^{-3}) to that found in larger trees from Italy (Alberti et al. 2005) and that of cider gum was similar (0.548 as opposed to 0.500 Mg m^{-3}) to that found in French plantations (AFOCEL 2003). The specific gravity was multiplied by stem volume enabling LAR (using stem weight rather than the conventional whole tree weight) to be calculated. This was compared by species and very highly significant differences were found between ash and all other species. Therefore ash supports a smaller leaf area per unit stem weight than sycamore, alder and cider gum (Figure 2). A shortcoming of this experiment was not to estimate branch and root weight of the trees, as this would have enabled a true LAR to be calculated and a better understanding of the relative allocation of resources between different parts of these trees. There are known to be differences in the allocation of resources between stem and leaves and roots between tree species (Poorter et al. 2012). At the trial differences were observed; some individuals of shining gum exhibited instability due to excessive above ground growth and had fallen over, despite being supported by a tree shelter and bamboo cane. The other species showed good stability.

Phenology of temperate trees is determined by temperature and photoperiod, with the importance of each of these factors varying with tree species (Basler & Körner 2012, Vitasse et al. 2012,). This study used visual assessment of budburst, which is the normal method used in field dormancy studies (Cooke et al. 2012). The pattern of bud burst and leaf fall between ash, alder and sycamore is illustrated in Figure 3. This shows that alder begins to come into leaf earlier than the other two species and also retains its foliage for longer into autumn and that ash flushes later and loses leaves earlier in autumn than the other two tree species. Basler and Körner (2012) found that there was no effect of photoperiod on bud burst of ash or sycamore, while a study (Vitasse et al. 2009) on the effect of temperature on budburst in seven temperate trees showed that of those planted at this trial, ash had the highest sensitivity to temperature, with sycamore being in the middle of the ranking. Spring 2011, when the assessment was made was particularly warm, being the warmest across the UK since 1910 (Met Office undated c). It is likely therefore that the growing season for 2011 was abnormally long for these species.

The phenology data for ash, sycamore and alder were based on monitoring the development and senescence of leaves on the terminal bud but development of leaf area in trees is complex. Focusing on the terminal bud does not allow the pattern of whole tree leaf area to be examined and

pioneer trees tend to adopt a different approach to climax species. Climax or forest tree species show a flushing habit of leaf development, whereas pioneers show a successive pattern of leaf development (Kikuzawa 1995). The patterns of flushing between alder, ash and sycamore showed differences (Figure 3). The progression of leaf unfolding started earlier in alder but was also more gradual in alder than in the other two species, which exhibited rapid flushing over a relatively short period.

Combining leaf area measurements from 2010 with growing season data from 2011 to create a resource capture index explained 56% of the differences in 2010 stem dry weight of the trees (Figure 4). The best fitting relationship in terms of R^2 was a curve but a linear relationship also provided a good fit ($R^2=0.548$). The nature of the relationship is difficult to identify precisely because of the lack of data at the higher end of the combined leaf area and growing season index. A possible explanation for a curved relationship between growth potential index and stem dry weight is that light interception by canopies is not linearly related to leaf area index, but follows a similar curved relationship due to mutual shading of leaves (Cannell et al. 1989). This is supported by the curvilinear relationship between leaf area and stem dry weight (Figure 5). Using leaf area alone explained 54% of the variation in stem dry weight, only marginally poorer than combining leaf area and growing season. This can be explained by the relatively small differences in growing period between the tree species. Had the origins been collected across a wider range of latitudes it is likely that variation in growing season between species would have been greater.

Growth is related to three variables; the site resources, the resource capture efficiency and the resource use efficiency (Stape et al. 2004). Multiplying leaf area by growing season provided a measure of the resource capture efficiency of the tree species at this trial. A combination of greater leaf area and longer period of growth has enabled alder to accumulate stem dry weight more rapidly than ash and sycamore (Figure 4). The rate of photosynthesis in a tree species is strongly linked to the nitrogen content of leaves due to large amount of leaf nitrogen devoted to chloroplasts (Poorter & Evans 1998) and alder, being a nitrogen fixing tree is likely to be able to devote larger concentrations of nitrogen to its leaves than the other species. This study also showed that alder exhibited a high SLA, (Figure 2) allocating relatively little biomass for every square metre of leaf area. Trees with high SLA are known to exhibit high photosynthetic nitrogen use efficiency (Poorter & Evans 1998) and in general high relative growth rates (Antinez et al. 2001). The higher leaf nitrogen concentration and this higher photosynthetic nitrogen efficiency may partly explain why alder has been able to build up leaf area rapidly and also use this leaf area efficiently. A further strength of alder is its relatively long growing season compared with sycamore and ash (Figure 3). Alder, is also known to have a short leaf longevity (Koike & Sanada 1989, Kikuzawa 1995), enabling it to replace damaged leaves rapidly.

The most rapid growing species, cedar gum was able to develop the highest leaf area of any of the species over two growing seasons (Table 5), a contributory factor being that it retains leaves for more than one growing season. Other factors contributing to the high productivity are the long period of photosynthetic activity and the high photosynthetic efficiency known of eucalypts, particularly under conditions of high stomatal conductance (Whitehead & Beadle 2004). Furthermore, the opportunistic nature of eucalypt growth is likely to have enabled cedar gum to exhibit a longer growing season than alder.

The leaf area of alder and sycamore were not significantly different (Table 5) and they both exhibit high SLA, yet alder had accumulated a greater stem dry weight (Table 4), due to a longer period in leaf and potentially due to higher leaf nitrogen content, allowing higher rates of photosynthesis (Koike & Sanada 1989). Ash was the slowest growing species, and had the lowest leaf area (Table 5) and the shortest growing period of the four tree species (Figure 3). Another factor may be low rate of photosynthesis in ash; a study by Koike and Sanada (1989) found that ash (*Fraxinus mandshurica*) had a relatively low rate of photosynthesis across soils with a range of levels of soil nitrogen, when compared with alder (*Alnus hirsuta*) and birch (*Betula maximowicziana*).

There were several limitations to this study. The method used to harvest leaves at the end of the growing season was unsuitable for ash and measuring leaf area at one point in time ignored potential turnover of leaves during the growing season. The approach used to measure growing season did not allow the estimation of growing season for cider gum, an evergreen tree. This prevented a comprehensive comparison of the trees surviving the first winter. If the study was repeated the root and branch biomass would also be estimated to enable calculation of the LAR. Furthermore, growing season and leaf area was assessed in two different years and it would have been more consistent to measure these in the same year.

Conclusion

It is clear that there are significant differences in growth between the tree species tested at this trial. By the end of the second growing season all shining gum, the fastest growing species had been killed by the cold winter of 2009-2010. Of the surviving three species, the species with greatest stem volume after two growing seasons was cider gum followed by alder (Table 3). After three growing seasons none of the original eucalypts survived and alder exhibited the largest stem volume, although the specific gravity of the wood was less than the other species (Table 4). Cider gum had developed a particularly high LA after two growing seasons, with alder and sycamore following with LA that were not significantly different (Table 5). However, alder had grown much more quickly and its longer growing season (Figure 3) may contribute to this higher growth rate. LAR (stem weight) was particularly low for ash (Figure 2), indicating that ash allocates less relative resources to leaves than to its stem. SLA was also low for ash, as well as cider gum indicating that these species invest relatively high resources in each unit area of leaf area, relative to alder and sycamore (Figure 2). The strong influence of LA and growing season on productivity was shown by creating a resource capture efficiency by multiplying growing season by LA, and this explained 56% of the variation in stem dry weight between trees (Figure 4), however this was only marginally better than LA alone (Figure 5). It is likely that growing season would have a greater influence on growth if origins had been selected from a wider range of latitudes.

The results show that for biomass production on similar sites, alder would be a good candidate, being capable of rapidly accumulating LA and exhibiting a long growing season, resulting in high productivity. However, studies have shown that stands of alder emit N₂O, which may possibly make it less attractive as a candidate tree for sequestration of greenhouse gas (Arnold et al. 2005, Mander et al. 2008).

List of abbreviations

L: length of leaf
LA: leaf area
LAR: leaf area ratio
LMF: leaf mass fraction
LDW: leaf dry weight
M: oven dry weight of stem
RGR: relative growth rate
S: length of petiole and main midrib in ash
SG: specific gravity
SLA: specific leaf area
V: stem volume
W: maximum width of leaf

Acknowledgements

We would like to acknowledge funding from the University of Cumbria and Alastair Chalmers of Pyro Classic stoves that made the establishment of the field trial possible. We would also like to thank staff and students at the University of Cumbria for their help in establishing and maintaining the trial

and to Graham and Chris Leslie for helping with assessments. Several people have provided useful information for this article; Ted Wooddisse of Nottinghamshire County Council and the late John Purse of Primabio on aspects of eucalypts and Matthew Wilkinson and Jason Hubert of Forest Research who provided methods for monitoring and assessing leaf phenology. The authors would like to acknowledge their important contribution and also that of Dr E. Bakr of the Plant protection Institute in Cairo who kindly provided the Compu Eye software used to measure leaf area. We also thank the two anonymous reviewers for making constructive suggestions for improvement to the original draft article.

References

- AFOCEL (2003) Information Eucalyptus, Itineraire technique et production. Fiche Information Eucalyptus No.2 - Avril 2003, AFOCEL, Nangis, France 4pp.
- Alberti G, Candido P, Peresstti A, Turco S, Piussi P and Zerbi G (2005) Aboveground biomass relationships for mixed ash (*Fraxinus excelsior* L. and *Ulmus glabra* Hudson) stands in Eastern Prealps of Friuli Venezia Giulia (Italy). *Annals of Forest Science* 62: 831-836
- Anon (2011) UK seasonal weather summary, winter 2010/2011. *Weather* 66: 99.
- Antinez I, Retamosa EC, Villar R (2001) Relative growth rate in phylogenetically related deciduous and evergreen woody species. *Oecologia* 128: 172-180.
- Arnold K von, Nilson M, Hinell B, Weslien P and Klemedtsson (2005) Fluxes of CO₂, CH₄ and N₂O from drained organic soils in deciduous forest. *Soil Biology and Biochemistry* 37: 1059-1071.
- Basler D and Korner C (2012) Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology* 165: 73-81.
- Cannell MGR, Sheppard LJ and Milne R. (1989) Light use efficiency and the woody biomass production of poplar and willow. *Forestry* Volume 61 (2): 125-136
- Centre for Ecology and Hydrology (2013) Soil data sheet for the Newton Rigg trial. Centre for Ecology and Hydrology, Lancaster, UK. 1p.
- Claessens H, Oosterbaan A, Savill P and Rondeux J (2010) A review of the characteristics of black alder (*Alnus glutinosa* (L.) Gaertn.) and their implications for silvicultural practices. *Forestry* 83 (2): 163-175.
- Cooke JE, Eriksson ME and Juntilla O (2012) The dynamic nature of bud dormancy in trees: environmental control and molecular mechanisms. *Plant, Cell and the Environment* 35 (10): 1707-1728.
- Cope M, Leslie AD and Weatherall A (2008). The potential suitability of provenances of *Eucalyptus gunnii* for short rotation forestry in the UK. *Quarterly Journal of Forestry* 102 (3): 185-194
- Evans J (1986). A re-assessment of cold-hardy eucalypts in Great Britain. *Forestry* 59 (2): 223-242.
- Forestry Commission (undated) Regions of provenance and seed zones in Great Britain (including zones for indigenous Scots pine.) Forestry Commission, Edinburgh, UK.
[http://www.forestry.gov.uk/pdf/FRMGuidelinesRoPmap.pdf/\\$FILE/FRMGuidelinesRoPmap.pdf](http://www.forestry.gov.uk/pdf/FRMGuidelinesRoPmap.pdf/$FILE/FRMGuidelinesRoPmap.pdf)
Accessed 2 January 2017.

586 Forest Research (no date) Assessing bud burst in rowan. Forest Research Northern Research
587 Station, Roslin, near Edinburgh, UK. 4pp.

588 Hardcastle PD (2006). A review of the impacts of short-rotation forestry. LTS International,
589 Edinburgh, UK [online] Last accessed on 27 June 2007:
590 [http://www.forestry.gov.uk/pdf/SRFFinalreport27Feb.pdf/\\$FILE/SRFFinalreport27Feb.pdf](http://www.forestry.gov.uk/pdf/SRFFinalreport27Feb.pdf/$FILE/SRFFinalreport27Feb.pdf)
591

592 Hubert J and Cundall E (2006). Choosing Provenance in Broadleaved Trees, Forestry Commission
593 Information Note 82. Forestry Commission, Edinburgh, UK 12pp.

594

595 Kikuzawa K (1995) Leaf phenology as an optimal strategy for carbon gain in plants. Canadian
596 Journal of Forest Research 73:158-163.

597

598 Koike T and Sanada M (1989) Photosynthesis and leaf longevity in alder, birch and ash seedlings
599 grown under different nitrogen levels. Annals of Forest Science 46 supplement: 476s-478s.

600

601 Laclau JP, Almeida JC, Gonçalves JL, Saint-André L, Ventura M, Ranger J, Moreira RM and
602 Nouvellon Y (2009) Influence of nitrogen and potassium fertilization on leaf lifespan and allocation of
603 above-ground growth in Eucalyptus plantations. Tree Physiology 29 (1):111-124.

604

605 Leslie AD, Mencuccini M, Purse J and Perks MP (2014) Frost damage to eucalypts in a short-rotation
606 forestry trial in Cumbria (England), iForest Volume 7.

607

608 Mander U, Lohmus K, Teiter S, Keiko U and Augustin J (2008) Gaseous nitrogen and carbon fluxes
609 in riparian alder stands. Boreal Environment Research 13: 231-242.

610

611 Matthews RW and Broadmeadow MSJ (2009) The potential of UK forestry to contribute to
612 Government's emissions reduction commitments, Chapter 8 in Read DJ, Freer-Smith PH, Morison
613 JIL, Hanley N, West CC and Snowdon P (eds) (2009). Combating climate change - A role for UK
614 forests. An assessment of the potential of the UK's trees and woodlands to mitigate and adapt to
615 climate change. The Stationery Office, Edinburgh, UK: 139-162.

616

617 **Met Office (undated a) 1971-2000 Averages.** Met Office, Exeter, UK.
618 <http://www.metoffice.gov.uk/climate/uk/averages/19712000/>. Accessed 23 May 2012.

619

620 **Met Office (undated b) Minimum temperature monthly lowest [deg C] January 1961-1990.** Met Office,
621 Exeter, UK. <http://www.metoffice.gov.uk/climate/uk/averages/19611990/exn/1.cjf>. Accessed 12 June
622 2009.

623

624 **Met Office (undated c) Spring 2011.** Met Office, Exeter, UK.
625 <http://www.metoffice.gov.uk/climate/uk/2011/spring.html>. . Accessed 7 July 2014.

626

627 Perks MP, Harrison AJ and Bathgate SJ (2006) Establishment Management Information System
628 [EMIS]: delivering good practice advice on tree establishment in the uplands of Britain. In Reynolds
629 KM, Ray D and Thomson AJ (eds.) Sustainable forestry: from monitoring and modelling to knowledge
630 management and policy science, CABI International, Oxford, UK pp412-424.

631

632 Poorter H and Evans JR (1998) Photosynthetic nitrogen-use efficiency of species that differ
633 inherently in specific leaf area. Oecologia 116: 26-37.

634

635 Poorter H and Remkes C (1990) Leaf area ratio and net assimilation rate of 24 wild species differing
636 in relative growth rate. Oecologia 83: 553-559.

637

Poorter H, Niklas KJ, Reich PB, Oleksyn J, Poot P and Mommer L (2012) Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *New Phytologist* 193: 30-50.

Prior J and Kendon M (2011) The UK winter of 2009/2010 compared with severe winters of the last 100 years. *Weather* 66: 4-9.

Purse J and Leslie AD (2016a) Eucalyptus – Part 1, species with forestry potential in the British Isles. *Journal of Forestry* 110 (2): 88-97.

Purse J and Leslie AD (2016b) Eucalyptus - Part 2, findings from trial plantings, and silvicultural requirements in the British Isles. *Quarterly Journal of Forestry* 110 (3): 161-168.

Pyatt DG, Ray D and Fletcher J (2001). An Ecological Site Classification for forestry in Great Britain. Bulletin 124. Forestry Commission, Edinburgh, UK.

Serdar U and Demirsoy H (2006) Non-destructive leaf area estimation in chestnut. *Scientia Horticulturae* 108: 227-230.

Stape JL, Blinkley D and Ryan MG (2004) Eucalyptus production and the supply, use and the efficiency of use of water, light and nitrogen across a geographical gradient in Brazil. *Forest ecology and Management* 193 (1-2): 17-31.

Tateno M (2003) Benefit to N₂-fixing alder of extending growth period at the cost of leaf nitrogen loss without resorption. *Oecologia* 137 (3): 338-343

Ugese FD, Baiyeri KP and Mbah BN (2008) Leaf area determination of the shea butter tree (*Vitellaria paradoxia* C.F. Gaertn) *International Agrophysics* 22: 167-170.

Valladares F and Niinemets U (2008) Shade Tolerance, a Key Plant Feature of Complex Nature and Consequences. *Annual Review of Ecology, Evolution, and Systematics* 39: 237-257.

Verwijst T and Wen D-Z (1996) Leaf allometry of *Salix viminalis* during the first growing season. *Tree Physiology* 16: 655-660.

Vitasse Y, Delzon S, Dufrene E, Pontallier J-Y, Louvet J-M, Kremer A and Michalet R (2012) leaf phenology sensitivity to temperature in European trees: Do within species populations exhibit similar responses? *Agricultural and Forest Meteorology* 149 (5): 735-744.

Wargo PM (1978) Correlations of leaf area with length and width measurements of leaves of black oak, white oak and sugar maple. Forest Service Research Note NE-256, Northeastern Forest Experiment Station, USDA Forest Service, Newtown Square, PA, USA. 3pp.

Whitehead D and Beadle CL (2004) Physiological regulation of productivity and water use in Eucalyptus: a review. *Forest Ecology and Management* 193:113-140.

Worrell R (1992) A Comparison Between European, Continental and British Provenances of Some British Native Trees: Growth, Survival and Stem Form. *Forestry* 65 (3): 253-280.

Worrell R (2006) Protocol for assessment for senescence and leaf size in birch. Forest Research, Northern Research Station, Edinburgh.

691 Wright, HL and Andrew IA (1976) Principles of experimental design in Burley J and Wood PJ (eds) A
692 manual on species and provenance research with particular reference to the tropics. Tropical
693 Forestry Papers No 10. Commonwealth Forestry Institute, Oxford, UK p67-81
694
695 Wright IJ and Westoby M (2002) Leaves at low versus high rainfall: coordination of structure, lifespan
696 and physiology. New Phytologist 155 (3): 403-416.
697
698

699 Table 1 Origins of trees used in the trial and recommendations for the origins. A map of regions of
700 provenance (Zones) for Great Britain is available in Forestry Commission (undated).

Species	Origin	Recommendations
Alder	Zone 108, South west Scotland	Use British provenances ¹ .
Ash	Zone 108, South west Scotland	Seed stand material or material slightly to the south of planting site ² .
Birch	Zone 202, central to north east Scotland.	Avoid origins from long distances away from the planting site (slightly southern/ eastern locations seems to give more rapid growth) ² .
Cider gum	Likely to be from a seed stand at Dipton, New Zealand ³ . Original origin unknown.	Origins from Lake McKenzie and Mount Cattley, Tasmania perform particularly well ^{4,5} .
Shining gum	Likely to be from a seed stand at Dipton, New Zealand ³ . Origin Central Victoria	Victoria provenances are most frost hardy ⁴
Sycamore	Zone 403, Midlands, England	Most British provenances grow well at most sites. May increase productivity by using origins from sites slightly to the south of the planting site ² .

701 Recommendations from ¹Worrell (1992) ²Hubert and Cundall (2006), ³Purse pers. comm. 2009a,
702 ⁴Evans (1986), ⁵Cope, Leslie and Weatherall (2008)

703
704
705 Table 2 Climate at the trial from the Establishment Management Information System¹ and from 1971
706 to 2000 average data from the Newton Rigg weather station² (Met Office undated a, except for
707 minimum temperature, which is from Met Office undated b)

AT5 ¹ (degrees yr ⁻¹)	CT ¹ (°C)	DAMS ¹	MD ¹ (mm)	Summer Rainfall (mm) ¹	Winter Rainfall (mm) ¹	Mean Min Temp (°C) ²	Min Temp (°C) ²	Frost days ²	Mean Max Temp (°C) ²
1503.4	6.3	14	148.2	386.4	396.2	0.4 (Jan)	-14°C	57.6	19.4 (Jul)

708 AT5 = Accumulated temperature above 5°C, CT=continentality, DAMS = Detailed aspect method of scoring, MD = moisture
709 deficit. A detailed description of these variables is found in Pyatt et al (2001). Annual rainfall = summer rainfall + winter
710 rainfall.
711
712

713 Table 3 Height, stem diameter and volume for each species after two and three growing seasons.

Species	2 growing seasons			3 growing seasons		
	Height ¹ (cm)	Diameter ² (mm)	Stem volume ² (cm ³)	Height ³ (cm)	Diameter ³ (mm)	Stem volume ² (cm ³)
Alder	155.5a	28.7a	9.6a	194.5a	43.8a	23.0a
Ash	119.0b	20.7b	6.4b	133.8b	27.4b	9.5b
Cider gum	199.5c	35.7c	18.0c	-	-	-
Sycamore	134.0ab	14.3b	4.9b	162.0ab	20.0b	8.1b

714 ¹Anova and Tukey’s test (Mean shown), ²Kruskal Wallis and Mann Whitney U test (Median shown),
715 ³natural logarithm transformed then Anova and Tukey’s test (Mean shown). The same lower case
716 letter (a, b or c) denotes no significant difference between species.
717
718

719 Table 4 Specific gravity (SG) and calculated stem dry weight (M) after two growing seasons from 12
720 randomly selected trees of each species.

Species	Specific gravity (g cm ⁻³ = Mg m ⁻³)	Stem dry weight (g)
Alder	0.391a	3.759a
Ash	0.563b	3.539ab
Cider gum	0.553c	9.896c
Sycamore	0.497c	2.390b

721 Kruskal Wallis and Mann Whitney U test applied to specific gravity and stem dry weight (Medians
722 shown). The same lower case letter (a, b or c) denotes no significant difference between species.

723 Table 5 Description of the models predicting leaf area where y is mean area of one leaf (LA) in cm²
 724 and x is L (cm) x W (cm) of the leaf, except for ash where x is leaf stalk length and median leaf area
 725 by species. Total tree LA was calculated by multiplying number of leaves by the mean area of one
 726 leaf and converted in m².

Species	Number leaves	Regression model	R ²	SEE	Median tree LA (m ²)
Alder	202a	$y=0.325x^{1.102}$	0.941	0.202	0.1919a
Ash	22b	$y = 0.1201x^{2.1891}$	0.707	0.524	0.0627b
Sycamore	25b	$y= 0.532x^{1.021}$	0.964	0.197	0.1856a
E. gunnii (mature)	657c	$y=0.052x^2+0.448x+1.032$	0.967	0.947	0.4999c
E. gunnii (juvenile)		$y=0.7714x^{0.943}$	0.881	0.216	

727 Table 6 Description of the models predicting leaf area where y is LDW of a leaf in grammes and x is
 728 L (cm) x W (cm) of the leaf, except for ash where x is leaf stalk length. Whole tree LDW was
 729 calculated by multiplying number of leaves by the mean dry weight of a leaf.
 730

Species	Regression model	R ²	SEE	Median tree LDW (g)
Alder	$y=0.054+0.001x+0.0000751x^2-0.000000292x^3$	0.967	0.041	21.75a
Ash	$y = 0.004x^2 + 0.005x - 0.029$	0.853	0.187	12.44a
Sycamore	$y=0.007-0.20x$	0.970	0.099	23.63a
E. gunnii (mature)	$y=0.010x+0.001x^2+0.029$	0.981	0.017	100.98b
E. gunnii (juvenile)	$y=0.012x+0.021$	0.932	0.300	

731
 732

733 Figure 1 Approach used to determine significance of differences between variables.
734
735 Figure 2 Leaf Area Ratios (LAR) and Specific Leaf Areas (SLA) for the four tree species (Different
736 letters above the bars indicates a significant difference, confidence intervals are not given because of
737 lack of normality).
738
739 Figure 3 Bud burst and leaf retention of ash, sycamore and alder over the growing season of 2011.
740
741 Figure 4 Stem dry weight against resource capture efficiency.
742
743 Figure 5 Stem dry weight against leaf area.
744
745
746
747