

Dalgarno, Gordon (2008) Therapy with music precisely tuned to the individual's own frequencies as indicated by muscle responses. In: Research Fest 2008, July 2008, University of Cumbria, UK. (Unpublished)

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/419/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
 - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

Therapy with music precisely tuned to the individual's own frequencies as indicated by muscle responses.

Abstract

This work arose from my research to find people's personal frequencies and to improve the effectiveness of therapy with music by tuning prominent notes in the music to match these frequencies.

These are early results and must be considered tentative, until confirmed with more subjects and under better controlled conditions.

In my work in *Attune to Health*, applying music in a high tech way for therapy, I originally used the methods of Applied Kinesiology to assess relative muscle tone, in response to music to assess whether the music seemed to be suitable therapeutically. Then almost accidentally, and to my surprise I found that, muscle tone was very strong at certain narrowly defined frequencies, and conversely very weak at slightly off these frequencies. In fact the frequency had to be correct to within 0.1% which corresponds to about 1/60th of a semitone. This tiny difference of pitch cannot be heard – it would need to be about 10 times that to be easily discernable under ordinary conditions – and yet certain muscles in the body have this large response.

I then found there is not just one frequency for a person but at least 11 and that these are spaced in a regular geometric pattern which is not related in any way to (at least Western) musical intervals.

Some very different pieces of music appear to show the same frequencies for a person, however, when correctly “tuned in” the relative strength of other muscles, e.g. individual fingers, appear to vary with the emotional affect of the music

There is also a time dependent effect. If one applies a slightly “wrong” frequency, 1/60th of a semitone too low, for say 2 minutes, the relevant muscle shows weak and it remains weak for perhaps 20 seconds after the “correct” frequency is applied. However if one instead of applying the ‘correct’ frequency, one applies one 1/60th of a semitone too high, the muscle goes strong within about 2 seconds; Naturally one must not leave this “wrong frequency” on for more than 10 seconds or so because it would cause the muscles to go weak.

Introduction

This work arose from my research to enhance therapy with music. My earlier work had been concerned with enabling the music to be felt through the whole body through a structured vibroacoustic system which incorporated a new design of transducers [1]. I believed that while useful, this was not enough and it could be enhanced further if a way could be found to find people's personal frequencies. It would then be relatively simple to tune prominent notes in the music to match these

frequencies.

The belief in people's personal frequencies is widespread. Checking in 8 languages there are sayings such as "He and I are on the same wavelength", "I am tuning into her", "I am finding his notes", "I am resonating with her." (Remember that $\text{frequency} = k/\text{wavelength}$). Most of us have had experiences of being in close proximity to a person we don't know and having a feeling of either being comfortable or uncomfortable. I have had a personal experiences of the great health benefits of music, which, without

my knowledge and in fact only discovered many years later, was tuned to my own frequencies. It was because of these benefits that I was motivated to find a way of helping others in this way.

It is universally accepted that one has to use the "right" music for the client and the occasion. The "right" music is not generally the clients favourite music but it has to be something which they like and which greatly moves them emotionally [2]. Without an appropriate choice any of the above attempts to enhance the therapeutic effect would be as naught.

Some year ago I trained as a practitioner of Applied Kinesiology and I believed I could use this to help select the appropriate music to use.

Applied Kinesiology

At the point its necessary to explain a little of the practice of Applied Kinesiology. The part of the practice which is diagnostic makes extensive use of testing specific muscles according to procedures to determine whether the muscle can resist a force applied in a specific way. The basis being that, for example, a substance to which the body is sensitive or allergic will make certain muscles go 'weak', while certain vitamins or other substances which the body needs will make specific muscles go 'strong'. The terms 'strong' and weak' and are a little misleading and are simply shorthand for ability to resist a specifically applied force. For more information see [3].

For the results to be valid, one has to first carry out some neurological tests and if necessary adjustment.

Unfortunately some unscrupulous salespeople are using what they call kinesiology so show clients that they should buy the products, without such tests, often with invalid procedures and sometimes the inappropriate muscles. Such demonstrations give the subject a bad name.

The purpose of this article is not to justify Applied Kinesiology and I ask you to believe, that despite being a subjective method. when properly carried out the results are reliable

I investigated whether I could apply these methods to find whether the music being used was 'right', and found that I that could. Initially the muscles which I tested were the pectoralis major clavicular (which is linked to the stomach meridian) and the latisimus dorsii (which is liked to the spleen meridian). Both of these are of course connected with emotions. Sometimes it was found to be appropriate to use a procedure in kinesiology known as the *relevance test*.

Of course kinesiology can only help to find the appropriate piece; one can only test with what one has, in this case the available music, in my case a selection of CDs which are believe to be therapeutic. Hence a wise selection of possible pieces is important, as clearly one cannot be testing too many in a therapy session.. If the client has an idea if which pieces are therapeutic for them, I strongly advice the client to bring these pieces with them.

An accidental discovery –application to finding personal frequencies.

While carrying out the above work ie to assess relative muscle tone, to determine whether the music seemed to be suitable therapeutically, ie noticed that on some pieces the muscle tone was extremely high whiles another recording of the same piece it was not [4] This might of course have been due to different musical interpretation and different instrumentation or voices, some of which might be disagreeable to the listener, but I also was alert to the possibility that it might sometimes be due to different tuning.


Investigated further by using pieces which were piano only, and which seems to have a similar tempo and interpretation. The same result was obtained, greatly strengthening the likelihood of it being connected with the frequency of tuning.

Investigating using 'variable pitch' playback

I began by using a variable pitch tape recoding deck which I had available. I used recording of piano pieces which had been found therapeutic and played them back at various pitches.

While it is generally called 'variable pitch' it is in fact variable frequency. Pitch is a subjective measurement while frequency is absolute and the play back is absolute in term of the variation in the % change in Hz from that at which it was recorded. EG if it was recorded with the tuning for 'A' of 438 Hz and the 'pitch' wheel is set to -1.0% than it would play back as if it had been recorded at A = 433.62

This quickly showed that the frequency of tuning hypothesis seemed valid: at certain spot setting of the pitch adjustment control the muscles would go extremely strong while just a factional movement of the control knob would cause then to go extremely weak while further away from the spot frequency they were normal. This is shown diagrammatically below.



The adjustment to find the spot was very fine – a ‘hairs breadth on the pitch adjustment knob, and being an analogue system, never designed for such a precise purpose, the position would not be reproducible hour to hour, or capable of being calibrated absolutely to a sufficient precision. Nevertheless it well served its purpose in finding that there were spot frequencies (I thought two) at which the muscles would exhibit this extremely sharp ‘tuning’ effect.

Further, and most important, I was able to show that it was not just an effect on the muscles. Some musical people found the music so disagreeable to listen to when it was just slightly off a “correct” setting for them, that they asked for it to be switched off.

Funding was obtained to buy a digital system based system, based on a ‘Tascam MD-CD1’. This allowed the frequency to be varied by + or – 16% to a precision of 0.1%. which correspond to about 1/60 of a semitone. (A semitone difference in frequency is about 6%)

To my surprise the fineness of setting required to exhibit the muscle effect was even greater than I had thought, corresponding to about the maximum resolution to which the machine can be set, ie about 1/60 of a semitone.

Indeed it would be desirable to have a machine with double the precision of setting because occasionally it seems the best setting is between two adjacent settings.

It seems likely that the reason why this phenomena has not been found before, is that unless one was looking for it one would not find it, due to the extreme narrowness of the frequency setting – within 1/60 of a semitone.

A number of personal frequencies – and their strange spacing.

Using this equipment, it became apparent that a person has considerable more personal frequencies than I first thought. With the range of the equipment there are 11 spot frequencies and typically they are spaced as below around which ever central frequency is strong. Actual figures depend on the individual person and the piece of music but the pattern is always similar. (The figures of 0.05% come from estimating between two settings of the frequency setting control.)


Frequency settings for client: Self			PIECE
Equally			Cents
tempered			100.0
semitone			
	tone		200.0
In just			Cents
intonation			
,			
Larger tone (e.g. C to D in key			203.9
of C)			
Smaller tone (e.g. D to E in key			182.4
of C)			
C to Db in key of C			111.7
C to C# in key of C			92.2
D to D# in key of C			70.7
D to Eb in key of C			90.2

The following table is another way of showing that these spacings of these spot frequencies do not correspond to (at least Western) musical scales.

Note	Scale Type			Just Intonation (Cents)	Pythagoras (Cents)	Equal Temp (Cents)
	Just Intonation	Pythagoras	Equal Temp			
C	1.000	1.000	1.000	0.00	0.00	0.00
C#	1.055	1.068	1.059	92.18	113.69	100.00
Db	1.067	1.053	1.059	111.73	90.22	100.00
D	1.125	1.125	1.122	203.91	203.91	200.00
D#	1.172	1.201	1.189	274.58	317.60	300.00
Eb	1.185	1.185	1.189	294.13	294.13	300.00
E	1.250	1.266	1.260	386.31	407.82	400.00
F	1.333	1.333	1.335	498.04	498.04	500.00
F#	1.406	1.424	1.414	590.22	611.73	600.00
Gb	1.422	1.405	1.414	609.78	588.27	600.00

G	1.500	1.500	1.498		701.95	701.95	700.00	
G#	1.563	1.602	1.587		772.63	815.64	800.00	
Ab	1.580	1.580	1.587		792.18	792.18	800.00	
A	1.667	1.688	1.682		884.36	905.86	900.00	
A#	1.758	1.802	1.782		976.54	1019.55	1000.00	
Bb	1.778	1.778	1.782		996.09	996.09	1000.00	
B	1.875	1.898	1.888		1088.27	1109.77	1100.00	
C'	2.000	2.000	2.000		1200.00	1200.00	1200.00	

Having seen that the spot frequencies do not correspond we can graph them as below so see if we can recognise any pattern.



My research work continues to find the explanation for this pattern and also for the extremely sharply tuned muscle response.

Variations between people and the phenomena of entrainment

Some people have frequencies so similar that one wonders whether they are indeed “on the same wavelength”.

Remember that $\text{wavelength} = k / \text{frequency}$

| |Freq for client: BA |

| |PIECE |Mozart |Piano |Conc. |Num |21 |Movemt | |Slow | | | | |Freq | |L5 |L4 |L3 |L2 |L1 | |F0 | |H1
|H2 |H3 |H4 |H5 | |% devi | |-14.3 |-7.6 |-4.5 |-2.4 |-1.3 | |0.0 | |1.5 |2.6 |4.8 |7.8 |14.5 | |

Initially I was finding that many of my friends and acquaintances were on very similar frequencies to me and I thought that meant that “they were on the same wavelength”. But there were too many similar for this to be a sufficient explanation, and it became even more obvious that this must be rang when I began to measure the frequencies of clients who were ill. The explanation was entrainment.

Entrainment: This is well know and understood phenomena in physics. For example if a number of pendulums, with approximately the same period are swinging in a room, they will soon all come into synchronicity.

If there are two pendulums and one is much heavier then the other which is swing with a slightly longer period, the lighter will speed up to a period almost the period of the heavier, while the heavier will slowdown very slightly (imperceptibly if it is much heavier) to the common period.

It was found that if a person approached me, usually within 3 feet our frequencies would change, with the ill person’s frequencies being pulled towards mine. For friends and acquaintances, it could go either way. Some people did not affect me at all, a few did as far away as 5 feet. In almost all cases touching the person produced no further frequency shift. A few people coming within 3 feet of me had no proximity effect but then when they touched me, a similar shift took place.

The above information may be valuable in the field of investigating unspoken communication.

There is a procedure in kinesiology involving holding what are known as the ESR point on the head, which can minimise or possibly remove such interactions, but whether that remove it completely in this case I don’t know.

I have been able to investigate the interaction between people to some extent by this means and also by using 3 way testing.

Time dependant effects

There is also a time dependent effect, and one needs to be aware of it and allow for it in this work. If one applies a slightly “wrong” frequency, 1/60th of a semitone too low, for say 2 minutes, the relevant muscle shows weak and it remains weak for

perhaps 20 seconds after the “correct” frequency is applied. However if one instead of applying the ‘correct’ frequency, one applies one 1/60th of a semitone too high, the muscle goes strong within about 2 seconds; Naturally one must not leave this “wrong frequency” on for more than 10 seconds or so because it would cause the muscles to go weak.

The need for an objective method of measurement and one which does not require being close to the subject

Because of this interaction between people, I feel it is very important to obtain an objective method of measuring the muscles without coming into proximity with the subject. Further this reason is, I believe, even more important than the objection that the technique of measurement is subjective, because with a person trained in kinesiology I believe the results are reliable. I have seen suitable equipment at Teesside University hospital but the cost was prohibitive for this work at about £20,000.

Further refinement on assessing the emotional effect on the muscles of the music

This work is at an early stage and needs to be confirmed.

To investigate this one first needs to be exactly on the person’s frequency. I noticed that the pectoralis major clavicular (which is linked to the stomach meridian) and the latisimus dorsii (which is linked to the spleen meridian) often gave slightly different answers and I hypothesises that this was due to different emotional effects. It seems that the best muscle to use for this purpose of setting the frequency is the piriformis. Having set the frequency precisely one can look at the secondary effects on these two muscles due to emotions.

I have also been assessing the relative strengths of the fingers on the left hand, again once the frequency is set precisely. For example, in Handel’s ‘Messiah’, the index finger is stronger in the 3 people I have tested in the ‘Halleluia Chorus’ and the little finger stronger in the ‘Amen’. In these 3 the effect is strong, and reproducible.

Why not use simple sounds (e.g. sine waves) instead of music for therapy?

A very fair question. I have also trained in sound therapy using computer generated sounds but I don’t practice it because of the inadequate results I have obtained. It true that some practitioners obtain good results but this may be down to them as individuals with special qualities.

I think the reason is that the brain quickly tires of listening to a simple sound and pays little attention after a short time, whereas music is a highly complex stimulus.

Nevertheless I think it would be a good idea add specific sounds to the music and I intend to pursue this, probably using low bass notes. If necessary, they could be tuned slightly differently from the existing bass notes and perhaps an octave lower, the ear-brain probably would not notice the slight mistuning (because pitch discrimination is much poorer with low pitch notes and because beat frequencies would be outside the audible range.)

Note

These are early results and must be considered tentative, until confirmed with more subjects and under better controlled conditions. These results are based on tests with only 31 people and these are far from a cross section of the population, being friends, acquaintances and clients.

Two people have independently verified some of my results without my involvement but using my equipment, so it is not entirely independent.

References

- [1] G. Dalgarno . “A vibroacoustic couch to improve perception of music by deaf people and for general therapeutic use.” 6th International Conference on Music Perception and Cognition. Keele University, ISBN 0-9539909-0-7 UK. 2000
- [2] G. Dalgarno “A Scientific Approach, including the use of Voice Measurements, to Sound and Music Therapy.” Positive Health, Sept 2005
- [3] See www.kinesiology.com and www.kinesiologyuk.com
- [4] G. Dalgarno. “Enhancing Therapy with Music.” TRAHMA conference held at Cumbria University, July 08

Acknowledgements

I would like to thank Dr Ian Fabian BA, MS, PhD, MB, ChB for many helpful discussions and practical help with experiments.

I am grateful to Community Champions and the Galbraith Trust for funding.

Gordon Dalgarno BSc (Hons. Physics), M.I.O.A., Adv. Cert. Kinesiology,
Reiki Master (Usui & Seichem), A.A.M.T. EFT Cert Practitioner
Former Research Fellow in Music and in Electronics at the University of York
Former Research Fellow in Music Psychology at Keele University

Glengarry Clinic, 3 Thornton Grove, Morecambe, LA4 5PU.
E-mail: AttuneToHealth@yahoo.co.uk Tel 01524 427919 Mobile 07903-367978