

Dr Nicola Relph^a; Dr Katie Small^b

^a*Corresponding Author*

Faculty of Health and Social Care

Edge Hill University

St Helens Road

Ormskirk

L39 4QP

0144695657056

Nicola.Relph@edgehill.ac.uk

^b University of Cumbria

Department of Medical and Sport Sciences,

Carlisle, UK.

The Influence of 9-marathons completed in 9 days on injury incidence and selected musculoskeletal tests

Key Points:

- Flexibility, proprioception and balance performance progressively decrease during a multi-day running event.
- Athletic trainers should design recovery and injury prevention strategies to be used with-in competition based on with-in event data collection.
- Future studies should adopt this novel with-in competition data collection method to appreciate the dynamic nature of musculoskeletal physiology.

Key words: Multi-day running; athletic therapy; musculoskeletal testing.

21 **Abstract**

22 Multi-day running events are increasingly popular however, research in these events is lacking
23 and fails to consider the dynamic nature of musculoskeletal physiology. Twenty-three athletes
24 completing a ten-day marathon event participated in the study. Proprioception, dynamic
25 balance, knee valgus and flexibility were assessed the day before the event and after one, five
26 and nine consecutive marathons. There were significant reductions in these measurements
27 across the event and reductions were more apparent in the non-dominant side. Each runner
28 suffered on average 4.2 injuries. Runners performed significantly worse in musculoskeletal
29 measurements, particularly on the non-dominant side, as the competition progresses.
30 Therefore, athletic trainers should design appropriate between-day recovery strategies during
31 events based on with-in event data collection.

33 **Introduction**

34 Ultra-distance running events continue to increase in popularity¹. The distance of these events
35 is greater than 26.2 miles with total running times typically over six hours, occurring over
36 multiple days². It is perhaps not surprising that there are higher injury rates in these events
37 relative to shorter running distances³, running of this nature may cause repeated stresses on
38 tissues in a fatigued state⁴. Indeed, injuries occurred in between 60% and 100% of all
39 competitors in previous research on ultra-distance events⁵⁻⁷. Furthermore, in a 7-day running
40 event 84.8% of 396 runners required medical attention, which was equivalent to 3.8 injuries
41 per runner or 7.2 injuries per 1000 hours³. However, there is a lack of research investigating
42 musculoskeletal physiology changes that may inform athletic practice during multi-day
43 running events.

It is important for practitioners to know how the musculoskeletal system changes to design effective with-in event treatments⁸. A dynamic approach to injury prevention has been developed in a model of etiology in sport injury⁹. Importantly, this incorporates repeated exposure to events that may lead to musculoskeletal adaptations⁹. This cyclical approach is one that is often missed in literature with clinical measurements being taken only once, before the race occurs; for example, static measures of alignment¹⁰. However, it is important to consider how the body may adapt across a performance taking the dynamic nature of injury occurrence into consideration.

There is very little research on the changes in musculoskeletal physiology during a multi-day running event⁵. One important measurement is neuromuscular control which has been linked to efficient running technique¹¹. Poor proprioceptive ability (the ability to perceive position, movement and force of the limbs during running¹²) may be linked to changes in running performance as the central nervous system may not receive effective afferent information (feedback mechanisms) and hence prepare the correct muscle activity for impending perturbations (feed-forward mechanisms)¹². Dynamic balance relies on good neuromuscular control and poor ability is possibly linked to running injuries due to excessive knee valgus positions from poor hip adductor strength during running¹³⁻¹⁵. An increased or decreased range of motion has previously been cited as a potential risk of injury¹⁶⁻¹⁷ due to a potential increase in compressive and tensile stress on lower-limb joints¹⁸. Therefore, it is important these measures are monitored across a multi-day event to identify when musculoskeletal physiology changes and if athletic therapy is required.

The lower-extremity is the most common location for running injuries in multi-day events¹⁹. Therefore, the current study considered knee joint position sense (JPS), lower-limb dynamic balance, knee neuromuscular control during single-leg landing and hip and ankle flexibility

adaptations during a multi-day event. The aim of this study was to evaluate the effect of running one, five and nine consecutive marathons on musculoskeletal physiology using in-competition data collection methods.

Methods

Participants

A total of 23 athletes (age 44.7 ± 7.59 years, mass 75.1 ± 12.99 kg, self-reported weekly mileage 43.3 ± 12.67 miles) participated in this prospective cohort study. The event involved completing 10 marathons in 10 consecutive days on the same course, however data was collected after day zero (D0), marathon one (M1), five (M5) and nine (M9). Table 1 describes participant characteristics. All participants provided voluntary, written informed consent and the rights of the participants were protected. The study was ethically approved by the University Review Board (ref: DC/SB 15/19).

Instrumentation

Knee JPS was collected using a camera (Casio Exilim, EX-FC100, Casio Electronics Co., Ltd. London, UK) on a fixed, level tripod. Dynamic balance was collected using a “Y symbol” taped to the floor (see figure 1) and tape measure. Knee neuromuscular control data was collected using a camera (Casio Exilim, EX-FC100, Casio Electronics Co Ltd, London, UK) mounted on a fixed, level tripod. Flexibility was collected using a 30cm goniometer (Baseline® Evaluation Instruments, White Plains, NY) and tape measure.

Tasks

Reliability statistics for tasks are reported in Table 2.

89 *Knee JPS*

90 Markers were placed on a point on a line following the greater trochanter to the lateral femoral
91 epicondyle, the lateral femoral epicondyle and the lateral malleolus on the dominant leg. The
92 athlete was seated on the end of a treatment table and blindfolded. The leg was passively moved
93 by the experimenter through 30°–60° of extension from a starting knee angle of 90° or 60°–
94 90° of flexion from a starting angle of 0° at an approximate angular velocity of 10°/s (see figure
95 2). The target angle was held by the athlete for 5s before the researcher returned the limb to the
96 starting position. The athlete then replicated the target position. Knee positions were captured
97 using photographs and digitising software (Kinovea, v0.8.15, Joan Chamant & Contrib, 2006–
98 2011). The delta score between target and reposition angles was taken as the absolute error
99 score in degrees and averaged across five trials. The protocol has been validated against an
100 isokinetic dynamometer²⁰.

101 *The Modified Star Excursion Balance Test (SEBT)*

102 Dynamic stability was measured using the modified SEBT. Briefly, this task involves single
103 leg squats, with the non-weight bearing leg reaching maximally towards anterior, posterior-
104 medial and posterior-lateral directions along a designated line and then returning to the start
105 position of single leg stance. Further detail of this protocol can be found in Munro and
106 Herrington²². Each runner completed four trials in each direction on both legs and results were
107 normalised to leg length.

108 *Knee Neuromuscular Control*

109 Knee neuromuscular control was measured using maximum knee valgus angle during single-
110 leg-landing. Markers were placed on the mid-point of the femoral condyles, the mid-point of
111 the ankle malleoli and the anterior superior iliac spine on both legs. The athlete stepped forward

from a 30-cm height, dropping as vertically as possible landing in a single-leg stance and holding for 3s, completing three trials on each leg. A trial was void if the non-stepping leg touched either the step or the floor during the task. Knee valgus angles were measured as the greatest angle between the line from the ASIS to the patella and the patella to the ankle marker during the landing performance using digitising software (Kinovea, v0.8.15, Joan Chamant & Contrib, 2006-2011) and the average taken. This task is correlated to forward running technique²⁵.

Lower-Limb Flexibility

Two experienced athletic trainers took flexibility measurements on both legs with consistent roles in each protocol. The pelvis was stabilised to avoid compensatory movements in hip measurements. The flexibility of the iliotibial band was collected using the Ober's protocol²⁶. Hip adductor flexibility measurements were taken in a supine position, the goniometer arms placed in-line with the contralateral anterior-superior iliac spine and the anterior mid-line of the ipsilateral femur. The runner actively performed maximal hip abduction and the value on the goniometer was recorded. Ankle dorsiflexion flexibility was measured using the knee to wall protocol detailed in Powden, Hoch and Hoch²⁸. The runner completed the test in a tandem stance, with the tibia progressing over the talus and the heel remaining fully on the ground until the knee touches the wall. Internal and external rotation of the hip was measured following the Bullock-Saxton and Bullock protocol²⁹. The runner was in prone and the knee passively flexed by the first examiner to 90°. The stationary arm of the goniometer was positioned parallel to the testing surface and the moving arm was placed along the tibia. The additional examiner then palpated the opposite posterior-superior iliac spine and the original examiner passively externally or internally rotated the limb. The measurement was taken at the point before the pelvis began rotating.

136 *Procedures*

137 Data was collected in a sports clinic between May 2015 and June 2016. The number of injuries
138 per athlete was recorded by two athletic trainers three times daily (further details of injury data
139 is reported elsewhere³⁰). An injury was defined as a specific musculoskeletal abnormality that
140 the runner perceived to affect performance⁵.

141 *Statistical Analysis*

142 Normality was checked using the Shapiro-Wilk test, if confirmed, the means and standard
143 deviations of parametric measures were calculated. One-way repeated measures ANOVAs
144 were used to explore the main effect of time and then post-hoc Bonferroni comparisons were
145 utilised when required to complete multiple (six) comparisons. Non-normal data was presented
146 as medians and interquartile ranges and analysed using Friedman's ANOVA and Wilcoxon
147 Signed Rank tests. The significance level was accepted at $p \leq 0.05$.

148 **Results**

149 In total 73%, 50%, 69% and 70% of the sample completed all knee JPS, SEBT, knee valgus
150 and flexibility testing respectively (see Table 3). Post-hoc analysis revealed no significant
151 differences between completers and drop-outs for baseline measures of each tests. Non-
152 parametric data (medians) and parametric data (means) is presented in Table 4.

153 *Knee JPS*

154 There were no effects of time on knee JPS into extension but was into flexion. JPS ability
155 improved from D0 to M1 by 2.4° and reduced from M0 to M5 by 1.3° . On one or more
156 occasions in the event 85% of runners demonstrated JPS difference scores above the SDD into
157 flexion.

158 *Dynamic Balance*

159 Results of the modified SEBT did not alter during the competition on the dominant leg for
160 anterior and posterior-lateral reach directions. However, posterior-medial reach distances
161 reduced from D0 to M5 by 15% of leg length and from D0 to M9 by 19% of leg length (both
162 above the SDD) and 80% of participants produced results above the SDD.

163 Anterior reach distance, posterior-medial reach distance and posterior-lateral reach distance all
164 reduced on the non-dominant side. The anterior reach distances reduced by 9% of leg length
165 from D0 to M9 (above the SDD). The posterior-medial reach distance reduced by 5% of leg
166 length from D0 to M1 (below the SDD) and by 13% of leg length from D0 to M9 (above the
167 SDD). The posterior-lateral reach distances reduced by 12% of leg length from D0 to M9
168 (above the SDD). 70% of runners produced differences greater than the SDD for anterior reach,
169 90% for posterior-medial reach data and 80% for posterior-lateral reach data.

170 *Knee Neuromuscular Control*

171 Knee neuromuscular control did not significantly change during the event for either the
172 dominant or non-dominant side, indeed 77% of runners did not demonstrate difference values
173 over the reported SDD.

174 *Lower-Limb Flexibility*

175 Adductor flexibility and ankle dorsiflexion flexibility on the dominant side of the body
176 reduced. Adductor flexibility reduced from D0 to M5 by 5.6° and M9 by 10.8°. This flexibility
177 also reduced from M1 to M9 by 11°. 91% of runners' data exceeded the reported SDD. Ankle
178 dorsiflexion flexibility was reduced from D0 to M1 (by 1.41cm), M5 (by 2.65cm) and M9 (by
179 3.40cm). There was also a significant reduction in this flexibility between M1 and M9 (by

2.00cm). 87% of runners had differences above the SDD during the event. The remaining flexibility measurements on the dominant side did not change.

All flexibility measures on the non-dominant side significantly reduced. ITB flexibility reduced from D0 to M9 (by 0.98cm) and from M1 to M9 (by 0.50cm). Flexibility of the adductor muscles again reduced from D0 to M9 (difference 9.1°) and M5 (difference 8.2°). This flexibility also significantly decreased when comparing M1 to M5 (difference 7.4°) and M9 (difference 8.3°). 87% of runners' data was above the reported SDD. A similar pattern was evident in ankle dorsiflexion flexibility, there were differences between D0 and M5 (difference 2.72cm) and M9 (difference 4.6cm). Ankle dorsiflexion flexibility was also worse after M9 compared to M1 (difference 3.6cm) and marathon five (difference 1.8cm). For this measurement 83% of runners displayed differences above the SDD. Internal hip rotation on the non-dominant side significantly increased between D0 and M9 (difference 6.0°). External hip rotation also significantly reduced over time from D0 to M5 by 7.6° and M9 by 10.5°. 78% of runners displayed differences greater than the SDD for hip rotation measures.

There were 4.2 injuries per runner; 89% of injuries involved the lower extremity; 24.1% in the foot, 18.5% the hip/buttock, 16.7% the ankle and 16.7% in the lower leg.

Discussion

The aim of this study was to measure the effects of a multi-day running event on knee proprioception, dynamic balance, knee neuromuscular control and flexibility. The results suggest these measures, particularly on the non-dominant side, decrease in performance from D0 to M5 and again to M9 during the event.

Knee Proprioception

There was an initial improvement in knee JPS into flexion from D0 to M1, but this difference was below SDD values²¹. However, knee JPS into flexion reduced from D0 to M5 and this difference was above SDD values²¹. This suggests knee JPS may not reduce after running one marathon but could be impaired after five marathons. To our knowledge this is the first paper to consider JPS ability during a multi-stage running competition. However, previous research has reported a reduction in JPS ability following treadmill running to fatigue³¹⁻³². Three theories have been proposed to explain the mechanisms behind this finding; impaired excitation of motor units³³, increase in knee laxity³⁴ and increase in pain³³. All explanations suggest the afferent signalling used by the CNS to process JPS information is disrupted with fatigue, therefore making this process unstable and increasing errors³⁵. Knee flexion occurs in running from initial touch down to mid-swing phases³⁶. If the runner is unable to correctly perceive the position of their knee this could lead to errors in efferent signalling used for movement preparation. The results of the current study suggest knee JPS ability into flexion reduces after completion of five marathons. Therefore, athletic trainers may incorporate proprioceptive exercises after five days of running.

Dynamic Balance

Dynamic balance significantly reduced in all reach directions on the non-dominant limb and the posterior-lateral direction on the dominant limb from D0 to M9 and all average differences were above SDD values²². Again, to the author's knowledge, this is the first study to measure dynamic balance ability during a multi-day running event. However, the findings from the current study support previous literature that reported a decrease in balance performance following shorter running activities; for example, Steib³⁷ reported a decrease in SEBT performance following treadmill running to exhaustion. Other authors used different methods of balance measurement to present a reduction in ability with fatigue³⁸⁻³⁹. A reduction in

dynamic balance has been suggested to potentially increase the risk of running injuries due to a loss in neuromuscular control in lower extremity joints^{31, 40}. The results of dynamic balance in the non-dominant leg got progressively worse across the event with the biggest performance decrease from D0 to M9. This has important implications for the timing of prevention and treatment strategies during an ultra-endurance event, athletic trainers should introduce dynamic balance exercises with-in ultra-running events.

Knee Neuromuscular Control

There were no significant changes to knee neuromuscular control on either leg. This is an unexpected finding however, Munro²⁴ stated the SDD as 7.54°-7.90° for the task and the greatest differences in this study were below the SDD. Therefore, 2D manual digitisation may not be sensitive enough to identify changes in knee valgus angle.

Lower Limb Flexibility

Increased flexibility may be desirable for optimal running performance⁴¹. The flexibility of the adductor muscles and ankle dorsi-flexors significantly reduced on both the dominant and non-dominant sides during the event. All adductor differences were above the reported SDDs apart from dominant leg, D0 to M5. Poor hip adductor flexibility has been linked to reduced stability at the hip and knee joint during gait and increased risk of ITB syndrome⁴². The flexibility of the adductor ankle dorsi-flexors also significantly reduced on both legs and differences were above the SDD except between D0 and M1 on the dominant leg. A reduction in ankle dorsi-flexion may change running mechanics at the ankle, specifically in preparation for foot strike; if the ankle is less flexed, this can modify foot strike patterns and lower-limb absorption mechanics and hence increase ground reaction forces⁴³. Reduced ankle dorsi-flexion has been

linked to injuries to the knee⁴⁴ and foot⁴⁵ due to an increase in force transmitted along the kinetic chain.

The non-dominant limb also had reduced flexibility in the remaining measurements. Hip internal rotation increased, and external rotation decreased in flexibility across the competition and all differences were above the SDD. A modification in hip internal movement has been associated with modified knee kinematics that may possibly be linked to injury⁴⁶. Poor hip control can lead to reduced neuromuscular control lower in the kinetic chain and potentially an increased risk of injury⁴⁶. Furthermore, a reduction in ITB flexibility may potentially cause patello-femoral pain⁴⁷ and ITB syndrome⁴⁸. These results suggest athletic trainers should consider flexibility recovery strategies after each day of a multi-day running event, particularly on the non-dominant side.

Limitations

Fatigue was not measured objectively, however, previous research has demonstrated fatigue will be present during ultra-marathon events⁴⁹. Reliability estimates were taken from prior studies. However, knee JPS measures were taken by the same assessor from the reliability study. Also, there is over a decade's worth of reliability literature on both flexibility and SEBT measurements. The dropout levels should also be acknowledged; however, appropriate statistical analysis was used based on the assumption of normality.

Clinical Implications

The results of this study suggest musculoskeletal physiology performance worsens after five days of marathon running and by nine days this may be significant. Athletic trainers should design individual interventions based on in-event testing that runners can perform both before and during events that target flexibility, knee neuromuscular control and dynamic balance.

272

273 *Future Research*

274 The Meeuwisse model¹¹ of injury prevention states risk of injury is cyclical, hence is event
275 and time dependent, therefore, it is recommended that the in-event data collection design
276 should be utilised in further work with larger sample sizes.

277

278 *Conclusion*

279 Multi-day running events can cause over four injuries per runner and musculoskeletal
280 physiology measures worsen progressively across competitions. Athletes should be aware of
281 the potential changes that will occur and prepare appropriately. Importantly, these
282 modifications became more apparent during the competition; these findings would not have
283 been identified if traditional research designs that do not take measurements within competition
284 had been used. Hence athletic trainers should consider in-event measurement with a view to
285 prescribe recovery strategies that incorporate this knowledge (i.e. balance and flexibility
286 recovery methods) in competition.

287 *Acknowledgements*

288 The authors would like to thank Brathay Trust for their assistance in recruiting runners to
289 participate in this research study and to all the runners themselves for their commitment to the
290 study.

291

292 **References**

293 1. van der Worp MP, ten Haaf DSM, de Wijer A et al. Injuries in runners; A systematic review
294 on risk factors and sex differences. PLoS ONE 2015;10:e0114937.

- 295 2. Knechtle B, Duff B, Schulze I et al. Anthropometry and pre-race experience of finishers and
296 nonfinishers in a multistage ultra-endurance run – Deutschlandlauf 2007. *Percept Mot Skills*
297 2009;109:105-118.
- 298 3. Krabak BJ, Waite B & Schiff MA. Study of injury and illness rates in multiday ultramarathon
299 runners. *Med Sci Sports Exerc* 2011;43:2314-2320.
- 300 4. Hreljac A. Impact and overuse injuries in runners. *Med Sci Sports Exerc* 2004; 36:845-849.
- 301 5. Bishop GW & Fallon KE. Musculoskeletal injuries in a six-day track race: Ultramarathoner's
302 ankle. *Clin J Sport Med* 1999;9:216-220.
- 303 6. Fallon KE. Musculoskeletal injuries in the ultra-marathon: the 1990 Westfield Sydney to
304 Melbourne run. *Br J Sports Med* 1996;30:319-323.
- 305 7. Graham SM, McKinley M, Connaboy CC et al. Injury occurrence and mood states during a
306 desert ultramarathon. *Clin J Sport Med* 2012; 22:462-466.
- 307 8. Borland ML, Rogers IR. Injury and illness in a wilderness multisport endurance event.
308 *Wilderness Environ Med.* 1997;8(2):82–88
- 309 9. Clansey A, Hanlon M, Wallace E, et al. Effects of Fatigue on Running Mechanics Associated
310 with Tibial Stress Fracture Risk. *Med Sci Sports Exerc* 2012;44:1917-1923.
- 311 8. Bahr R, Krosshaug T. Understanding injury mechanisms: a key component of preventing
312 injuries in sport. *Br J Sports Med* 2005;39:324-329.
- 313 9. Meeuwisse WH, Tyreman H, Hagel B et al. A dynamic model of etiology in sport injury:
314 The recursive nature of risk and causation. *Clin J Sport Med* 2007; 17:215-219.

- 315 10. Hespanhol Junior LC, De Carvalho ACA, Costa LOP et al. Lower limb static alignment
316 characteristics are not associated with running injuries in runners: Prospective cohort study.
317 Eur J Sport Sci 2016;16:1137-1144.
- 318 11. Glofcheskie GO & Brown SHM. Athletic background is related to superior trunk
319 proprioceptive ability, postural control, and neuromuscular response to sudden perturbations.
320 Hum Mov Sci 2017;52:74-83.
- 321 12. Switlick T, Kernozek TW & Meardon S. Difference in joint-position sense and vibratory
322 threshold in runners with and without a history of overuse injury. J Sport Rehabil 2015;24:6-
323 12.
- 324 13. Baltich J, Emery CA, Stefanyshyn D et al. The effects of isolated ankle strengthening and
325 functional balance training on strength, running mechanics, postural control and injury
326 prevention in novice runners: design of a randomized controlled trial. BMC Musculoskelet
327 Disord 2014;15:407-418.
- 328 14. Francis, P., Gray, K. and Perrem, N. The Relationship Between Concentric Hip Abductor
329 Strength and the Performance of the Y-Balance Test (YBT). International Journal of Athletic
330 Therapy and Training. 2018;23:42-47.
- 331 15. Munro A, Herrington L. & Comfort P. The Relationship Between 2-Dimensional Knee-
332 Valgus Angles During Single-Leg Squat, Single-Leg-Land, and Drop-Jump Screening Tests. J
333 Sport Rehabil. 2017;26(1):72-77.
- 334 16. Shanley E, Kissenberth MJ, Thigpen CA, Bailey LB, Hawkins RJ, Michener LA, Tokish
335 JM, Rauh MJ. Preseason shoulder range of motion screening as a predictor of injury among
336 youth and adolescent baseball pitchers. J Shoulder Elbow Surg. 2015 Jul;24(7):1005-13

- 337 17. Winkelmann ZK, Anderson D, Games KE, Eberman LEJ. Risk Factors for Medial Tibial
338 Stress Syndrome in Active Individuals: An Evidence-Based Review. *Athl Train*. 2016
339 Dec;51(12):1049-1052.
- 340 18. Mann KJ, Edwards S, Drinkwater EJ, Bird S. A Lower Limb Assessment Tool for Athletes
341 at Risk of Developing Patellar Tendinopathy. *Medicine & Science in Sports & Exercise* 2013;
342 45(3):527-533.
- 343 19. Scheer BV & Murray A. Al Andalus ultra trail: An observation of medical interventions
344 during a 219-km, 5-day ultramarathon stage race. *Clin J Sport Med* 2011;21:444-446.
- 345 20. Relph N & Herrington L. Criterion-related validity of knee joint-position-sense
346 measurement using image capture and isokinetic dynamometry. *J Sport Rehabil*
347 2015;Technical Notes(10).
- 348 21. Relph N & Herrington L. Interexaminer, intraexaminer, and test-retest reliability of
349 clinical knee joint-position-sense measurements using an image-capture technique. *J Sport*
350 *Rehabil* 2015;Technical Notes(12).
- 351 22. Munro AG & Herrington LC. Between-session reliability of the star excursion balance test.
352 *Phys Ther Sport* 2010;11:128-132.
- 353 23. Gribble PA, Hertel J & Plisky P. Using the star excursion balance test to assess dynamic
354 postural-control deficits and outcomes in lower extremity injury: A literature and systematic
355 review. *J Athl Train* 2012;47:339-357.
- 356 24. Munro A, Herrington L, Carolan M. Reliability of 2-dimensional video assessment of
357 frontal-plane dynamic knee valgus during common athletic screening tasks. *J Sport Rehabil*.
358 2012;21(1):7-11

- 359 25. Atkin K, Herrington L, Alenezi F, Jones P & Jones R. The Relationship Between 2d Knee
360 Valgus Angle During Single Leg Squat (SLS), Single Leg Landing (SLL), And Forward
361 Running. Br J Sports Med 2014 48: 563.
- 362 26. Gajdosik RL, Sandler MM, & Marr, HL. (2003). Influence of knee positions and gender on
363 the ober test for length of the iliotibial band. Clin Biomech 2003;18:77–79.
- 364 27. Cejudo A, Ayala F, De Baranda PS & Santonja F. Reliability of Two Methods of Clinical
365 Examination of the Flexibility of the Hip Adductor Muscles. Int J Sports Phys Ther. 2015;
366 10(7): 976–983.
- 367 28. Powden CJ, Hoch JM & Hoch MC. Reliability and minimal detectable change of the
368 weight-bearing lunge test: A systematic review. Man Ther. 2015;20(4):524-32
- 369 29. Bullock-Saxton JE & Bullock MI. Repeatability of Muscle Length Measures around the
370 Hip. Physiotherapy Canada 1994; 46(2): 105-109.
- 371 30. Small K & Relph N. Musculoskeletal Injury Rates in Multiday Marathon Runners
372 Performing Ten Consecutive Marathons on a Repeat Course. Journal of Athletic Enhancement
373 2017; 6 (5).
- 374 31. Baharlue J & Khayambashi K. The effect of sub-maximal and maximal running on knee
375 joint position senses and static balance in healthy young male athletes. World Journal of Sport
376 Sciences 2012;6:242-246.
- 377 32. McMullen KL, Cosby NL, Hertel J. et al. Lower Extremity Neuromuscular Control
378 Immediately After Fatiguing Hip-Abduction Exercise. J Athl Train 2011;46:607-14.

- 379 33. Miura K, Ishibashi Y, Tsuda E, et al. The effect of local and general fatigue on knee
380 proprioception. *Arthroscopy* 2004;20:414-418.
- 381 34. Hutton RS & Nelson DL. Stretch sensitivity of Golgi tendon organs in fatigued
382 gastrocnemius muscle. *Med Sci Sports Exerc* 1986;18:69-74.
- 383 35. Fortier S & Basset FA. The effects of exercise on limb proprioceptive signals. *J*
384 *Electromyogr Kinesiol* 2012;22:795-802.
- 385 36. Dugan, S. A., & Bhat, K. P. Biomechanics and analysis of running gait. *Physical medicine*
386 *and rehabilitation clinics of North America*, 2005;16(3):603-621.
- 387 37. Steib S, Zech A, Hentschke C et al. Fatigue-induced alterations of static and dynamic
388 postural control in athletes with a history of ankle sprain. *J Athl Train* 2013;48:203-208.
- 389 38. Burdet C & Rougier P, Effects of utmost fatigue on undisturbed upright stance control. *Sci*
390 *Sports* 2004;19:308-316.
- 391 39. Lepers R, Bigard AX, Diard JP et al. Posture control after prolonged exercise *Eur J Appl*
392 *Physiol Occup Physiol*. 1997;76:55-61.
- 393 40. Wright KE, Lyons TS & Navalta JW. Effects of exercise-induced fatigue on postural
394 balance: a comparison of treadmill versus cycle fatiguing protocols. *Eur J Appl Physiol*
395 2013;113:1303.
- 396 41. Overmoyer GV, Reiser RF 2nd. Relationships between lower-extremity flexibility,
397 asymmetries, and the Y balance test. *J Strength Cond Res*. 2015;29(5):1240-7.

42. Louw M, Deary C. The biomechanical variables involved in the aetiology of iliotibial band syndrome in distance runners - A systematic review of the literature. *Phys Ther Sport*. 2014;15(1):64-75
43. Carter J & Greenwood M. Does Flexibility Exercise Affect Running Economy? A Brief Review. *Strength and Conditioning Journal* 2015 37(3):12-21
44. Malliaras P, Cook JL, Kent P. Reduced ankle dorsiflexion range may increase the risk of patellar tendon injury among volleyball players. *J Sci Med Sport*. 2006;9(4):304-9.
45. Wearing SC, Smeathers JE, Urry SR, Hennig EM, Hills AP. The pathomechanics of plantar fasciitis. *Sports Med*. 2006;36(7):585-611.
46. Bedi A, Warren RF, Wojtys EM, Oh YK, Ashton-Miller JA, Oltean H, Kelly BT. Restriction in hip internal rotation is associated with an increased risk of ACL injury. *Knee Surg Sports Traumatol Arthrosc*. 2016;24(6):2024-31.
47. Hudson Z, Darthuy E. Iliotibial band tightness and patellofemoral pain syndrome: a case-control study. *Man Ther*. 2009;14(2):147-51
48. Noehren B, Schmitz A, Hempel R, Westlake C, Black W. Assessment of strength, flexibility, and running mechanics in men with iliotibial band syndrome. *J Orthop Sports Phys Ther*. 2014;44(3):217-22.
49. Millet GY. Can Neuromuscular Fatigue Explain Running Strategies and Performance in Ultra-Marathons?: The Flush Model. *Sports Med* 2011;41:489-506.

419

420

421

422

423

424

For Peer Review




Figure 1. Modified Star Excursion Balance Test (SEBT) set-up
215x279mm (150 x 150 DPI)




Figure 2. Example of Knee Joint Position Sense (JPS) data collection set-up
665x563mm (144 x 144 DPI)

Table 1. Participant Characteristics (mean±SD unless stated).

Flexibility Testing (n=23)	
Age (years)	44.7±7.59
Mass (kg)	75.1±12.99
Gender (males/females)	16/7 (number)
Knee JPS and Neuromuscular Control (n=13)	
Age (years)	46.8±5.03
Mass (kg)	72.3±13.02
Gender (males/females)	9/4
Dynamic Balance (n=10)	
Age (years)	42.0±9.61
Mass (kg)	78.8±12.66
Gender (males/females)	7/3 (number)

Table 2. Reliability statistics

Protocol	Test-retest reliability (ICC)	Smallest Detectable Difference (SDD)
Knee Joint Position Sense ²¹	Knee Flexion 0.92 Knee Extension 0.86	Knee Flexion 1.10° Knee Extension 1.67°
Star Excursion Balance Test ²²⁻²³	0.84-0.92	Anterior reach 6.87% leg length Posterior-medial reach 8.15% leg length Posterior-lateral reach 7.11% leg length
Knee Neuromuscular Control ²⁴	Men 0.80 Women 0.82	Men 7.54° Women 7.90°
Ober’s Protocol ²⁶	0.82-0.92	Not available
Hip Adductor Flexibility ²⁷	0.92-0.99	5.1° - 7.6°
Ankle Dorsiflexion Flexibility ²⁸	0.99	1.57cm
Internal Hip Rotation ²⁹	0.98	4.29°
External Hip Rotation ²⁹	0.99	6.11°

Table 3. Drop out data (absolute number of runners at each time phase).

Risk Factor	Day zero	Marathon 1	Marathon 5	Marathon 9	Completion
Flexibility	23	23	18	16	70%
Knee JPS	13	12	11	8	73%
Knee Neuromuscular Control	13	12	10	9	69%
SEBT	10	10	6	5	50%

Table 4. Mean±SD measurements for parametric data and Median [IQR] measurements for nonparametric data at day zero (D0), marathon one (M1), five (M5) and nine (M9) for parametric data. SEBT = Star Excursion Balance Tests. *p≤0.05, **p≤0.01, ***p≤0.001.

Variable	D0	M1	M5	M9	Main Effect p-value	Significant Post-hoc analysis
Knee Joint Position Sense Extension (°)	3.6±1.9	3.7±1.8	3.3±2.3	2.8±1.2	0.65	N/A
Knee Joint Position Sense Flexion (°)	4.7 [3.3]	2.9 [2.2]	3.3 [2.9]	3.3 [3.4]	0.03	D0 v M1** M1 v M5*
Dominant Leg SEBT Anterior (% of leg length)	85±8.3	85±5.0	80±11.5	75±6.6	0.15	N/A
Dominant Leg SEBT Posterior Medial (% of leg length)	101 [18.0]	90 [11.2]	86 [13.6]	82 [17.6]	0.01	D0 v M5* D0 v M9*
Dominant Leg SEBT Posterior Lateral (% of leg length)	86±10.7	84±14.5	80±14.2	79±11.4	0.42	N/A
Non-Dominant Leg SEBT Anterior (% of leg length)	86 [12.4]	83 [11.6]	80 [8.7]	77 [6.4]	0.05	D0 v M9*
Non-Dominant Leg SEBT Posterior Medial (% of leg length)	96±8.6	91±10.4	84±10.9	83±7.4	0.004	D0 v M* D0 v M9*
Non-Dominant Leg SEBT Posterior Lateral (% of leg length)	86 [15.4]	85 [12.7]	82 [19.1]	74 [118.6]	0.02	D0 v M9*
Dominant Leg Knee Valgus (°)	5.8±2.0	5.5±3.1	6.5±5.4	6.9±5.1	0.57	N/A
Non-Dominant Leg Knee Valgus (°)	7.6±4.9	8.6±3.5	5.9±4.7	7.8±3.7	0.85	N/A
Dominant Leg Iliotibial Band Flexibility (cm)	14.9±1.7	14.8±1.8	15.3±1.8	15.6±2.0	0.21	N/A
Non-Dominant Leg Iliotibial Band Flexibility (cm)	14.6±1.9	15.1±1.7	15.6±1.8	16.8±2.0	0.001	D0 v M9** M1 v M9*

Dominant Leg Ankle Dorsiflexion Flexibility (cm)	11.0±3.1	9.6±3.3	8.4±3.3	7.6±2.9	0.001	D0 v M1** D0 v M5 *** D0 v M9 *** M1 v M9*
Non-Dominant Leg Ankle Dorsiflexion Flexibility (cm)	10.5±3.0	9.6±3.7	7.8±2.8	5.9±2.5	0.001	D0 v M5*** D0 v M9*** M1 v M9** M5 v M9**
Dominant Leg Internal Hip Rotation (°)	31.8±9.5	35.3±9.7	34.7±9.4	34.1±5.4	0.32	N/A
Dominant Leg External Hip Rotation (°)	29.8±11.4	30.5±13.5	30.4±9.6	29.1±8.5	0.24	N/A
Non-Dominant Leg Internal Hip Rotation (°)	31.1±9.9	32.9±10.6	34.3±7.4	37.1±6.5	0.01	D0 v M9**
Non-Dominant Leg External Hip Rotation (°)	34.7±11.6	31.8±12.3	27.2±8.0	24.2±6.9	0.001	D0 v M5* D0 v M9*