

A review of the suitability of eucalypts for short rotation forestry for energy in the UK.

Leslie, A.D¹.; Mencuccini, M².; Perks, M.P³ and Wilson, E.R⁴.

¹Senior Lecturer, National School of Forestry, University of Cumbria, Ambleside Campus, Rydal Road, Ambleside, LA22 9BB, UK. Andrew.leslie@cumbria.ac.uk. +44 1539 430291

²Professor, CREAF, University Autonomous Barcelona, Campus de Bellaterra, Edifici C 08193 Cerdanyola del Vallès, Barcelona, Spain.

³Project Leader, Centre for Forestry and Climate Change, Forest Research, Northern Research Station, Roslin, Midlothian EH25 9SY, UK.

⁴Walsh Fellow in Silviculture, University College Dublin Forestry, Belfield, Dublin 4, Ireland

Abstract

Eucalyptus has been identified as a genus with potential for short rotation forestry in the UK. This article assesses the suitability of *Eucalyptus* for biomass production. The first part of the article compares *Eucalyptus nitens* and *Eucalyptus gunnii* against short rotation forestry (SRF) species proposed by Hardcastle (2006), while the second part discusses limitations to the growing of eucalypts in the UK and how they may be overcome.

Eucalypts compare favourably with other tree species in the UK in terms of rapid growth (up to $30 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$) over short rotations of ten to fifteen years. The only genus that is potentially as productive in the UK is *Nothofagus*. Furthermore, most species will readily coppice, enabling regeneration after damage and avoiding the costs of replanting. The wood characteristics compare positively with other SRF species, exhibiting a moderate wood density, but limitations are a relatively high moisture and chlorine content. Many of the SRF species listed in Hardcastle (2006) are now damaged or under threat from damage by exotic pests or diseases. Eucalypts are currently relatively free from such damage.

It is cold temperatures that most limits the use of eucalypts in the UK. Eucalypts, particularly when young are vulnerable to damage from cold weather events, particularly when temperatures drop rapidly. However, the risk can be reduced by planting appropriate species and provenances, facilitating rapid growth as smaller trees are more vulnerable and by focusing on species that coppice following damage.

Key words:

Eucalyptus, United Kingdom, short rotation forestry, biomass

Acknowledgements

The authors would like to thank Professor Julian Evans and Dr Bill Mason for encouraging the writing of this review, originally as part of a PhD thesis. They would also like to acknowledge the assistance of Dr James Morison of Forest Research, the two anonymous referees and the editor for providing constructive comments on the draft manuscript.

Introduction

In the United Kingdom (UK) there are two main aims of the Government's Renewable Energy Strategy (DECC 2012); to reduce emissions of carbon dioxide and to improve energy security. This is to be achieved through producing 15% of the energy in the UK through renewable means by 2020, which represents an increase of seven times the 2009 contribution within a decade (DECC 2012). The lead scenario generated within the Strategy suggests that 30% of electricity and 12% of heat could be provided through use of renewable sources of energy (DECC 2012). By 2014, 7% of energy was derived from renewable sources, of which 72% was obtained from bioenergy (DECC 2015). Of this approximately 40% was from wood or other plant biomass (DECC 2015). The transition from fossil fuels to renewable sources of energy has been supported by the Renewable Heat Incentive (RHI) and the Feed-In Tariff (FIT). The RHI is focussed on providing payments to encourage the production of heat from renewable energy sources (OFGEM 2019a), while the FIT is aimed at increasing renewable electricity generation (OFGEM 2019b). The FIT scheme is closing to most applications in April 2019 but the RHI continues.

As a source of renewable energy, biomass has certain attractions; it can produce energy at times of peak demand, it involves low fossil fuel inputs for production and tried and tested technology is available for its efficient conversion. The UK Bioenergy Strategy (DECC 2012) predicted that while biomass imports will form the bulk of supply, domestic production can provide an important, stable and reliable resource. If demand for biomass were to be met domestically, the area under energy crops would need to increase dramatically. The UK Biomass Strategy (DEFRA 2007) anticipated that 350,000 ha of perennial energy crops would be required by 2020, which contrasts with the 2009 area of 15,500 ha of Short Rotation Coppice (SRC) and *Miscanthus* (SAC 2009). Woody biomass is an attractive option compared with agricultural biomass crops as it requires the input of relatively low levels of fertiliser, pesticides and herbicides and can also be established on marginal land, thereby not competing with food production (Hastings et al. 2014).

Two approaches to bio-energy production using tree species have been adopted: short rotation coppice (SRC) and short rotation forestry (SRF) (Read et al. 2009). These two approaches are compared in Table 1. Since the 1990s the development of dedicated woody energy crops in the UK has focused on using SRC of clones of willow (*Salix* spp.) or poplar (*Populus* spp.). However, a comparative analysis of both systems identified SRF using fast growing tree species as being the most cost-effective and time-efficient approach to reducing greenhouse gases (Matthews and Broadmeadow 2009). However, there are very few examples of SRF in the UK, with the largest plantation an area of 24.2 ha of eucalypts established at Daneshill in Nottinghamshire, eastern England as an energy forest (Wooddisse¹ pers comm). SRF contrasts with conventional production forestry in several ways.

Conventional production forestry uses conifers, which are more productive than broadleaves

¹ Land Management Officer, Nottinghamshire City Council

1 in the UK over traditional rotations of 40-70 years. Normally SRF does not include thinning
2 during the rotation, whereas after canopy closure production forestry stands are thinned on a
3 cycle of 5 -10 years. Due to the short rotation, measures are taken to accelerate establishment
4 and maximise growth such as highly intensive weed control and fertiliser application (Purse
5 and Leslie 2016b).
6
7
8
9
10

11 The silvicultural attributes of an ideal SRF tree crop are described in Table 2. Ideally it
12 should grow rapidly, with mean annual increment peaking early, have a low environmental
13 impact and the wood should have ideal properties of a fuel, such as low moisture content and
14 appropriate chemical composition. A review by Hardcastle (2006) identified species and
15 genera with potential for SRF in the UK, these being ash (*Fraxinus excelsior*), alder (*Alnus*
16 *glutinosa*), birch (*Betula pendula*), *Eucalyptus gunnii*, *Eucalyptus nitens*, *Nothofagus* spp,
17 poplar (*Populus* spp.) and sycamore (*Acer pseudoplatanus*). Of these ash, alder, birch and
18 some poplars are tree species native to the UK. Since the publication of Hardcastle's (2006)
19 report, the introduction of exotic pathogens has reduced the viability of some species, e.g. ash
20 dieback (*Hymenoscyphus fraxineus*) on ash (Thomas 2016).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 In the UK there has been research to identify site suitability criteria for eucalypts through an
39 understanding of the limitations to survival and growth (e.g. Evans 1980, Evans 1986, Purse
40 and Richardson 2001, Leslie et al. 2011, Leslie et al. 2012). However, the scale of planting
41 of eucalypts in the UK has been modest; in the five years between 2011 and 2016 nurseries in
42 the UK have sold about 220,000 seedlings (Purse and Leslie 2016b), which represents less
43 than 100 ha of establishment. As such, there remains uncertainty about the risks of large-
44 scale planting of *Eucalyptus* SRF in the UK.
45
46
47
48
49
50
51
52
53
54
55

56 In this review paper, we compare eucalypts with other appropriate species or genera as a
57 source of biomass for energy production then identify the limitations of eucalypts in the UK
58
59
60
61
62
63
64
65

1 and potential ways in which they may be overcome. There was found to be limited literature
2 on short rotation forestry in general in the UK. The review has therefore relied on compiling
3 information from two sources. The first is literature on cold-tolerant eucalypts that has been
4 collected by the authors over a period of fifteen years and includes material no longer on-line.
5 The information on other species, has relied on extracting relevant material from publications
6 covering more general aspects of the silviculture, growth and wood properties of the tree
7 species. A constraint to comparing growth of SRF species is that there were few data for trees
8 grown under intensive short rotation management in the UK. Whilst yield models have been
9 developed for commercial stands under longer rotations these only provide estimates of stand
10 growth after canopy closure (from 10 to 25 years of age) depending on the maximum mean
11 annual increment (Edwards and Christie 1981).
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 This review is timely as recent research studies (eg Harrison 2011, Leslie et al. 2014b,
28 McEvoy 2016, Leslie et al. 2018) and reviews (eg McKay 2011, Leslie et al. 2012, Purse and
29 Leslie 2016a, Purse and Leslie 2016b) have led to a better understanding of the potential of
30 eucalypts but also their limitations. The findings are focused on the UK, but are relevant to
31 other areas with cool oceanic temperate climates in western Europe.
32
33
34
35
36
37
38
39
40
41
42
43

44 ***Yield of eucalypts and other SRF species***

45
46
47
48
49

50 The main attraction of eucalypts is their high productivity, being among the most productive
51 trees in the UK. Volume growth rates have been estimated from small plot trials, to be as
52 high as 30 m³ ha⁻¹ y⁻¹ for *E. nitens* (Purse and Richardson 2001). In Ireland a plantation of *E.*
53 *nitens* felled at 16 years of age, with 740 stems ha⁻¹, yielded a mean annual increment (MAI)
54 of 26.1 m³ ha⁻¹ y⁻¹ (Hutchinson et al. 2011). Historic Forestry Commission data provides
55
56
57
58
59
60
61
62
63
64
65

evidence of MAIs of $16 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$ for *E. gunnii* over a twenty year rotation (Leslie et al. 2018). Purse and Richardson (2001) conclude that higher yields of 10-15 oven dry Mg $\text{ha}^{-1} \text{ y}^{-1}$ (approximately $20\text{-}30 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$) over 8 to 10 year rotations are possible from plantations of *E. gunnii*. In Redmarley, Gloucestershire in the south of England (Figure 1), *E. gunnii* was estimated to have grown at a MAI of $25 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$ over a 11 or 12 year rotation (Purse and Richardson 2001). The coppice from this stand was assessed at 10 years of age and the standing volume of mainly *E. gunnii* with some *E. dalrympleana* combined was $317 \text{ m}^3 \text{ ha}^{-1}$ or $31.7 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$ with 4,746 stems ha^{-1} (McKay 2010). Furthermore, when *E. gunnii* and *E. nitens* that had been damaged or killed by extreme cold in the winter of 2010-2011 was harvested at Daneshill in Nottinghamshire in the north of England (Figure 1) at 5 years old an average of 85 Mg were extracted per hectare giving a green weight production of $17 \text{ Mg ha}^{-1} \text{ y}^{-1}$ (Wooddisse pers comm).

While eucalypts can be highly productive in the UK yields vary considerably across sites due to edaphic and climatic conditions, the quality of silvicultural practices and the genotype of the planting stock (Kerr and Evans 2011). This was highlighted by a study of four trials from the 1980s that demonstrated the difficulties in consistently achieving high levels of productivity (Kerr and Evans 2011). It was only at one trial at the New Forest (Figure 1), in southern England where the potential of fast growth of eucalypts was realised; biomass at seven years of age of *E. gunnii* and *E. glaucescens* was at least three times that of the other species planted, which were *Alnus cordata*, *Nothofagus obliqua*, *Populus balsamifera* and *Pteryocarya x rehderiana*. This trial was planted at two spacings and the results showed that increasing stocking significantly increased yield over a seven year rotation. For *E. gunnii* a MAI of $2.7 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$ was achieved at 2.8 m spacing and of $13.9 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$ at 1.4 m spacing. For *E. glaucescens* the respective MAI was $8.1 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$ and $17.4 \text{ m}^3 \text{ ha}^{-1} \text{ y}^{-1}$.

SRF species should exhibit a peak in mean annual increment (MAI) at a relatively young age.

The potentially most productive genus other than *Eucalyptus* is *Nothofagus* and UK yield models predict maximum mean annual increments (MMAI) of 10 to 18 m³ ha⁻¹ y⁻¹ (Tuley 1980). The mid MMAI of this range for *Nothofagus* shows MAI peaking at 14.0 m³ ha⁻¹ y⁻¹ at 29 years of age (Tuley 1980). Poplars have been used in short rotation coppice due to their rapid early growth (Mitchell et al. 1993) and have been predicted to produce yields of 30.5 m³ ha⁻¹ y⁻¹ or 22 Mg ha⁻¹ y⁻¹ (at 35% moisture content) at age 15 years (Forest Research 1997). Stokes et al. (2017) has demonstrated the fast growth of hybrid aspen (*Populus x wettsteinii*) and commercial hybrid poplar clones in two Scottish trials at Moray and Kilmichael (Figure 1). An assessment was made after 22 growing seasons at Moray and 21 growing seasons at Kilmichael. Clones of commercial poplar hybrids attained an average height of 19.98 m and 14.15 m respectively and a diameter at breast height (dbh) of 39.98 cm and 24.29 cm. Hybrid aspen exhibited more rapid growth with a height of 17.38 m and dbh of 30.99 cm at Kilmichael. Native aspen (*Populus tremula*) grew more slowly, achieving a height of 10.75 m and dbh of 14.97 cm at Moray and a height of 6.54m and dbh of 8.86 cm at Kilmichael.

A review of the silviculture of alder across Europe showed that current annual increment (CAI) peaks at 20 years and MAI at between 30 and 50 years (Claessens et al. 2010). In Europe sycamore also exhibits an early peak in CAI and MAI and it is described as growing more rapidly than beech (*Fagus sylvatica*) up to an age of around 40 years and as being more productive than ash, on both poor and fertile sites (Hein et al. 2009). In most European countries birch is slower growing than sycamore or ash (Hynynen et al. 2009). The mean and range of MMAI and biomass productivity for these SRF species are presented in Table 3, while biomass productivity is based on SRF rotations, the MAI of species other than the eucalypts is based on conventional silviculture and rotations.

Wood properties and important physiological and morphological characteristics

There are physiological, morphological and wood characteristics that are attractive in a SRF species. The first is the ability to coppice (Dickman 2006, Hinchee et al. 2009, Guidi et al. 2013), as it avoids the costs of replanting and results in enhanced growth rates in the second and subsequent rotations (Pukkala and Pohjonen 1990). All tree species described in Hardcastle (2006) will coppice or sucker, except for *E. nitens* which has a very limited ability to coppice (Neilan and Thompson 2008). While birch coppices after cutting it only does so weakly and so this not a recommended means of regenerating stands (Cameron 1996). For eucalypts the ability to coppice also confers a degree of resilience to damage. During the winter of 2010-11 stems and foliage of *E. gunnii* were killed at a planting at Daneshill, Nottinghamshire in the north of England, however many of the trees later produced coppice shoots and remain healthy to this date, whereas trees of *E. nitens* were killed. This supports the recommendation that in the UK, only species that coppice, such as *E. glaucescens*, *E. gunnii* and *E. rodwayi* (Sims et al. 2009) should be planted, rather than the few species like *E. nitens* that do not have this capability (Neilan and Thompson 2008).

For wood used for solid fuel, rather than conversion into liquid fuels, the main factors determining its suitability are moisture content, heating value, proportion of fixed carbon and volatiles, the ash content and the alkali metal content (McKendry 2002). Moisture content strongly affects the net heating value of wood when it is burned (Huhtinen 2006). There is considerable variation in moisture content between the tree species selected by Hardcastle (2006) (Table 3). Eucalypts have relatively high wood moisture content and experience in Ireland has shown that drying *E. gunnii* can be problematic. The wood only dried rapidly when the bark was removed and this itself was difficult using machinery because of its

1 fibrous nature (Leslie 2013). The heating value of dry wood of different tree species varies
2 relatively little with conifers having a higher value per unit of dry wood than broadleaves,
3
4 due to higher lignin content and the presence of resins (McKendry 2002). However, wood
5
6 density varies considerably between the tree species. Dense wood is an attractive trait for
7
8 fuelwood (Senelwa and Sims 1999), as higher densities represent a higher mass of material
9
10 and therefore energy per unit volume and allows a higher mass of wood to be transported per
11
12 unit volume. Alder and poplar exhibit a low wood density, the eucalypts and *Nothofagus*
13
14 moderate density and sycamore, ash and birch exhibit higher density wood (Table 3).
15
16
17
18
19

20 Many of the species have not been burned for energy on an industrial scale in contrast to
21
22 short rotation coppice. Ash, is known to produce good domestic fuel wood (Savill 2013)
23
24 while in Sweden, birch is widely used as a source of domestic heat (Hedberg et al. 2002). A
25
26 study of the effects of torrefaction, a heating process that improved the quality of wood fuel
27
28 in terms of combustion and gasification on wood from trees including birch, aspen and
29
30 eucalypts showed that the two eucalypt species tested contained chlorine concentrations eight
31
32 times that of aspen and six times that of birch (Keipi et al. 2014). Chlorine can be corrosive
33
34 to boilers and pipework in power plants. High levels of chlorine have also been reported
35
36 from eucalypts grown in Ireland (Teagasc 2015).
37
38
39
40
41
42

43 A regular, straight stem enables more efficient handling, storage and processing (Walker et
44
45 al. 2013), although if the material is to be chipped on site when harvested this is less
46
47 important. Potential SRF species known to exhibit good stem form include *E. nitens* (Neilan
48
49 and Thompson 2008), *Populus* (Savill 2013) and silver birch (Hynynen et al 2009), while the
50
51 stem form of *Nothofagus alpina* is as good as poplar (Tuley 1980). The other SRF species
52
53 show a wide variation in stem form between individuals. Ash is sensitive to frost damage and
54
55 this can result in poor stem form through death of the leader and so frost prone sites should be
56
57
58
59
60
61
62
63
64
65

avoided (Dobrowolska et al 2011). Sycamore shows considerable variation in stem form (Hein et al 2009). Young alder often exhibits a straight stem with a compact pyramidal crown, but stem form becomes more variable as the trees age (Savill 2013). Stem form of *E. gunnii* is variable and often poor (Purse 2010, Marriage 1977), but improved material used in France exhibits good stem form (AFOCEL 2007).

Abiotic and biotic limitations

To be a productive tree for biomass, it must be well adapted to the abiotic and biotic environment. The following sections examine the limitations imposed on eucalypts and the other tree species identified by Hardcastle (2006).

Abiotic limitations

Most of the tree species in Hardcastle's (2006) list of potential SRF species are well adapted to climate of the UK and will grow well on a range of soil types. However, periods of low temperatures can be damaging to *Eucalyptus* spp. and *Nothofagus* spp. (Deans et al. 1992). Eucalypts are at the margins of their climatic limits in most of the UK: over the last decade there have been two winters (2009-2010 and 2010-2011), with extended periods of <-10°C overnight temperatures, causing widespread mortality to *E. gunnii* and *E. nitens* across a range of experimental sites planted in England (Harrison 2011) and in Scotland (McEvoy 2016). However, 2009-2010 was the coldest winter in thirty years and in some parts of England in 100 years (Prior and Kendon 2011) and 2010-2011 was only a little less severe (Met Office 2011). Furthermore, in a review of the impact of climate change on eucalypt plantations in general, Booth (2013) assessed their vulnerability as being moderate. He also noted that the short rotations often associated with eucalypts, compared with conventional forest rotations offered greater opportunities to change genotypes and silvicultural practices.

1 Murray et al. (1986) noted the high risk of *Nothofagus* being damaged: however, if southern
2 provenances are used and the hardiest individuals are selected, they are suitable for planting
3 in most lowland parts of the UK.
4
5
6
7

8 Biotic limitations 9

10
11 The risk of damage to trees from biotic agents is predicted to increase (Logan et al. 2003,
12 Sturrock et al. 2011) and is already having major impacts on forestry in the UK. European
13 larch (*Larix decidua*) is no longer planted due to the impact of *Phytophthora ramorum*
14 (Forestry Commission 2014), while planting of Corsican pine (*Pinus nigra* ssp. *laricio*) has
15 ceased because of damage from *Dothistroma septosporum* (Brown and Webber 2008).
16
17 Wainhouse et al. (2016) provide useful predictions of those pests and pathogens likely to be
18 particularly damaging to trees in the UK in the future while another resource for appraising
19 the risk from insect pests and pathogens is the online UK Plant Health Risk Register (UK
20 Plant Health Risk Register no date). This does not provide a combined overall risk rating for
21 a tree species. However, the risk from pests and pathogens to Hardcastle's (2006) potential
22 SRF tree species can be broadly described as follows.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 There are tree species currently at high risk from damage, such as ash which is no longer
41 being planted due to the predicted damage from ash dieback (Woodward and Boa 2013).
42 Furthermore, an additional risk is that from Emerald ash borer (*Agrilus planipennis*), which is
43 now present in Russia (Straw et al. 2013), with the combined impacts likely to be severe
44 (Thomas 2016). Sycamore can suffer extreme damage by grey squirrels (*Sciurus*
45 *carolinensis*) and planting this species for timber is uneconomic in areas where high squirrel
46 populations are present (Savill 2013). For pathogens of sycamore, Webber et al. (2011) note
47 that *Phytophthora* spp and *Verticilium* wilt can be damaging in nurseries or newly planted
48 stock. *Cryptostroma corticale* also affects sycamore and remains dormant in the tree until it
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 becomes stressed by prolonged dry conditions and the pathogen then causes an ailment
2 known as sooty bark disease, which results in crown dieback and can cause death of the tree
3
4 (Savill 2013).
5
6
7

8 Of the tree species there are those where potentially serious pests or pathogens are already
9 established in Britain. This group includes alder and *Nothofagus* which are at threat of
10 damage from *Phytophthora alni* (Gibbs et al. 1999) and *Phytophthora pseudosyringae* (Scanu
11 et al. 2012) respectively. This group also includes poplars, plantations of which have been
12 severely damaged by rusts (*Melampsora* spp) in the UK (Forestry Commission 2005).
13
14
15

16 Damage by *P. alni* was first noted in Britain 1993 primarily infecting and killing the native
17 alder, but also grey alder (*A. incana*) and Italian alder (*Alnus cordata*) (Gibbs, et al. 1999).
18
19

20 Other recent work (Černý et al. 2012) demonstrates that cold temperatures will kill the
21 pathogen and suggests that with predicted increases in winter temperatures due to climate
22 change, persistence of this pathogen may increase damage.
23
24
25

26 Infection by *P. pseudosyringae* of *Nothofagus* was first noted in 2009 in a stand of *N. obliqua*
27 in Cornwall in the south of England, where in four plots between 50% and 72% of trees had
28 become infected. The susceptibility of *Nothofagus* to this disease prompted Scanu et al.
29 (2012 p27) to comment ‘A consequence of this damaging new disease is that future use of *N.*
30 *obliqua* and *N. alpina* in UK forestry as suitable species for climate change adaptation
31 strategies could be limited’. This view is supported by a recent review of *Nothofagus* in
32 Britain (Mason et al. 2018) which identified *N. obliqua* and *N. alpina* as having potential,
33 provided *P. pseudosyringae* does not prove to be highly damaging. Poplars are susceptible to
34 attack by rusts (*Melampsora* spp). Rusts cause premature leaf fall and can also disrupt
35 hardening in some hosts and other damage can include a reduction in growth, shoot die back
36 and when severe, tree death. Developing varieties of poplar resistant to rusts and to the
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

highly damaging *Xanthomonas populi* that causes stem cankers is the main strategy to produce disease free stands. In the past, the Forestry Commission published a list of resistant varieties (Forestry Commission 2005) and a mix of resistant clones was planted to reduce risk further, however at present there are no fully rust resistant varieties available (Tabbush and Lonsdale 1999).

Birch is currently relatively free of major damaging biotic agents, although it is susceptible to attack by *Armillaria* (Webber et al. 2011). Furthermore, there have been problems of crown dieback reported in recent plantings of birch, in Scotland, due to three pathogens; *Anisogramma virgultorum*, *Marssonina betulae* and *Discula betulina* (Green 2005).

However, it is the threat from a pest, currently absent from the UK, that gives greatest cause for concern. The bronze birch borer (*Agrilus anxius*), if introduced would have a devastating impact (Nielsen et al. 2011) as silver birch and downy birch (*Betula pubescens*) are highly susceptible. Within 8 years of planting in a trial in the USA, all individuals of these birch species had been killed by the borer (Nielsen et al. 2011). The probability of detection of bronze birch borer in wood chips was extremely low using the current protocols in Europe (Okland et al. 2012), although the European and Mediterranean Plant Protection Organisations (2011) risk assessment suggested current measures meant the likelihood of bronze birch borer arriving in Britain is relatively low.

Finally there are the eucalypts which currently are probably at the lowest risk of damage from pests and diseases in the UK as few native pests of eucalypts have been introduced to plantations outside Australia (Fanning and Barrs 2013). The eucalypts identified as being suited to SRF in Britain are not those most susceptible to *Phytophthora* spp or to foliar pathogens (Webber et al. 2011). There are no records of major pest outbreaks in the UK, however there have been outbreaks of pests in Ireland. In the late 1990s a psyllid,

1 *Ctenarytaina eucalypti* was introduced to Ireland (Chauzat et al. 2002). Chemical control was
2 not particularly effective and so a parasitic wasp, *Psyllaephagus pilosus* was introduced and
3 this effectively controlled the psyllid (Chauzat et al. 2002). In 2007 a leaf beetle,
4
5 *Paropsisterna selmani* caused severe defoliation in multi species plantings of eucalypts
6
7 (Fanning and Barrs 2013, Horgan 2012). Fanning and Barrs (2013) describe the beetle as
8
9 being a serious threat to eucalypts in Ireland, the UK and more widely in Europe as the adults
10
11 are strong fliers, capable of surviving long periods without food and can tenaciously cling to
12
13 various forestry residues.
14
15
16
17
18
19

20 **Discussion**

21
22 Eucalypts possess many of the silvicultural attributes that are attractive for SRF for biomass,
23
24 namely rapid early growth, good stem form and methods for propagation and establishment
25
26 that are well understood. This review of evidence supports the proposition that eucalypts
27
28 could provide a productive source of biomass in the UK and in other countries with oceanic
29
30 climates in Europe. In Ireland, well-adapted species of eucalypts have been identified (Neilan
31
32 and Thompson 2008) and spacing and other trials are underway (Tobin et al. 2016). In
33
34 France there has been a programme establishing pulp plantations of cold-tolerant eucalypts
35
36 that has been established for 35 years in the mid-Pyrennes. About 2,000 ha have been planted
37
38 of *E. gunnii* and *Eucalypts x gundal*, its hybrid with *Eucalyptus dalrympleana* (FCBA 2018).
39
40
41
42
43
44
45

46 The climate of the UK however is generally colder than Australia and the parts of France
47
48 where cold-tolerant eucalypts are planted. Over most of the UK absolute minimum
49
50 temperatures dip to below -14°C over a thirty year period (Figure 1), a temperature that will
51
52 damage or kill most eucalypts, particularly young trees or when temperatures have dropped
53
54 rapidly. It is mainly coastal or southern areas where absolute minimum temperatures remain
55
56 higher (Met Office no date). Eucalypts are particularly vulnerable to cold damage because,
57
58
59
60
61
62
63
64
65

1 unlike most temperate broadleaves they do not have a defined dormant period. Growth of the
2 naked buds begins above a certain minimum temperature threshold, which contributes to their
3 high productivity (Beadle et al. 1995). A study of *E. nitens* in Tasmania showed maximum
4 winter growth rates were only marginally less than those in summer (Davidson et al. 1995).
5 Hardening in cold-tolerant eucalypts usually commences below a temperature of between 2°C
6 (Paton 1983) and 4°C (Davidson and Reid 1987) and there are differences between species in
7 the rapidity of hardening and in the minimum temperature they will tolerate before damage.
8 Work by Black (no date) in Ireland developed an index of hardiness for eucalypt species in
9 Ireland based on lethal minimum temperature and rate of hardening and the index ranked
10 species tested in the following order, from most hardy to least hardy; *Eucalyptus rodwayi*, *E.*
11 *glaucescens*, *Eucalyptus subcrenulata*, *Eucalyptus delegatensis*, *E. gunnii*, *E. coccifera* and
12 *E. nitens*. This is a useful approach, although it raises questions as to why *E. gunnii*, a highly
13 cold-tolerant species ranked so low. Details of the provenance used was not provided and it
14 may be that a less cold-hardy one was used.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35 The limited area of both *Eucalyptus* (Purse and Leslie 2016b) and *Nothofagus* (Mason et al.
36 2018) makes matching species to site imprecise but colder and more exposed sites must be
37 avoided. To identify climatically suitable sites for planting *Nothofagus* and *Eucalyptus*, Ray
38 and Sing (2006) created a map of England, Scotland and Wales that showed areas where
39 minimum temperatures of -14°C or below occur more than once every 50 years (Figure 1) as
40 a very broad indication of sites that had a low risk of cold damage (*cf.* Murray et al. 1986).

41
42
43
44
45
46
47
48
49 An indication of relative cold of a site is currently not incorporated into Forest Research's
50 Ecological Site Classification (ESC) (Pyatt et al 2001) decision support system, a web-based
51 programme for matching species to site.

1 An important means of reducing risk is ensuring suitable species and provenances of
2 *Eucalyptus* are planted. Results from trials established in the 1980s (Evans 1986) or informal
3 planting (Purse and Leslie 2016a) have yielded useful information on species and
4 provenances that are well-adapted to parts of the UK. It is clear a wider range of eucalypts
5 could be planted (Purse and Leslie 2016a) than *E. gunnii* and *E. nitens* as proposed in
6 Hardcastle (2006) and incorporated into ESC. This has been recognised and the species most
7 planted in the UK between 2011 and 2015 was *Eucalyptus glaucescens* (Purse and Leslie
8 2016b). Table 4 describes recommended provenances for five promising eucalypts and the
9 areas of the UK where they are best suited. However, obtaining seed of superior provenances
10 has proven difficult and it is likely that sub-optimal origins have been planted in recent
11 decades in the UK.

12 The risk of damage by cold can be reduced further by focusing on species that readily
13 coppice as they are likely to recover from the main stem being killed by cold and thus
14 replanting costs can be avoided. Furthermore, coppice rotations are more productive than
15 first-rotations (Pukkala and Pohjonen 1990). A further approach to reduce damage is the use
16 of intensive silviculture to accelerate establishment and growth as larger trees are more
17 resistant to cold than smaller ones (Leslie et al. 2014a). Finally, to reduce the risk of
18 complete failure, plantations should comprise stands reflecting the full range of ages, from
19 those recently established to those of rotation age, as younger stands are more susceptible to
20 damage from cold.

21 A further positive influence on the suitability of eucalypts to the climate in the UK and other
22 temperate maritime environments is the likely effect of climate change on the frequency and
23 duration of extreme cold events. Climate models predict a reduction in the frequency of cold
24 events during winters in Western Europe, and in general a reduced severity (Peings et al.

2013). Gloning et al. (2013) predict warmer winter temperatures and no increase in the frequency of extreme winter temperature events for the UK. However, warmer springs may lead to greater frost damage; a study by Augspurger (2013) at a woodland in Illinois, USA described increased risks of frost damage to woody species over the last 100 or more years associated with an increased frequency of periods of warming in March followed by frosts in April. The species other than eucalypts listed by Hardcastle (2006) are pioneer species and so their phenology is influenced more by photoperiod and less by temperature (Basler and Körner 2012). It is likely that climate change will extend their growing season, but this will also increase the risk of frost damage (Basler and Körner 2012). A potential negative factor for eucalypts are increased levels of carbon dioxide. Heightened levels (twice ambient concentrations) of atmospheric carbon dioxide have been shown to increase susceptibility to frost damage through an increased ice nucleation temperature in cold-tolerant eucalypts, probably caused by bacteria on the leaves (Lutze et al. 1998).

The main constraints for using the wood from eucalypts as a source of biomass is the high moisture content and chemical composition of the wood. The high moisture content and requirement for some species for debarking for effective drying require additional inputs to produce a quality wood fuel. However, most of the plantations of woody energy crops in the UK have been SRC and wood from this source has as high a moisture content as eucalypt SRF. The high chlorine content of eucalypt wood creates acid gases and corrosion of boilers, however the concentration of chlorine can be considerably reduced (as much as 90%) by pre-heating the wood before burning through torrefaction (Kiepi et al. 2014).

If eucalypts were to be planted on a large scale in the UK, the environmental impacts would need to be quantified. The risk of invasiveness of *E. gunnii* and *E. nitens* is low. For both species, seed germination can be poor and the seedlings are susceptible to frost damage and,

1 for *E. gunnii*, it is also palatable to deer. Furthermore, for both species the small seed size
2 means that seedlings have few reserves and are vulnerable to competition from other plants;
3
4 there are few UK sites where natural regeneration of *E. gunnii* has been observed and none
5
6 where *E. nitens* has been recorded (GB Non Native Species Secretariat a & b). In general,
7
8 Booth (2013) highlights the low risk of invasiveness of eucalypts in frost prone areas of the
9
10 world. *Nothofagus* is also not considered to be a threat and is included in the exotic tree
11
12 species where planting is not regulated under the Wildlife and Natural Environment Act of
13
14 2011 (Forestry Commission Scotland 2015).
15
16
17
18
19

20 There are limited studies on the effects of eucalypts on flora and fauna in the UK. Surveys of
21
22 fungi showed that most of the mycorrhizal fungi associated with eucalyptus had originated
23
24 from Australia, with a limited number of native British species (Pennington et al. 2011),
25
26 although another survey identified rare fungal species, including three species representing
27
28 three new genera to the UK (Hobart 2012). A study of earthworms under SRF and in
29
30 comparison with pasture (Rajapaksha et al. 2013) suggested SRF should focus on native
31
32 species but also *Eucalyptus* spp., which also supported dense populations of earthworms.
33
34
35
36

37 Under the Great Britain Non-native Species Risk assessments (GB Non Native Species
38
39 Secretariat no date a, & b), reviews were conducted of the environmental risk associated with
40
41 *E. gunnii* and *E. nitens*. The conclusion of the analysis for both species was that they were
42
43 both in the upper level of the 'low' risk category for environmental impact.
44
45
46

47 In a review of predicted impacts of SRF on water quality and supply no problems were
48
49 associated specifically with eucalypts, rather with intensively grown plantations (Nisbet et al.
50
51 2011). Furthermore, evidence shows that eucalypts are very efficient at using water to
52
53 produce biomass (Dvorak 2012).
54
55
56
57
58
59
60
61
62
63
64
65

Conclusion

Eucalypts have many properties that make them attractive as a source of woody biomass in the UK. Where eucalypts have the greatest advantages is their potential growth rates that exceed those of the other species and are achieved over short rotations, although in practice high yields are often not achieved due to poor survival due to low temperatures. The risk of poor survival can however be mitigated by adopting four management approaches. The first is to focus on planting eucalypts that readily coppice as they are likely to recover from the main stem being killed by cold and thus replanting costs can be avoided. Furthermore, coppice rotations are more productive than first-rotations (Pukkala and Pohjonen 1990). A second means of reducing risk is the use of origins that have been proven to be best adapted to UK conditions. There is a growing body of information on the most suitable provenances for a limited range of eucalypts (Table 4). A third approach to reduce damage is the use of intensive silviculture to accelerate establishment and growth as larger trees are less damaged by cold than smaller ones (Leslie et al. 2014a). Finally, plantations should comprise stands reflecting the full range of ages, from those recently established to those of rotation age, as younger stands are more susceptible to damage from cold.

The high moisture content of green wood and the high chlorine content released during their combustion (Keipi et al. 2014) are constraints to the use of eucalypt wood as a fuel.

However, the moisture content is not higher than some other species, such as poplars (Table 3) and for SRC willow and methods have been devised for drying this material (Whittaker et al 2018). The problem of chlorine release on combustion can also be addressed through pretreatment by torrefaction (Keipi et al. 2014).

References

AFOCEL (2004). Information *Eucalyptus*, *Eucalyptus* et environnement. Lettre

d'information semestrielle eucalyptus. Numero 3, Vol. 2 Juillet 2004, AFOCEL, France.

AFOCEL (2007) *Eucalyptus*: 35 ans d'experimentation dans le sud de la France.

Informations Foret, No 2-2007, Fiche No 747. AFOCEL, France.

Augspurger CK (2013) Reconstructing patterns of temperature, phenology, and frost damage over 124 years: Spring damage risk is increasing. *Ecol* 94:41–50.

Basler, D. and Körner, C., (2012). Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agric and For Meteorolog* 16:73-81.

Beadle CL, Honeysett JL, Turnbull CRA and White DA (1995) Site limits to achieving genetic potential. In *Eucalypt Plantations: Improving Fibre Yield and Quality*. Proceedings CRC Temperate Hardwood Forestry-IUFRO Conference:19-24.

Black K (no date) Freeze tolerance of Eucalypt seedlings: a tool for assessing frost hardiness. University College Dublin, Ireland. 6pp.

Booth TH (2013) Eucalypts and Their Potential for Invasiveness Particularly in Frost-Prone Regions. *International Journal of Forest Research*. 1-7.

<http://www.hindawi.com/journals/ijfr/2012/837165/citations/>. Accessed on 18 June 2015

Brown A and Webber J (2008) Red band needle blight of conifers in Britain. *Forestry Commission Research Note* 2.

[http://www.forestry.gov.uk/pdf/FCRN002.pdf/\\$FILE/FCRN002.pdf](http://www.forestry.gov.uk/pdf/FCRN002.pdf/$FILE/FCRN002.pdf). Accessed 20 July 2017

Cameron AD (1996) Managing birch woodlands for the production of quality timber. *Forestry* 69:357-371.

Černý K, Filipová N and Strnadová V1 (2012) Influence of low temperature and frost duration on *Phytophthora alni* subsp. *alni* viability. *For Syst* 21:337-342.

Chauzat M-P, Purves G and Dunn R (2002) Release and establishment of a biological control agent, *Psyllaephagus pilosus* for eucalyptus psyllid (*Ctenarytaina eucalypti*) in Ireland. *Ann Appl Biol* 141:293-304.

Christersson L (2010) Wood production potential in poplar plantations in Sweden. Biomass Bioenerg 34 (9):1289–1299.

Claessens H, Oosterbaan A, Savill P and Rondeux J (2010). A review of the characteristics of black alder (*Alnus glutinosa* (L.) Gaertn.) and their implications for silvicultural practices. Forestry 83:163-175.

Cope MH, Leslie A and Weatherall A (2008) The potential suitability of provenances of *Eucalyptus gunnii* for short rotation forestry in the UK. Q J For 102:185-194.

Davidson NJ and Reid JB (1987). The influence of hardening and waterlogging on the frost resistance of subalpine eucalypts. Aust J Bot 35:91-101

Davidson NJ, Battaglia M and Beadle CL (1995) Photosynthesis of *Eucalyptus nitens* is reduced by mild frosts. Eucalypt Plantations: Improving Fibre Yield and Quality: 339-343.

Deans JD, Billington HL and Harvey FL (1992) Winter Frost Hardiness of two Chilean Provenances of *Nothofagus procera* in Scotland. Forestry 62: 205-212.

DECC (2012) UK Bioenergy Strategy. Department for Transport, DECC, DEFRA. <https://www.gov.uk/government/publications/uk-bioenergy-strategy>. Accessed 23 April 2016.

DECC (2015) Digest of United Kingdom Energy Statistics,.DECC https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/450302/DUKES_2015.pdf. Accessed 23 April 2016.

DEFRA (2007) UK Biomass Strategy, DEFRA. http://www.globalbioenergy.org/uploads/media/0705_Defra_-_UK_Biomass_Strategy_01.pdf. Accessed on 22 September 2017.

Dickmann, D.I. 2006. Silviculture and biology of short-rotation woody crops in temperate regions: then and now. Biomass Bioenerg 30:696–705.

Dobrowolska D, Hein S, Oosterbaan A, Wagner S, Clark J, Skovsgaard JP (2011) A review of European ash (*Fraxinus excelsior* L.): implications for silviculture. Forestry 84: 133-148.

1 Dvorak WS (2012) Water use in plantations of eucalypts and pines: a discussion paper from a
2 tree breeding perspective. *Int For Rev* 14:110-119.

3
4 Edwards PN and Christie JM (1981) *Yield Models for Forest Management*. Forestry
5 Commission Bulletin 48. HMSO, London.

6
7
8
9 European and Mediterranean Plant Protection Organisation (2011) Pest Risk Analysis for
10 *Agrilus anxius*. [https://secure.fera.defra.gov.uk/phiw/riskRegister/plant-](https://secure.fera.defra.gov.uk/phiw/riskRegister/plant-health/documents/agrilusAnxious.pdf)
11 [health/documents/agrilusAnxious.pdf](https://secure.fera.defra.gov.uk/phiw/riskRegister/plant-health/documents/agrilusAnxious.pdf). Accessed 7 August 2015.

12
13
14
15 Evans J (1980) Prospects for Eucalypts as forest trees in Great Britain. *Forestry* 53:129-143.

16
17
18 Evans, J (1986) A Re-assessment of cold-hardy eucalypts in Great Britain. *Forestry* 59:223-
19 242.

20
21
22
23 Fanning PD and Barrs J-R (2013) Biology of the *Eucalyptus* leaf beetle *Paropsisterna*
24 *selmani* (Chrysomelidae: Paropsini): a new pest of *Eucalyptus* species (Myrtaceae) in Ireland.
25 *Agr For Entomol* 16:45–53.

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3 Forestry Commission (2014) *Phytophthora ramorum* - Frequently Asked Questions. at URL:
4 <http://www.forestry.gov.uk/forestry/infd-5ubesn>. Accessed 20 July 2017
5
6

7 Forestry Commission Scotland (2015) Managing invasive and non-native forestry species.
8 Guidance for forest managers and owners.
9
10 [http://scotland.forestry.gov.uk/images/corporate/pdf/managing-invasive-and-non-native-](http://scotland.forestry.gov.uk/images/corporate/pdf/managing-invasive-and-non-native-forestry-species.pdf)
11 [forestry-species.pdf](http://scotland.forestry.gov.uk/images/corporate/pdf/managing-invasive-and-non-native-forestry-species.pdf). Accessed 2 November 2017.
12
13
14

15
16 GB Non Native Species Secretariat (no date a) Non-native Species Risk assessments
17 *Eucalyptus gunnii* – Cider gum. www.nonnativespecies.org/downloadDocument.cfm?id=756.
18 Accessed 6 August 2015
19
20
21

22 GB Non Native Species Secretariat (no date b) Non-native Species Risk assessments
23 *Eucalyptus nitens* – Shining gum.
24 www.nonnativespecies.org/downloadDocument.cfm?id=756A. Accessed 6 August 2015
25
26
27

28
29 Gibbs JN, Lipscombe MA and Peace AJ (1999) The impact of *Phytophthora* disease on
30 riparian populations of common alder (*Alnus glutinosa*) in southern Britain. Eur J For Pathol
31 29:39-50.
32
33
34

35
36 Gloning P, Estrella N and Menzel A (2013) The impacts of climate change on the winter
37 hardiness zones of woody plants in Europe. Theor Appl Climatol 113: 683–695.
38
39

40 Gordon DR, Tancig KJ, Onderdonk DA, Gantz CA (2011). Assessing the invasive potential
41 of biofuel species proposed for Florida and the United States using the Australian Weed Risk
42 Assessment. Biomass Bioenerg 35: 74-79.
43
44
45

46 Green S (2005) Birch dieback in Scotland. Forestry Commission Information Note 72.
47 [http://www.forestry.gov.uk/PDF/fcin072.pdf/\\$FILE/fcin072.pdf](http://www.forestry.gov.uk/PDF/fcin072.pdf/$FILE/fcin072.pdf). Accessed 22 August 2015.
48
49
50

51 Guidi W, Pitre FE, Labrecque M (2013). Short-rotation coppice of willows for the production
52 of biomass in Eastern Canada, In: Matovic, M.D. (Ed.), Biomass– Now - Sustainable Growth
53 and Use. InTech, pp. 421–448. [http://intechopen.com/books/biomass-now-sustainable-growth-](http://intechopen.com/books/biomass-now-sustainable-growth-and-use)
54 [and-use](http://intechopen.com/books/biomass-now-sustainable-growth-and-use). Accessed 18 June 2016.
55
56
57
58
59
60
61
62
63
64
65

Hardcastle PD (2006). A review of the impacts of short-rotation forestry. LTS International, Edinburgh.

[http://www.forestry.gov.uk/pdf/SRFFinalreport27Feb.pdf/\\$FILE/SRFFinalreport27Feb.pdf](http://www.forestry.gov.uk/pdf/SRFFinalreport27Feb.pdf/$FILE/SRFFinalreport27Feb.pdf). Accessed 27 June 2007.

Harrison A (2011) Growing Energy Forestry: trials update, Research update presentation, 17 March 2009.

[https://www.forestry.gov.uk/pdf/Energy_Forestry_020611_Harrison_A.pdf/\\$FILE/Energy_Forestry_020611_Harrison_A.pdf](https://www.forestry.gov.uk/pdf/Energy_Forestry_020611_Harrison_A.pdf/$FILE/Energy_Forestry_020611_Harrison_A.pdf). Accessed 9 August 2016.

Hastings A, Tallis MJ, Casella E, Matthews RW, Henshall PA, Milner S, Smith P and Taylor G (2014) The technical potential of Great Britain to produce ligno-cellulosic biomass for bioenergy in current and future climates. *Glob Change Biol Bioenerg* 6:108–122.

Hedberg E, Kristensson A, Ohlsson M, Johansson C, Johansson P, Swietlicki E, Veselya V, Wideqvista U, Westerholme R (2002) Chemical and physical characterization of emissions from birch wood combustion in a wood stove. *Atmos Environ* 36:4823–4837

Hein S, Collet C, Ammer C, Le Goff N, Skovsgaard and Savill P (2009) A review of growth and stand dynamics of *Acer pseudoplatanus* L. in Europe: implications for silviculture. *Forestry* 82:361-385

Hinchee M, Rottmann W, Mullinax L, Zhang C, Chang S, Cunningham M, Pearson L, and Nehra N (2009) Short-rotation woody crops for bioenergy and biofuels applications. *In Vitro Cell Dev Biol Plant* 45:619–629.

Hobart C (2012) *Eucalyptus* a host for fungal aliens new to the UK. *Field Mycol* 13:51-56.

Horgan FG (2012) Effects of leaf damage on oviposition choice in an invasive paropsine beetle. *J Appl Entomolog* 136:271–281.

Hutchinson K, Thompson D and Berkery B (2011) The Potential of *Eucalyptus* Species as a Source of Biomass / Fibre in Ireland. Powerpoint presentation.

Huhtinen M (2006) Wood biomass as a fuel. https://www.hnee.de/obj/CD8F42A0-B396-40E7.../5Eures_WoodPropertiesPDF.pdf. Accessed 3 January 2019

- Hynynen J, Niemisto P, Vihera-Aarnio A, Brunner A, Hein S and Velling P (2009) Silviculture of birch (*Betula pendula* Roth and *Betula pubescens* Ehrh.) in northern Europe. Forestry 83:103-119.
- Keipi T, Tolvanen H, Kokko L and Raiko R (2014) The effect of torrefaction on the chlorine content and heating value of eight woody biomass samples. Biomass Bioenerg 66: 232-239.
- Kent T, Kofman P, Owens E, Coates E and Cooley S (2009) FORESTENERGY, Harvesting and processing forest biomass for energy production in Ireland. Coford, Dublin, Ireland.
- Kerr G (2011). A Review of the growth, yield and biomass distribution of species planted in the English network trials of Short Rotation Forestry, in McKay, H. (ed.) (2011) Short Rotation Forestry: review of growth and environmental impacts. Forest Research Monograph, 2, Forest Research, Surrey, pp135-164.
- Kerr G and Evans J (2011). Eucalypts for Short Rotation Forestry: a case study from the 1980s. Q J Forest 105:109-117.
- Kibblethwaite RP, Johnson BI and Shelbourne JA (2001) Kraft pulp qualities of *Eucalyptus nitens*, *E. globulus* and *E. maidenii* at ages 8 and 11 years. N Zeal J For Sci 30:447-457.
- Leslie, A.D (2013) Eucalypts in the Republic of Ireland. Q J Forest 107:62-64.
- Leslie AD, Mencuccini M and Perks MP (2011) Eucalyptus in the British Isles. Q J Forest 205:43-53.
- Leslie AD, Mencuccini M and Perks MP (2012) The potential for *Eucalyptus* as a wood fuel in the UK. Appl Energ 89:176-182.

1 Leslie AD, Mencuccini M and Perks MP (2014a) Frost damage to eucalypts in a short-
2 rotation forestry trial in Cumbria (England), iForest Volume 7.

3
4 Leslie A, Mencuccini M, Purse JG and Perks MP (2014b) Results of a species trial of cold
5 tolerant eucalypts in south west England. Q J Forest, 108:18-27.

6
7
8
9 Leslie AD, Mencuccini M and Perks MP (2018). Preliminary growth functions for
10 *Eucalyptus gunnii* in the UK. Biomass Bioenerg 108:464-469.

11
12
13
14 Logan JA, Régnière J, and. Powell JA (2003). Assessing the impacts of global warming on
15 forest pest dynamics. For Ecol Environ 1:130-137

16
17
18
19 Lutze JL, Roden JS, Holly CJ, Wolfe J, Egerton JJG and Ball MC (1998). Elevated
20 atmospheric [CO₂] promotes frost damage in evergreen seedlings. Plant Cell Environ 21:
21 631-635.

22
23
24
25 Marriage A. (1977) Fast-growing firewood. Q J For 71:199-203.

26
27
28 Mason B, Jinks R, Savill, P and Wilson SM (2018) Southern Beeches (*Nothofagus* species).
29 Q J For 112: 30-43.

30
31
32
33 Matthews RW and Broadmeadow MSJ (2009) The potential of UK forestry to contribute to
34 Government's emissions reduction commitments, Chapter 8 in Read DJ, Freer-Smith PH,
35 Morison JIL, Hanley N, West CC and Snowdon P (eds) (2009). Combating climate change -
36 A role for UK forests. An assessment of the potential of the UK's trees and woodlands to
37 mitigate and adapt to climate change. The Stationery Office, Edinburgh: 139-162.

38
39
40
41
42
43 McEvoy C (2016) Short Rotation Forestry Trials in Scotland, Progress report 2015.
44 [http://scotland.forestry.gov.uk/images/corporate/pdf/energy-forestry-trials-progress-report-](http://scotland.forestry.gov.uk/images/corporate/pdf/energy-forestry-trials-progress-report-2016.pdf)
45 [2016.pdf](http://scotland.forestry.gov.uk/images/corporate/pdf/energy-forestry-trials-progress-report-2016.pdf). Accessed 13 June 2017.

46
47
48
49 McKay H (2010) Is planting Eucalypts a good idea? Powerpoint presentation at the National
50 School of Forestry, Newton Rigg, 21 January 2010.

51
52
53
54 McKay H (ed.) (2011) Short Rotation Forestry: review of growth and environmental impacts.
55 *Forest Research Monograph*, 2, Forest Research, Surrey, 212pp.

56
57
58
59
60
61
62
63
64
65

McKendry P (2002). Energy production from biomass (part 1): overview of biomass. *Bioresour Technol* 83:37-46.

Met Office (no date) Minimum temperature monthly lowest (deg C) January 1961-90. The Met Office, London.

Met Office (2011) Winter 2010/ 2011. <http://www.metoffice.gov.uk/climate/uk/2011/winter.html>. Accessed 3 September 2012

Milch J, Vavrcik H, Tippner J and Brabec M (2015) The effect of growth conditions in specific areas of Croatia and the Czech Republic on the physical and mechanical properties of black alder wood (*Alnus glutinosa* Gaertn.) *Turk J Agric For* 30:1-6.

Mitchell CP, Ford-Robertson JB and Waters MP (1993). Establishment and monitoring of large scale trials of short rotation coppice for energy. Wood Supply research group, Department of Forestry, University of Aberdeen: 76pp plus Appendices.

Murray MB, Cannell MGR and Sheppard LJ (1986). Frost Hardiness of *Nothofagus procera* and *Nothofagus obliqua* in Britain. *Forestry* 59:209-222.

Neilan J and Thompson D (2008). *Eucalyptus* as a potential biomass species for Ireland. COFORD Connects, Reproductive Material No.15. COFORD, Ireland: 8pp

Nielsen DG, Mullenberg VL and Herms DA (2011) Interspecific variation in resistance of Asian, European and North American birches (*Betula* spp) to bronze birch borer (Coleoptera: Buprestidae). *Environ Entomolog* 40: 648-653.

Nisbet T, Thomas H and Shah N (2011) Short Rotation Forestry and Water in McKay, H. (ed.) (2011) Short Rotation Forestry: review of growth and environmental impacts. Forest Research Monograph 2, Forest Research, Surrey. 13-36.

OFGEM (2019a) Non-domestic Renewable Heat Incentive. <https://www.ofgem.gov.uk/environmental-programmes/non-domestic-rhi>. Accessed 1 January 2019.

OFGEM (2019b) About the FIT. <https://www.ofgem.gov.uk/environmental-programmes/fit/about-fit-scheme>. Accessed 1 January 2019.

- Okland B, Haack RA and Wilhelmsen G (2012) Detection probability of forest pests in current inspection protocols – a case study of the bronze birch borer. *Scand J For Res* 27: 285-297.
- O'Reilly C, Tobin B and Farrelly N (2014) Can short rotation forestry play a role in renewable energy demands? *For Eng Rev* 4: 32-34.
- Paton DM (1983) Physiology of frost resistance in *Eucalyptus*. In: “Colloque International sur les eucalypts au froid”. Nangis (France) 26-30 September 1983. AFOCEL, Nangis, France:107-125.
- Peings Y, Cattiaux J and Douville H (2013) Evaluation and response of winter cold spells over Western Europe in CMIP5 models. *Clim Dynam* 41:3025–3037.
- Pennington HG, Bidartondo MI and Barsoum N (2011) A few exotic mycorrhizal fungi dominate eucalypts planted in England. *Fungal Ecol* 4:299-302
- Prior J and Kendon M (2011). The UK winter of 2009/2010 compared with severe winters of the last 100 years. *Weather* 66:4-10.
- Pukkala T and Pohjonen V (1990) Yield models for *Eucalyptus globulus* fuelwood plantations in Ethiopia. *Biomass* 21 (2): 129-143.
- Purse J (2010) *Eucalyptus gunnii*–the pioneer Australian tree in Britain. *Aust Plant* 25(205): 376-382.
- Purse JG and Richardson KF (2001) Short rotation single stem tree crops for energy in the UK- an examination with Eucalyptus. *Aspects of Applied Biology*, No.65, 2001, Biomass Eng Crop II:.1-8.
- Purse JG and Leslie AD (2016a) Eucalyptus - Part 1 Species with forestry potential in the British Isles. *Q J For* 110:88-97.
- Purse, J and Leslie, A.D. (2016b) Eucalyptus – Part 2. Findings from trial plantings, and silvicultural requirements in the British Isles. *Q J For* 110:161-168.
- Pyatt DG, Ray D and Fletcher J (2001) An ecological site classification for forestry in Great Britain. Forestry Commission Bulletin 124 Forestry Commission, Edinburgh.

1 Rajapaksha NSS, Butt KR, Vanguelova EI and Moffat AJ (2013) Effects of Short Rotation
2 Forestry on earthworm community development in the UK. For Ecol Manag 309:96-104.
3

4 Ranney JW and Mann LK (2004) Environmental considerations in energy crop production.
5 Biomass Bioenerg 6:211-228.
6

7 Ramsay, P. (2004) Viable alternatives. For Mach J 9: 18-19.
8

9 Ray D and Sing L (2006) Annex 4: Climate Change and Species Suitability Analysis using
10 Ecological Site Classification (ESC) in Hardcastle (2006) A review of the impacts of short-
11 rotation forestry. LTS International.
12
13 [http://www.forestry.gov.uk/pdf/SRFFinalreport27Feb.pdf/\\$FILE/SRFFinalreport27Feb.pdf](http://www.forestry.gov.uk/pdf/SRFFinalreport27Feb.pdf/$FILE/SRFFinalreport27Feb.pdf).
14 Accessed 27 June 2007:
15
16

17 Read DJ, Freer-Smith PH, Morison JIL, Hanley N, West CC and Snowdon P (2009) (Eds)
18 Combating climate change: a role for UK forests. An assessment of the potential of the UK's
19 trees and woodlands to mitigate and adapt to climate change. The Stationery Office Limited,
20 London.
21

22 SAC (2009). An assessment of the potential impact on UK agriculture and the environment
23 of meeting renewable feedstock demands. SAC Consulting, Edinburgh.
24
25

26 Savill P (2013) The silviculture of trees used in British forestry. 2nd Edition, CABI, Oxford.
27

28 Scanu B, Jones B, Webber JF (2012). A new disease of *Nothofagus* in Britain caused by
29 *Phytophthora pseudosyringae*. New Dis Rep 25: 27.
30

31 Senelwa K and Sims REH (1999) Fuel characteristics of short rotation forest biomass.
32 Biomass Bioenerg 17: 127-140.
33

34 Sims REH, Senelwa K, Maiava T, Bullock BT (1999) Eucalyptus species for biomass energy
35 in New Zealand, Part II: Coppice performance. Biomass Bioenerg 17: 333-343.
36

37 Solid Fuel Association (no date) Solid Fuel association Guide to wood and multifuel.
38 http://www.solidfuel.co.uk/pdfs/guide_to_wood_and_multifuel.pdf. Accessed 10 August
39 2015.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Stokes V, Cottrell J and A'Hara S (2017) Summary report on Forest Research aspen clonal trials. Forest Research, Roslin.

Straw NA, Williams DT, Kulinich O, Gninenko YI (2013) Distribution, impact and rate of spread of emerald ash borer *Agrilus planipennis* (Coleoptera: Buprestidae) in the Moscow region of Russia. Forestry 86:515–522.

Sturrock RN, Frankel SJ, Brown AV, Hennon AV, Kliejunas JT, Lewis KJ, Worrall JJ and Woods AJ (2011) Climate change and forest diseases, review. Plant Pathol 60: 133–149

Tabbush P and Lonsdale D (1999) Approved poplar varieties. Forestry Commission Information Note 21. Forestry Commission, Edinburgh.

Teagasc (2015) Preliminary results on the physical & chemical properties of *Eucalyptus* in Ireland.
http://www.teagasc.ie/forestry/docs/research/shortfor_euc_handout_DPlant_210515.pdf.
 Accessed 20 May 2016.

Tharakan PJ, Volk TA, Abrahamson LP and White EH (2003) Energy feedstock characteristics of willow and hybrid poplar clones at harvest age. Biomass Bioenerg 25:571-580.

Thomas PA (2016) Biological Flora of the British Isles: *Fraxinus excelsior*. J Ecol 104: 1158-1209.

Trebrown (no date) Plant hardiness zones of the British Isles.
<http://www.trebrown.com/hrdzone.html> Accessed 4 January 2019.

Tobin B, Foreman S and O'Reilly C (2016) Short Rotation Forestry in Ireland - new research trials. For Eng Rev 6: 28–30.

Tuley G (1980) *Nothofagus* in Britain. Forestry Commission Forest Record 122. HMSO, London.

UK Plant Health Risk Register (no date) UK Plant Health Risk Register.L:
<https://secure.fera.defra.gov.uk/phiw/riskRegister/>. Accessed 24 June 2016

USDA Forest Service (no date) Wood technical fact sheet – *Nothofagus*.

http://www.fpl.fs.fed.us/documnts/TechSheets/Chudnoff/SEAsian_Oceanic/htmlDocs_SEAsian/Nothofagusspp.html. Accessed 30 July 2015

Wainhouse D, Inward DJG, Denman S, Green Sand Webber JF (2016), Agriculture and Forestry Climate change report card technical paper 7. Insect Pests and Pathogens of Trees.

<http://www.nerc.ac.uk/research/partnerships/ride/lwec/report-cards/agriculture-source07/>

Accessed 18 December 2017.

Walker JCF, Butterfield BG, Harris JM, Langrish TAG and Prichard JM (2013) Primary Wood Processing: Principles and Practice. Springer.

Webber J, Tilbury C, Steele H and Rose D (2011) Potential Impacts of Pests and Pathogens on Short Rotation Forestry in Britain in McKay, H. (ed.) (2011) Short Rotation Forestry: review of growth and environmental impacts. Forest Research Monograph 2. Forest Research, Surrey. pp165-190.

Web of Science (2018) Web of Science for UK Education. <http://wok.mimas.ac.uk/>.

Accessed 7 January 2019.

Whittaker C, Yates NE, Powers SJ, Misselbrook T and Shield I (2018) Dry matter losses and quality changes during short rotation coppice willow storage in chip or rod form. Biomass Bioenerg 112:29-36.

Woodward S and Boa E (2013) Ash dieback in the UK: a wake-up call. Mol Plant Pathol 14:856-860.



Figure 1 Locations of sites with shaded area representing the -14°C minimum temperature monthly lowest (January) isotherm for (1961-90) (Met Office no date). The locations are as follows: (1) Moray, (2) Kilmichael, (3) Glenbranter, (4) Wark, (5) Daneshill, (6) Thetford, (7) Redmarley, (8) Exeter, (9) New Forest, (10) Chiddingfold.

Table 1 Characteristics of SRC and SRF (modified from SAC 2009)

	SRC	SRF
Production practices	Established at high planting densities using willow cuttings which are harvested every 2-4 years	Established from transplants at lower planting densities and harvested every 8-12 years
Inputs	Pre-planting herbicide. N application in year 2 after cutting. Few additional inputs	Pre-planting herbicide. N application to reflect crop uptake and maintain crop vigour. Few additional inputs
Yields	7-12 Mg (oven dried) ha ⁻¹ y ⁻¹	5-15 Mg (oven dried) ha ⁻¹ y ⁻¹ estimated – depending on species

Table 2 Attributes of the ideal SRF tree crop.

Silvicultural/ agronomic attributes	Wood property characteristics	Environmental characteristics
<ul style="list-style-type: none"> • Fast growth and high biomass yield (Guidi et al 2013), with mean annual increment (MAI) peaking early. • Resistant to pests and diseases and extremes in climate, such as cold and drought. • The ability to coppice (Dickman 2006, Hinchee et al 2009, Guidi et al 2013), or to sucker which avoids the costs of replanting and also enhances growth rates in the second and subsequent rotations (Dickman 2006, Hinchee et al 2009, Guidi et al 2013); • Produces straight stems; lowering harvesting, handling and transportation costs (Walker et al 2013); 	<ul style="list-style-type: none"> • Higher density wood in SRF species resulting in higher energy outputs per unit volume (Ramsay 2004) • Wood with a low moisture content and other characteristics that reduce the drying requirement (Senelwa and Simms 1999) • Wood with suitable chemical characteristics for combustion (Senelwa and Simms 1999), • Wood attractive to markets other than biomass, reducing market risk. 	<ul style="list-style-type: none"> • Low negative impacts on the environment, such as soil nutrients and moisture (Ranney and Mann 1994). • Reproductive or other characteristics that limit the likelihood of invasiveness (Gordon et al 2011) or hybridisation with native populations of trees (Felton et al 2013)

Table 3 Estimates of mean annual increment and biomass yield for potential SRF species (cf.Hardcastle 2006).

Species	Mean MMAI (m ³ ha ⁻¹ y ⁻¹) and range	Biomass yield Mg ha ⁻¹ y ⁻¹ (oven dried)	Specific gravity	Green Moisture content
<i>Eucalyptus nitens</i>	26 to 30 ⁷	N/A	0.45 ⁹ , 0.435 to 0.446 ²¹	56.2-59.3% ²¹
<i>Eucalypts gunnii</i>	16 ⁶	1.5 to 8.2 ⁵	0.50 ¹¹	-
<i>Nothofagus</i>	14 (10 to 18) ³	3.0 to 10.5 ⁵	0.6 ¹² , 0.45 to 0.53 ¹³	-
Poplar	9 (4 to 14) ¹	4.2 ⁵	0.36 ⁵ 0.335 ¹⁴ (aspen 0.48 ¹⁵)	64% ¹⁶ , 49- 56% ¹⁹
Sycamore	8 (4 to 12) ¹	0.6 to 5.7 ⁵	0.63 ⁸ (MC 12- 17%)	41% ¹⁶ , 48% ²²
Alder	4.5 to 14.6 ²	0.9 to 4.8 (red alder) ⁵	0.54 ¹⁰ (MC 12%), 0.43-0.49 ¹⁸	53% ²²
Birch	4 to 10 ⁴	0.5 to 5.7 ⁵	0.662 ¹⁷ , 0.53 ²⁰	43% ¹⁶
Ash	6 (2 to 10) ¹	0.5 to 4.7 ⁵	0.674 ¹¹	32% ¹⁷ , 40% ²³

¹Edwards and Christie (1981), ²Claessens et al (2010), ³Tuley (1980), ⁴Hynenen et al (2009),
⁵Kerr (2011) Table 16, ⁶Leslie et al. 2018, ⁷O'Reilly, Tobin and Farrelly (2014), ⁸Hein et al
(2009), ⁹Kibblewhite et al (2000), ¹⁰Claessens et al (2010), ¹¹AFOCEL(2004), ¹²Tuley
(1980), ¹³USDA (no date), ¹⁴Christersson (2010), ¹⁵Harrison (2009), ¹⁶Forestry Commission
(2011), ¹⁷Solid Fuel association no date. ¹⁸Milch et al (2015). ¹⁹Tharakan et al (2003),
²⁰Cameron (1996), ²¹Teagasc (2015), ²²Kent et al (2009).

Table 4 Recommended origins for selected eucalypts. Refer to Figure 1 for locations of sites.

Species	Recommended origin	Source	Notes
<i>E. delegatensis</i>	Var <i>tasmaniensis</i> from Ben Lomond, Tasmania	One of the more cold hardy origins in Evans (1986) and confirmed in Leslie et al (2014b).	Some mainland Australian origins are also hardy (Evans 1986). Recommended for warm areas of south west Britain with fast growth and good survival at a trial at Haldon Forest near Exeter. A valuable timber tree (FAO 1981).
<i>E. glaucescens</i>	Guthega, New South Wales	The most cold hardy of six origins in Evans (1986)	Adaptable to a range of sites and is unpalatable. One of the hardiest of eucalypts (Purse and Leslie (2016a)
<i>E. gunnii</i>	Lake MacKenzie, Tasmania.	One of the most cold hardy origins in Evans (1986) and confirmed Cope et al. (2008).	Performs well over a range of locations but variable growth; at 3 years of age trees at Exeter were twice the height of those at Chiddingfold, Thetford, Glenbranter or Wark. (Leslie et al. 2018)
<i>E. nitens</i>	Higher altitude provenances from Victoria.	Evans (1986)	Rapid growth but only to be planted in the least cold and least exposed sites.
<i>E. subcrenulata</i>	Mount Cattley, Tasmania	Evans (1986) recommended central or southern Tasmanian origins. Mount Cattley origin recommended in Leslie et al. (2014b)	To be planted in warm areas of south west Britain. Excellent survival and growth at a trial at Haldon Forest near Exeter (Leslie et al. 2014b).