

Tubbs, Sarah E., Baş, Aylin Akkaya, Côté, Gabrielle, Jones, Amy L. and Notman, Gill ORCID: <https://orcid.org/0000-0002-7531-2709> (2019) Sighting and stranding reports of Irrawaddy dolphins (*Orcaella brevirostris*) and Dugongs (*Dugong dugon*) in Kep and Kampot, Cambodia. *Aquatic Mammals*, 45 (5). pp. 563-568.

Downloaded from: <https://insight.cumbria.ac.uk/id/eprint/5101/>

***Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.***

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

**provided that**

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
  - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

**You may not**

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing [insight@cumbria.ac.uk](mailto:insight@cumbria.ac.uk).

Short Note

**Sighting and Stranding Reports of Irrawaddy dolphins (*Orcaella brevirostris*) and Dugongs (*Dugong dugon*) in Kep and Kampot, Cambodia**

Sarah E. Tubbs<sup>1</sup>, Aylin Akkaya Baş<sup>2</sup>, Gabrielle Côté<sup>3</sup>, Amy L. Jones<sup>4</sup> and Gillian M. Notman<sup>4</sup>

<sup>1</sup>*Marine Conservation Cambodia, Koh Ach Seh, Kep Archipelago, Kep Province, Cambodia.*

*E-mail: sarahtubbs123@gmail.com*

<sup>2</sup>*Marine Mammals Research Association, Antalya, 07070, Turkey*

<sup>3</sup>*Collégial international Sainte-Anne 1300, boulevard Saint-Joseph Lachine (Québec) H8S 2M8, Canada*

<sup>4</sup>*University of Cumbria, Rydal Road, Ambleside, LA22 9BB, England.*

In Cambodia, information on the distribution of Irrawaddy dolphins *Orcaella brevirostris* and dugongs *Dugong dugon* is limited (Beasley & Davidson, 2007; Hines et al., 2015a). Irrawaddy dolphin presence is confirmed in the coastal waters of Thailand's Trat Province, Cambodia's Koh Kong and Sihanoukville provinces and Vietnam's Kien Giang Province (Figure 1) (Beasley & Davidson, 2007; Ponnampalam, 2013; Smith et al., 2014; Vu et al., 2015; Hines et al., 2015b; Smith et al., 2016). Unconfirmed fisher reports identify the presence of small and sporadic dugong populations at both the Thai-Cambodian and Vietnamese-Cambodian border regions (Perrin et al., 1995, 1996; Nelson, 1999; Beasley & Davison, 2007; Hines et al., 2008). The coastal distribution of both species exposes them to anthropogenic threats, specifically habitat degradation, and fisheries bycatch (Reeves et al. 2003; Hines et al., 2008; Smith et al. 2008; Jaaman et al. 2009; Marsh et al. 2011; Peter et al., 2016a; Peter et al. 2016b; Pilcher et al. 2017). Dugongs are also affected by targeted hunts (Hines et al., 2008; Marsh et al. 2011; Robards & Reeves 2011). Irrawaddy dolphin and dugong populations are declining across Southeast Asia (IWC, 1994; Robards & Reeves 2011), with conservation statuses reflecting these declines.

## Sightings and Strandings in Cambodia

Irrawaddy dolphins were reclassified from ‘vulnerable’ to ‘endangered’ by the IUCN in November 2017 (Minton et al., 2017), with dugongs classified as 'vulnerable' since 2015 (Marsh & Sobotzick, 2015). According to Cambodian Fisheries Law *Sub-decree no. 123* (2009), both species are considered as endangered fisheries resources and are therefore protected by law (*Law on Fisheries*, 2007). In September 2017, The Cambodian Marine Mammal Conservation Project (CMMCP) was established to fill marine mammal knowledge gaps in Cambodian waters and highlight the need for marine mammal conservation strategies in the region. As an initiative of CMMCP, the current study aims to provide initial sighting and occurrence records of Irrawaddy dolphins and dugongs in Cambodia’s Kep and Kampot region. Additionally, we have initiated a photo-identification catalog for the region’s Irrawaddy dolphins, which is part of on-going survey work in the region.

### Methodology

The Kep and Kampot coastline spans from Sihanoukville Province, Cambodia, to Kien Giang Province, Vietnam, on the eastern coast of the Gulf of Thailand (Figure 1). Coastal waters range between depths of 2 and 12m, with the exception of a deeper channel off the western Kampot coast reaching 25m. Kep supports 13 offshore islands, collectively known as the Kep Archipelago (Figure 1), home to fringing coral reef, seagrass meadows, and commercially valuable fish and crustacean species (Cockerell et al., 2016).

(insert Figure 1 here)

Between October 2017 and May 2018, CMMCP conducted 14 boat-based cetacean surveys in Kep’s coastal waters (Table 1). Surveys lasted for a minimum of three hours, during sea state conditions  $\leq$  Beaufort 3. Surveys were conducted on a 20m long by 4.5m wide

## Sightings and Strandings in Cambodia

converted pair trawling boat with a 200HP inboard engine and a viewing platform 3.8m above sea level. The vessel travelled at an average speed of 4 knots around the Kep Archipelago, following a repeated circular route that gave visual coverage of all islands and regions in the Kep Archipelago with known marine mammal occurrence (Figure 2). 100% of the survey route was covered at least once on each survey day, with routes were recorded using a Garmin 64s GPS. Each survey employed four cetacean observers, two scanning the sea surface with Bushnell 8 x 42 binoculars and two with the naked eye. When a group or an individual was sighted, time, species, group size and the number of sub-adults were recorded. Sub-adults were distinguished based on their size. GPS coordinates of dolphin sightings were calculated using boat GPS coordinates, the distance of the group or individual from the boat and the angle of the group or individual from north. Resultant sighting locations were transformed into line data *Esri*<sup>®</sup> *ArcGIS*<sup>™</sup> (ESRI, 2014) and used to create kernel density maps, weighted by the number of animals per group. Mask extraction and percent volume contour, where 50% contours were selected as the core zones, were then used to produce raster maps. 50% contours were selected, as in Bertrand et al. (1996) and Gubbins (2002), as most animals do not use their full home range, but focus their activities in certain areas (Dixon & Chapman, 1980; Samuel et al., 1985; Hauser et al., 2007). Finally, buffers of 1 km radius were created around the core zones. Due to the level of sample effort to date, we did not adjust densities for bias due to survey effort (e.g., McBride et al. 2018). However, this approach will be employed in future surveys.

(insert Figure 2 and Table 1)

Over a total of 14 survey days (40 hours 12 minutes), Irrawaddy dolphins were sighted on 8 days (18 groups). No other cetacean species were encountered during the surveys. Core dolphin habitat was found south of Koh Tonsay, south-east of Koh Tbal and west of Koh Poh

(Figure 1; Figure 3). It also should be noted that project scientists remained in the study area, residing on Koh Ach Seh island (Figure 1), over the entire survey period. Over this time, Illegal, Unreported, and Unregistered (IUU) fishing activities, specifically bottom and electric trawling, were observed and logged in the study region at least once per day.

(insert Figure 3 here)

For all marine mammals sighted between 4 January and 15 May 2018 photographs of dorsal fins and/or tail flukes were captured with a Canon Rebel T6i Digital SLR and EF 75-300mm f/4-5.6 III lens. The photographer avoided bias towards more distinctive individuals by trying to capture images of both sides of each individual in the group. Photograph resolution was used to sort photographs in ‘good’ and ‘poor’ quality folders, with only ‘good’ quality photographs used for post-analysis to identify individuals. Photographs were also assessed for the number of distinctive characteristics shown (e.g., nicks, notches, unusual fin shapes, fin scars, body scars, body deformities). If a ‘poor’ quality image showed two or more distinctive characteristics, it was also included in the analysis. Images of calves and unmarked individuals were discarded to avoid misidentification. *Discovery software* (Gailey & Karczmarski, 2012) was then used to catalogue individuals.

15 individual Irrawaddy dolphins were identified through photo-identification techniques, as reported in CMMCP’s 2018 technical report (Tubbs, 2018). One identified individual possessed deep scars characteristic of entanglement in fishing gear (Figure 4).

(insert Figure 4 here)

On 20 February 2018, a C-POD - Continuous PORpoise Detector - (Chelonia Ltd. 2014a) was deployed in the Kep Archipelago (10° 21' 29"N, 104° 19' 16"E), suspended 0.5m above the

seabed, at a water depth of 2m. C-PODs are fully automated static data loggers, logging peak frequency, duration, intensity and bandwidth for sounds between 20 and 160kHz. *CPOD* software (Chelonia Ltd. 2014b) was used to identify click trains using the KERNO classifier algorithm (Chelonia Ltd. 2014b). Click trains were sorted into ‘high’, ‘moderate’, low’ or ‘doubtful’ quality groups. Based on parameters including frequency and click interval, ‘moderate’ and ‘high’ click trains were further sorted into species classification groups.

The C-POD recorded data for 62 days, with post analysis revealing 60 click trains characterised as belonging to cetaceans. The *CPOD* software’s KERNO classifier (Chelonia Ltd. 2014b) does not allow for species-specific identification to be made; however, as Irrawaddy dolphins were the only species sighted through observational surveys, it is likely that acoustic data belongs to this species.

Between September 2017 and May 2018, CMMCP ran a marine mammal stranding and bycatch network in the Kep and Kampot region. The network is comprised of local fishers and fisheries officers, who report events to project scientists. Over this period, four Irrawaddy dolphin carcasses were stranded in Kep Province and one dugong caught via bycatch in Kampot Province (Table 2), as reported in CMMCP’s 2018 technical report (Tubbs, 2018). No necropsies were conducted, hence causes of death for strandings could not be determined. The figures reported here are likely underestimates of true stranding and bycatch figures, due to lack of rigour of data compilation by fishers or fisheries authorities and/or lack of basic information about species occurrence or identification (Reeves et al., 2005).

(insert Table 2 here)

Baseline knowledge on species presence is the essential first step in the creation of conservation strategies. Here, we report the first observations of Irrawaddy dolphins in Kep and report one incident of dugong bycatch in Kampot. Field observations of IUU fishing were noted during the current study, with previous studies by Nelson (1999) and Beasley and Davidson (2007) also reporting presence of IUU activities in the region. Past and present observations of IUU fishing, combined with marine mammal entanglement, stranding and bycatch cases reported during the current study, identify the urgency for regional conservation strategies. As a starting point, we recommend that the Irrawaddy dolphin core and buffer zone habitat presented in Figure 3 should be managed for the purposes of Irrawaddy dolphin conservation. This study highlights the need for more research activities in the Kep and Kampot region, so data can be used to design tailored conservation strategies for the populations. Conservation strategies for both species will require ongoing, comprehensive data collection on their population, ecology and area use. We, therefore, also recommend expanding survey efforts for Irrawaddy dolphins to the Sihanoukville and Kien Giang Province borders, as well as dedicated dugong studies in Kep province to fill these knowledge gaps.

Author contributions:

Study design and fieldwork: ST, GC

Data analysis: ST, AAB, GC

Writing the article: ST, AAB, AJ, GN

## Acknowledgements

The authors would like to thank Marine Conservation Cambodia for funding the research and Cambodia's Fisheries Administration for their collaboration, specifically Phay Somany, Chheng Touk, and Vibol Ouk. The authors would further like to thank the Marine Mammals Research Association for research advice and support.

## Literature Cited

- Beasley, I. & Davidson, P. (2007). Conservation status of marine mammals in Cambodian waters, including seven new cetacean records of occurrence. *Aquatic Mammals*, 33(3), 368-379.
- Bertrand, M., DeNicola, A., Beissinger, S. & Swihart, R. (1996). Effects of parturition on home ranges and social affiliations of female white-tailed deer. *The Journal of Wildlife Management*, 60(4), 899.
- Chelonia Ltd. (2014a). *C-POD specification*. Available at: <https://chelonia.co.uk/cpods specification.htm> [accessed 21 January 2019].
- Chelonia Ltd. (2014b). *CPOD.exe: a guide for users*. Available at: <https://chelonia.co.uk/downloads/CPOD.pdf> CMS Technical Series No. 32 [accessed 2 February 2018].
- Cockerell, B., Haïssoune, A. & Ferber, P. (2016). *Koh Seh marine environmental assessment*. Unpublished report for the Cambodian Ministry of Agriculture, Forestry and Fisheries.
- Dixon, K. & Chapman, J. (1980). Harmonic mean measure of animal activity areas. *Ecology*, 61(5), 1040-1044.
- ESRI. (2014). *ArcGIS Desktop: Release 10.3*. Redland, CA: Environmental Systems Research Institute.
- Gailey G. & Karczmarski L. (2012). *Discovery: Photo-identification data-management system for individually recognizable animals*. Available at: <https://biosch.hku.hk/ecology/staffhp/1k/Discovery> [accessed 8 May 2018].
- Gubbins, C. (2002). Use of home ranges by resident bottlenose dolphins (*Tursiops truncatus*) in a South Carolina estuary. *Journal of Mammalogy*, 83(1), 178-187.

## Sightings and Strandings in Cambodia

- 172 Hauser, D., Logsdon, M., Holmes, E., Van Blaricom, G. & Osborne, R. (2007). Summer  
173 distribution patterns of southern resident killer whales *Orcinus orca*: core areas and  
174 spatial segregation of social groups. *Marine Ecology Progress Series*, 351, 301-310.
- 175 Hines, E., Adulyanukosol, K., Somany, P., Ath, L., Cox, N., Boonyanate, P. & Hoa, N. (2008).  
176 Conservation needs of the dugong *Dugong dugon* in Cambodia and Phu Quoc Island,  
177 Vietnam. *Oryx*, 42(1).
- 178 Hines, E., Ponnampalam, L., Jamal Hisne, F., Tara S., Jackson-Ricketts, J., Kuit, S. &  
179 Acebes, J. (2015a). *Report of the Third Southeast Asian Marine Mammal Symposium*  
180 *(SEAMAM III)* UNEP / CMS Secretariat, Bonn, Germany.
- 181 Hines, E., Strindberg, S., Junchumpoo, C., Ponnampalam, L., Ilangakoon, A., Jackson-Ricketts,  
182 J. & Monanunsap, S. (2015b). Line transect estimates of Irrawaddy dolphin abundance  
183 along the eastern Gulf Coast of Thailand. *Frontiers in Marine Science*, 2.
- 184 IWC (1994) *Report of the workshop on mortality of cetaceans in passive fishing nets and traps*.  
185 Report of the International Whaling Commission (Special Issue) 1, 1-71.  
186
- 187 Jaaman, S., Lah-Anyi, Y. & Pierce, G. (2009). The magnitude and sustainability of marine  
188 mammal by catch in fisheries in East Malaysia. *Journal of the Marine Biological*  
189 *Association of the United Kingdom*, 89, 907-920.
- 190 *Law on Fisheries* (2007) Cambodia. Available at:  
191 <http://extwprlegs1.fao.org/docs/pdf/cam82001.pdf>. [accessed 18 January 2019].
- 192 Marsh, H. & Sobotzick, S. (2015). *Dugong dugon*. *The IUCN Red List of Threatened Species*  
193 *2015*. Available at: [http://dx.doi.org/10.2305/IUCN.UK.2015-](http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T6909A43792211.en)  
194 [4.RLTS.T6909A43792211.en](http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T6909A43792211.en). [accessed 3 June 2018].
- 195 Marsh, H., O'Shea, T. & Reynolds III, J. (2011). *The ecology and conservation of Sirenia:*  
196 *dugongs and manatees*. Cambridge, England: Cambridge University Press.
- 197 Minton, G., Smith, B., Braulik, G., Kreb, D., Sutaria, D. & Reeves, R. (2017). *Orcaella*  
198 *brevirostris*. In *The IUCN Red List of Threatened Species 2017*. Available at:  
199 <http://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T15419A50367860.en>. [accessed 3  
200 June 2018].
- 201 Nelson, V. (1999). *The coastal zone of Cambodia – Current status and threats: Volumes I and II*.  
202 Unpublished report for the Ministry of Environment and DANIDA.
- 203 Perrin, W., Leatherwood, S., & Dolar, M. (1995). *The status of marine mammal research in*  
204 *Southeast Asia* (UNEP/SEA95/BP6, IBI Reports No. 5), 16.

## Sightings and Strandings in Cambodia

- 205 Perrin, W., Dolar, M., & Alava, M. (1996). *Report of the workshop on the biology and*  
 206 *conservation of small cetaceans and dugongs of Southeast Asia*. Dumaguete: United  
 207 Nations Environment Programme.  
 208
- 209 Peter, C., Poh, A., Ngeian, J., Tuen, A. & Minton, G. (2016a). Identifying habitat characteristics  
 210 and critical areas for Irrawaddy Dolphin, *Orcaella brevirostris*: Implications for  
 211 Conservation. In I. Das & A. Tuen (Eds), *Naturalists, Explorers and Field Scientists in*  
 212 *South-East Asia and Australasia* (pp. 225-238). Springer.
- 213 Peter, C., Ngeian, J., Minton, G., Zulkifli Poh, A., Grinang, J. & Tuen, A. (2016b). *Artisanal*  
 214 *fisheries and cetaceans in Kuching Bay, Sarawak, East Malaysia: Threats and potential*  
 215 *mitigation*. Report presented to the meeting of the Scientific Committee of the  
 216 International Whaling Commission SC/66b/SM09.
- 217 Pilcher, N., Adulyanukosol, K., Das, H., Davis, P., Hines, E., Kwan, D., Marsh, H.,  
 218 Ponnampalam, L. & Reynolds, J. (2017). A low-cost solution for documenting  
 219 distribution and abundance of endangered marine fauna and impacts from fisheries.  
 220 *PLOS ONE*, 12(12).
- 221 Ponnampalam, L., Hines, E., Monanunsap, S., Ilangakoon, A., Junchompoo, C., Adulyanukosol,  
 222 K., & Morse, L. (2013). Short note: behavioral observations of coastal Irrawaddy  
 223 Dolphins (*Orcaella brevirostris*) in Trat Province, Eastern Gulf of Thailand. *Aquatic*  
 224 *Mammals*, 39(4), 401-408.
- 225 Reeves, R., Smith, B., Crespo, E. & Notarbartolo di Sciara, G. (2003). Dolphins, whales and  
 226 porpoises: 2002-2010 Conservation Action Plan for the World's Cetaceans. *IUCN*,  
 227 Gland, Switzerland & Cambridge, UK.
- 228 Reeves, R., Berggren, P., Crespo, E., Gales, N., Northridge, S., di Sciara G., Perrin, W., Read,  
 229 A., Rogan, E., Smith, B., & Van Waerebeek, K. (2005). Global priorities for reduction of  
 230 cetacean bycatch. *WWF*. Available at:  
 231 [https://www.researchgate.net/profile/Koen\\_Van\\_Waerebeek/publication/228345242\\_Global\\_priorities\\_for\\_Reduction\\_of\\_Cetacean\\_Bycatch/links/0fcfd5057ee79c5e7b000000.pdf](https://www.researchgate.net/profile/Koen_Van_Waerebeek/publication/228345242_Global_priorities_for_Reduction_of_Cetacean_Bycatch/links/0fcfd5057ee79c5e7b000000.pdf)  
 232 [bal\\_priorities\\_for\\_Reduction\\_of\\_Cetacean\\_Bycatch/links/0fcfd5057ee79c5e7b000000.p](https://www.researchgate.net/profile/Koen_Van_Waerebeek/publication/228345242_Global_priorities_for_Reduction_of_Cetacean_Bycatch/links/0fcfd5057ee79c5e7b000000.pdf)  
 233 [df](https://www.researchgate.net/profile/Koen_Van_Waerebeek/publication/228345242_Global_priorities_for_Reduction_of_Cetacean_Bycatch/links/0fcfd5057ee79c5e7b000000.pdf) [accessed 5 July 2018]
- 234 Robards, M. & Reeves R. (2011). The global extent and character of marine mammal  
 235 consumption by humans: 1970–2009. *Biological Conservation*, 144, 2770–2786.
- 236 Samuel, M., Pierce, D. & Garton, E. (1985). Identifying areas of concentrated use within the  
 237 home range. *The Journal of Animal Ecology*, 54(3), 711.
- 238 Smith, B., Kong, S., & Saroeun, L. (2014). Conservation status and the use of Irrawaddy  
 239 dolphins as a flagship species for climate adaptation in the Peam Krasop Wildlife  
 240 Sanctuary, Cambodia. *IUCN Asia*, Bangkok, Thailand. Available at:

## Sightings and Strandings in Cambodia

[http://cmsdata.iucn.org/downloads/bcr\\_dolphin\\_survey\\_report\\_final.pdf](http://cmsdata.iucn.org/downloads/bcr_dolphin_survey_report_final.pdf) [accessed 23 January 2019].

Smith, B., Ahmed, B., Mowgli, R. & Strindberg, S. (2008). Species occurrence and distributional ecology of nearshore cetaceans in the Bay of Bengal, Bangladesh, with abundance estimates for Irrawaddy dolphins *Orcaella brevirostris* and finless porpoises *Neophocaena phocaenoides*. *Journal of Cetacean Research and Management*, 10, 45-58.

Smith, B., Sonim, V., Boonchai, R., Joehl Cadena, A. & Manopawitr, P. (2016). *Final Report on Dolphin Conservation along the Coastline of the Thai and Cambodian Border*. Available at: [https://www.iucn.org/sites/dev/files/content/documents/2016/final20report20iucn20cambodia20thailand20dolphin20project\\_final2028229.pdf](https://www.iucn.org/sites/dev/files/content/documents/2016/final20report20iucn20cambodia20thailand20dolphin20project_final2028229.pdf) [accessed 3 July 2018].

*Sub-decree no. 123 Determination of a taxon of fishery and endangered fishery product* (2009). Cambodia. Available at: <https://www.ajne.org/sites/default/files/resource/laws/7217/sub-decree-123-endangered-fisheries-products-2009.pdf> [accessed 18 January 2019].

Tubbs, S. (2018). *The Cambodian Marine Mammal Conservation Project*. Technical report for the Cambodian Ministry of Agriculture, Forestry and Fisheries.

Vu, L., Tho, T., Ny, T., Tham, N., Duy, L., Hung, N. & Duc, H., (2015). *Conservation of cetacean in Kien Giang Biosphere Reserve, Vietnam*. Proceedings at the 21<sup>st</sup> Biennial Conference of the Society for Marine Mammalogy, San Francisco, CA.

## Figures

**Figure 1.** The eastern coast of the Gulf of Thailand and the islands of Cambodia's Kep Archipelago. UPLOADED AS A SEPARATE DOCUMENT.

**Figure 2.** Boat survey tracklines and study area covered between October 2017 and May 2018. UPLOADED AS A SEPARATE DOCUMENT.

**Figure 3.** Boat route core zones, and Irrawaddy dolphin core and buffer zones, produced from tracklines and sightings between October 2017 and May 2018. UPLOADED AS A SEPARATE DOCUMENT.

**Figure 4.** Irrawaddy dolphin with scars characteristic of fishing gear entanglement, Kep Archipelago, 15th May 2018 (Tubbs, 2018). UPLOADED AS A SEPARATE DOCUMENT.

## Tables

**Table 1.** Dates of boat surveys undertaken. UPLOADED AS A SEPARATE DOCUMENT.

## Sightings and Strandings in Cambodia

275 **Table 2.** Stranding and bycatch records from Kep and Kampot provinces, between September  
276 2017 and May 2018. UPLOADED AS A SEPARATE DOCUMENT.

277

278