

Blagrove, Richard, Howe, Louis ORCID: <https://orcid.org/0000-0003-2001-2802> , Howatson, Glyn and Hayes, Philip (2020) Strength and conditioning for adolescent endurance runners. *Strength and Conditioning Journal*, 42 (1). pp. 2-11.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/5530/>

***Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.***

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

**provided that**

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
  - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

**You may not**

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing [insight@cumbria.ac.uk](mailto:insight@cumbria.ac.uk).

## Strength and Conditioning for Adolescent Endurance Runners

### ABSTRACT

For the adolescent athlete who chooses to specialize in endurance running, strength and conditioning (S&C) activities provide a means of enhancing several important determinants of performance and may reduce the risk of overuse injury. It is recommended that adolescent endurance runners include at least two S&C sessions per week that comprise of movement skills training, plyometric and sprint training, resistance training, plus exercises designed to target specific tissues that are vulnerable to injury. This article describes how these modalities of training can be integrated into the routine of adolescent endurance runners.

**Key words:** paediatric, endurance, distance running, youth, concurrent training

### INTRODUCTION

Endurance running is a popular choice of sport for young athletes. For example, in 2016-17, cross-country was the fourth and fifth largest sport by participation for boys and girls respectively in USA high schools (see [www.nfhs.org/ParticipationStatistics](http://www.nfhs.org/ParticipationStatistics)), and endurance running represented the second most popular sport (18.7%) in a survey ( $n=7794$ ) of Scandinavian 14 year olds (83). A young athlete should be exposed to a wide range of sports and physical activities during their adolescence, however the priority should be placed on the development of rudimentary motor skills and muscular strength (49). Endurance training during early-adolescence (11-14 years old) should form part of an active healthy lifestyle but should not take precedence over other modalities of sport-training (80). Endurance sports are typically associated with a high volume of training (80), which places the developing body of a young athlete under a high level of stress that could leave them susceptible to overtraining syndrome, illness and overuse injury (54). Therefore, specialization in endurance running should not occur until late-adolescence, when a young athlete's body is sufficiently mature and well-conditioned to cope with the rigours of this type of training. Strength and conditioning (S&C) activities might contribute towards lowering the risk of injury in athletes (45, 82), therefore providing sport-specific recommendations for this vulnerable population is important.

Endurance running is primarily limited by cardiovascular and metabolic factors, however there is an abundance of research showing that strength training (ST) activities (resistance training (RT), explosive resistance training (ERT) and plyometric training (PT)) can provide performance benefits to middle- (0.8-3 km) and long-distance (>3 km) runners (16, 29). A plethora of literature also exists that

demonstrates ST activities are also a safe and effective way of enhancing proxy measures of athletic performance in adolescents of both sexes (10, 41, 46). Specifically, compared to sport-only training, various forms of ST augment improvements in maximal strength, explosive strength, muscular endurance, sprint speed, agility test time, tennis serve velocity, kicking velocity, throwing velocity, and general motor skills (10, 40, 41, 46). However, there are currently no papers which have specifically summarized the effect of ST modalities on aerobic-related qualities in young athletes. For practitioners working with young distance runners in particular, it would be useful to establish whether ST activities offer any benefit to performance-related factors and how such training techniques could be applied in practice. Therefore, the aims of this article are to briefly review the literature that has investigated the efficacy of ST on the determinants of endurance running in adolescent runners, and provide guidelines for best practice to improve performance and minimize the occurrence of overuse injury.




Figure 1. Primary determinants of endurance running performance and the modalities of training to improve each.  $\dot{V}O_{2\max}$  = maximal oxygen uptake,  $s\dot{V}O_{2\max}$  = speed at maximal oxygen uptake.

## DETERMINANTS OF ENDURANCE RUNNING PERFORMANCE

Endurance running performance is determined by several key physiological variables, which are summarized in figure 1. The physiological determinants of performance for adolescents appears to be similar to those of adult runners (3, 27). A number of investigations have confirmed that maximal

oxygen uptake ( $\dot{V}O_{2max}$ ) is a significant predictor ( $r=0.5-0.9$ ) of performance for 1.5 km (1, 3), 3 km (1, 50, 87), 5 km (1, 27) and cross-country (24, 34) in young (10-18 years) groups of runners. The proportion of  $\dot{V}O_{2max}$  that can be sustained for a given duration (known as 'fractional utilization'), has also been shown to significantly correlate with endurance running performance in adolescents (50, 87). Running economy (RE), defined as the metabolic cost of running a given distance (79), is related to middle- (3, 87) and long-distance running performance (24, 34), and importantly, is influenced by neuromuscular related qualities, which can potentially be improved with ST activities (16, 29). Additionally, speed at  $\dot{V}O_{2max}$ , which is a product of  $\dot{V}O_{2max}$  and RE, correlates well with distance running performance in adolescents (1, 3, 24, 27).

The contribution of anaerobic factors to endurance running performance in adults is well-established (20), however the influence of anaerobic determinants on performance in young endurance runners has not been fully delineated. This is likely due to the unspecific nature of the tests (Wingate test, isokinetic strength tests, counter-movement jump height) utilized to quantify anaerobic and neuromuscular capacities in studies that have investigated young distance runners (3, 24, 28). Anaerobic capacity and neuromuscular capabilities are thought to play a large role in discriminating performance in runners who are closely matched from an aerobic perspective (22, 67). Mahon and co-authors (50) also showed that 55 m sprint and counter-movement jump were significant predictors of 3 km time trial in preadolescent children, although given the age of the participants, this finding could simply be a reflection of individuals possessing high or low levels of athletic ability across the range of the tests utilized. Speed at  $\dot{V}O_{2max}$  probably provides the most sport-specific representation of neuromuscular capabilities in distance runners, however measures of maximal running speed and anaerobic capacity are also important attributes (65). For an 800 m specialist in particular, near-maximal velocities of running are reached during the first 200 m of the race (74), which necessitates a high capacity of the neuromuscular and anaerobic system. Similarly, the quickest finisher at the end of a middle- or long-distance race often determines the winner (85), thus possessing a higher top speed is potentially crucial for achieving success in distance running. Regardless of the capacity for anaerobic and neuromuscular factors to predict endurance performance in adolescents, activities to develop sprint speed and muscular strength-qualities as part of a well-rounded physical training programme is recommended during adolescence irrespective of whether sport-specialization has occurred (47, 49).

## **EFFECT OF STRENGTH TRAINING ON AEROBIC-RELATED PARAMETERS**

Based upon the findings of recent reviews (16, 29), it is suggested that supplementing the training of an endurance runner with ST is likely to provide improvements in RE, time-trial (1.5 km – 10 km) performance and anaerobic parameters such as maximal sprint speed. Improvements in RE in the absence of changes in  $\dot{V}O_{2max}$ , speed at  $\dot{V}O_{2max}$ , blood lactate and body composition parameters suggests

that the underlying mechanisms predominantly relate to alterations in intra-muscular co-ordination and increases in stiffness (16). Specifically, ST brings about increases in motor unit recruitment, firing frequency and musculotendinous stiffness, which are thought to optimize the length-tension and force-velocity relationships of active skeletal muscle, thus reducing the metabolic cost of running (36). It is clear that the inclusion of ST also does not adversely affect  $\dot{V}O_{2max}$ , blood lactate markers or body composition (16). Concurrently, RE showed improvements of 2-8% with ST compared to a running-only control group following a 6-14 week intervention that includes 2-3 ST sessions per week (16).

### *Efficacy in Adolescent Runners*

Three studies have investigated the effect of ST specifically in young (<18 years) middle- or long-distance runners and these are summarized in table 1. A recent study by Blagrove et al. (17) found that two weekly sessions of ST (mainly PT and RT) added to the programme of post-pubertal adolescent distance runners (17 years) for ten weeks was 'possibly beneficial' for RE (effect size: 0.31-0.51) and 'highly likely beneficial' for maximal sprint speed. However, only the maximal speed improvement reached statistical significance ( $p<0.05$ ) compared to the change observed in the control group. Mikkola et al. (58) took a group of trained male and female distance runners (mean: 17 years,  $\dot{V}O_{2max}$ : 62.5 mL·kg<sup>-1</sup>·min<sup>-1</sup>) and following eight weeks of ERT, PT and sprint training, noted a difference (-2.7%) in RE at 14 km·h<sup>-1</sup> and improvements in anaerobic capabilities (speed during the maximal anaerobic running test and 30 m sprint) compared to a running-only group. It is noteworthy that both of these investigations (17, 58) included sprints (3-10 x 30-150 m) as part of the intervention, which provides a highly specific overload to the neuromuscular system in endurance runners. Interestingly, participants in the ST intervention groups in these studies reduced their weekly running volume to accommodate the additional S&C activities. Total time spent training was however very similar between intervention and control groups.

Bluett and associates (18) found that ten weeks of concurrent aerobic and ST provided little strength advantage and no change in 3 km time trial performance in 10-13 year old competitive runners compared to running only. This study utilized mainly single joint machine-based RT and did not measure any physiological parameters, which may explain the lack of effect observed. The authors speculated that excessive fatigue resulting from the concurrent training regimen may have compromised both strength and endurance adaptations (18). Interestingly, the blunting of strength adaptation which is often observed in adult performers when both strength- and endurance-training are included in the same training session (90) appears not to occur in children (53) and adolescents (75, 76). As the interference phenomenon is mediated by training volume and recovery from sessions (5), it seems likely that the volumes of each training modality included in the aforementioned studies were insufficient to negatively impact upon strength-related adaptation. Indeed, in elite youth soccer players (17 years) who

utilize higher workloads compared to younger performers, larger improvements were evident in strength and sprint performance after five weeks when ST was performed after sport-specific endurance training on two days per week, compared to a group who adopted a ST followed by endurance training sequence (33).

Accepted

Authors	Participants <i>n</i> (I/C), sex, age	Study duration	Running volume	ST frequency	ST prescription	Main strength outcomes	Main running-related outcomes
Blagrove et al. (18)	18 (9/9), both, 17.2 years	10	I: 151 min.wk <sup>-1</sup> C: 213 min.wk <sup>-1</sup>	2 per week	PT (3-4 x 6-8/15 m): box jumps, A-skips, hurdle jumps  RT (2-3 x 6-8 reps): back squat, RDL, rack pull, deadlift, step-ups, leg press, calf raise	MVC (15.4%, ES=0.86, p<0.05), vGRF <sub>jump</sub> (6.1%, ES=0.93)  CMJ, little difference compared to C	RE@ sLTP (3.2%, ES=0.31), sLTP -1 km.h <sup>-1</sup> (3.7%, ES=0.47), sLTP -2 km.h <sup>-1</sup> (3.6%, ES=0.51)  20 m sprint (3.6%, ES=0.32, p<0.05)
Bluett et al (19)	12 (6/6), both, 10-13 years	10	Both groups: 2 x runs per week (1x25-30 min continuous, 1xintervals)	2 per week	RT (3-4 sets x 10-12 reps): leg curl, leg extension, leg press, upper limb/trunk exercises	Increase (12.2-24%) loads lifted during ST but no change in peak torque pre-post I	3 km TT no change in either group
Mikkola et al (59)	25 (13/12), both, 17.3 years	8	I: 8.8 h.wk <sup>-1</sup> (19% running replaced with ST), C: 8.5 h.wk <sup>-1</sup>	3 per week	PT: alternative jumps, calf jumps, squat jumps, hurdle jumps  ERT (2-3 x 6-10 reps): half-squats, knee extension, knee flexion, leg press, calf raise	MVC (8%), 1RM (4%), RFD (31%) on leg press; all p<0.05.  CMJ and 5-bounds little difference compared to C	RE@ 14 km.h <sup>-1</sup> (2.7%, ES=0.32, p<0.05), @10,12,13 km.h <sup>-1</sup> , NS  BL@12 km.h <sup>-1</sup> (12%, p<0.05), @14 km.h <sup>-1</sup> (11%, p<0.05)  sSMART (3.0%, p<0.01), s30 m sprint (1.1%, p<0.01)

C=control, I=intervention, ST=strength training, wk=week, PT=plyometric training, RT=resistance training, RDL=Romanian deadlift, MVC=maximum voluntary contraction, ES=effect size, vGRF<sub>jump</sub>=vertical ground reaction force during squat jump, CMJ=counter-movement jump, RE=running economy, sLTP=speed at lactate turnpoint, TT=time trial, ERT=explosive resistance training, 1RM=one repetition maximum, RFD=rate of force development, NS=not statistically significant, BL=blood lactate

Table 1. Summary of studies (*n*=3) that have investigated the effects of a strength training intervention on adolescent endurance runners.

Research investigating the impact of ST techniques on performance-related measures in young athletes has tended to use participants from field-based sports, martial arts, court sports, aquatic sports, gymnastics and strength-based sports (40, 46). A number of studies using adolescent participants from other sports that require high-levels of aerobic fitness have observed superior improvements in Yo-Yo test (44, 51, 72, 91) and middle-distance time-trial performance (70, 73) after various modalities of ST were added (6-12 weeks) to a sports-specific training programme, compared to only practicing the sport.

Taken together, it appears that the addition of 2-3 ST sessions to the weekly routine of adolescent endurance runners provides a small but potentially meaningful benefit to RE and maximal sprint speed following an 8-10 week intervention. Evidence for improvements in performance exists for adult runners (16), however there is currently a lack of research in younger endurance runners. Benefits are likely to be larger for interventions of a longer duration (29) and for ST programmes that are supervised by qualified practitioners (16). Although the majority of studies in adults supplement a runners training with ST, there also appears to be no disadvantage to reducing weekly running volume to accommodate the addition of two weekly ST sessions.

## **PRACTICAL RECOMMENDATIONS**

### *Timing of Specialization and Long-Term Athlete Development*

Adolescence represents an important period of development in young athletes where significant alterations in hormonal status causes rapid physical growth (52). Contemporary views of long-term youth development suggest adolescents should avoid training routines that focus on intensive training in a single sport (for >8 months per year), or a total weekly training volume (in hours) that exceeds the athletes age in years, until late-adolescence (47, 49, 63). Evidence from several endurance sports shows elite senior athletes tend to specialize at a later age, and participate in a diverse range of sports during their earlier years (26, 60). Recent work has also shown that very few middle-distance runners ranked in the UK top 20 in the under-13 and under-15 age-groups experience success as senior runners (43). Young athletes who adopt an early-diversification, late-specialization approach to their development have fewer injuries, are at less risk of overtraining, and play sports longer than those who specialize in one sport before puberty (21, 30).

The youth athlete development model suggests a wide range of physical activities and training modalities should be utilized during adolescence, however movement skills training (MST) and development of strength qualities should be prioritized (47, 49). The emphasis on ST activities throughout an athletes development is thought to maximize adaptations to inter- and intra-muscular coordination, during a period when neuroplasticity is high (64). Improvements in muscular strength and motor control during this period have also been shown to improve physical performance (10, 46) and



lower the risk of sustaining an injury (62, 82). It is recommended that endurance training (and metabolic conditioning) is not emphasised, relative to other biomotor abilities, until late-adolescence (49), as typically this type of training is associated with high volumes of work, which may lead to injury or overtraining (54). Moreover, pre-pubertal children have tended to show smaller changes (<10%) in aerobic measures following endurance training interventions compared to post-pubertal adolescents and adults (54, 56). A recent study also showed that pre-pubertal boys (11 years) were metabolically comparable to well-trained endurance athletes and experienced less fatigue during high-intensity exercise compared to untrained adults (14). It was suggested that pre-pubertal children avoid specific training to develop aerobic metabolic qualities and shift priority during post-pubertal years once movement technique and mechanical competency have been developed (14). Due to the risks associated with early-specialization, it is recommended that adolescent athletes younger than 15 years old do not solely specialize in endurance running, but should participate in a wide range of sports and physical activities, including ST.

#### *Organization of the Training Microcycle*

Prior to specialization in a sport of a young athlete's choice, physical training should be semi-structured and not emphasize peaking for competitions (26, 63). Conversely, an adolescent endurance runner will typically run 45 to 55 miles weekly in preparation for a race (80), which when combined with academic and social commitments, can place a high level of physical and psychological stress on a young athlete (54). This necessitates a well-organized approach to training that caters to the needs of individual athletes and ensures sufficient periods of recovery between bouts of training.

Two seven-day microcycle designs are shown in Table 2a and 2b to illustrate how an adolescent endurance runner could incorporate S&C activities into their routine. Adolescent distance runners typically perform 2-3 high-intensity running sessions per week (15), and these should form the priority sessions in the programme (Table 2a and 2b; Tuesday, Thursday and Saturday). Similarly, a minimum of two ST sessions per week are suggested for adolescents (11, 48) and endurance runners (16). RT sessions should ideally take place at least three hours after a running session (6) and at least 24 hours recovery should follow after ST before an intensive running session (31). A novel approach to organizing S&C activities around training and lifestyle commitments with young runners is to incorporate shorter periods of activity ('training units') as part of running sessions wherever possible (see Table 2b). This type of programming is useful for young runners who perhaps cannot access a specialist S&C facility, and therefore perform a largely home-based routine, or are unable to commit to two full S&C sessions per week. Each training unit takes 10-20 min to complete, thus making it easy to integrate some purposeful S&C prior to or after running sessions. It is important to note that studies in

adolescent distance runners (17, 58) have shown that including weekly ST sessions are more effective than increasing running volume, at least in the short-term (8-10 weeks).

Assuming runners are of a non-strength trained status, it appears that a variety of ST modalities can be used to achieve similar outcomes. However to maximize long-term adaptations in young athletes, it is suggested that a periodized approach is adopted with fundamental skills training and RT prioritized initially (9, 25, 47). Figure 2 provides an overview of the session design and characteristics of specific training units recommended for adolescent distance runners. A similar session design framework has also been employed successfully in other investigations that used distance runners embarking upon a ST programme for the first time (9, 17, 58).




Figure 2. Recommended structure of a strength and conditioning session for adolescent endurance runners. Characteristics and example exercise prescription for individual training units are also shown. Prescription is sets x repetitions (unless stated).

<i>Day</i>	Mon	Tues	Wed	Thurs	Fri	Sat	Sun
<i>Session</i>	(am) Easy 30 min run (pm) S&C session	Interval training session	(am) Cycle or swim 30 min (pm) S&C session	Tempo run* (20-30 min)	Rest	Race or interval training session	Easy 45 min run

S&C = strength and conditioning. \* Continuous fast run performed at approximately 10 km race-pace (speed at lactate turn-point)

Table 2a. Example of a seven day microcycle for an adolescent endurance runner.

<i>Day</i>	Mon	Tues	Wed	Thurs	Fri	Sat	Sun
<i>Session</i>	Easy 30 min run + PT and RT	MST + Interval training session	Cycle or swim 30 min + RT and SC	PT + Tempo run (20-30 min)	Rest	Race or interval training session + SC	Easy 45 min run + MST

PT = plyometric training, RT = resistance training, MST = movement skills training, SC = specific conditioning

Table 2b. Example of a seven day microcycle for an adolescent endurance runner with strength and conditioning activities organised as training units before or after running sessions. Training units should last 10-20 min with the focus on movement quality rather than inducing high levels of fatigue.

### *Movement Skills Training*

The inclusion of MST in the routine of adolescent distance runners is recommended, and is likely to reduce long-term injury risk (45, 62, 82). This form of conditioning is ideal to include as part of a movement preparation warm-up routine prior to running and ST sessions, or as an independent training unit (82). MST should include activities to enhance both general (fundamental) and specific (running-related) movement skill and control, balance and dynamic stability, and low-level resistance training targeting specific muscle groups, such as the gluteals (37).

### *Plyometrics and Sprint Training*

Low intensity plyometric-based exercise that aim to develop ankle stiffness, such as skipping, low-box re-bound jumps, mini hurdle jumps and short range hopping tasks, offer a potent stimulus to the neuromuscular system and have independently been shown to enhance RE and time trial performance (12, 68, 71, 81, 86). It is suggested that 30-60 foot contacts per session are utilized initially with adolescent distance runners (17). Sprint training has also been used in several investigations showing enhancements in maximal speed and performance-related factors (17, 58, 59, 66). Three to five sets of short distance (30-60 m) technical and maximal sprints performed 2-3 times per week is likely to provide benefits to adolescent endurance runners.

### *Resistance Training*

Resistance training, which should include both ERT and heavy RT, increases motor unit recruitment and firing frequency, thus enhances a runners ability to appropriately control and express force during ground contact. Although changes in fat-free mass appear to be minimal following a ST intervention in distance runners (16), a targeted RT programme, that aims to increase muscle mass specifically around the proximal region of the lower limb may enhance biomechanical and physiological factors, which positively influence RE (36). Exercises, such as squats, deadlifts, step-ups and lunge patterns, which possess similar kinematic characteristics to running gait, are likely to provide the greatest transfer (8) and have been utilized in several previous investigations (9, 17, 58). Loaded jump squats, medicine ball throwing and weightlifting are examples of suitable ERT activities that can also be utilized (8, 9, 59). Upper limb exercises such as press-ups, rowing exercises and overhead presses, should also be incorporated to offset the vertical angular momentum created by the lower limbs and aid in controlling excessive rotation forces (42, 69, 77). One to three sets of each exercise performed in a moderate repetition range (8-12 repetitions) is likely to provide non-strength trained individuals with a stimulus sufficient to drive neuromuscular adaptation whilst developing skill in each exercise (9, 17, 59). Higher loads ( $\geq 80\%$  one repetition maximum) and lower repetition ranges (3-8 repetitions) are likely to be

required to provide further overload in strength-trained adolescents, with volume of work moderated via an increase in sets.

### *Specific Conditioning*

Many young endurance runners are motivated to include S&C activities to reduce injury risk more so than improve their performance (15). Youth endurance athletes have been identified as a high-risk group due to the rigorous training that they undertake during a critical period of their physical and emotional development (54, 80). Indeed, injury incidence rate has been reported to be higher in adolescent elite endurance runners compared to other endurance sports (88). Moreover, female adolescent runners tend to display higher rates of low bone mineral density and bone stress injuries compared to young female athletes in other sports (78). Overuse injuries occur over multiple running sessions when structure specific cumulative load exceeds capacity (13). MST, PT and resistance-based exercises are likely to contribute towards lowering risk of injury via enhancements in motor control and increased bone mineral density and tissue resilience (45, 62, 89). Exercises designed to expose specific muscles or tissues to a high magnitude of load are also likely to provide benefits to tendon stiffness (35) and tolerance to repetitive stress (7, 19, 61, 84, 89). It is recommended that such exercises are positioned in final part of a session or performed separately, as pre-fatiguing muscles in isolation is likely to be detrimental to performance in multi-joint tasks (4). Specifically for distance runners, targeted conditioning exercises should focus on the specific structures which are vulnerable to injury, or the muscles that contribute towards controlling the positioning of joints within the lower limb, such as: the intrinsic joints of the feet, the calf-Achilles complex, gluteal and hamstring muscles (2, 32, 38, 55, 57, 61). In addition, specific exercises that target proximal musculature around the lumbopelvic-hip complex ('core stability') are likely to offset the risk of several types of common overuse injuries in runners (23). Specifically, exercises which facilitate greater strength and control of the hip abductors and external rotators are likely to provide benefits (23, 39).

## **CONCLUSION**

Endurance running performance is constrained by several important physiological variables, however anaerobic and neuromuscular factors have also been recognized as being important. For the young athlete, participating in a broad range of sporting and physical pursuits is recommended during early-adolescence. Age-appropriate S&C should form an integral part of a well-rounded approach to the long-term physical development of all young sports performers. Participating in endurance running events can certainly form part of a programme of activities during adolescence, however it is suggested young athletes should not solely specialize in endurance running until late-adolescence. For the young

endurance runner, adding ST sessions twice per week that includes RT, PT and sprinting, is likely to provide benefits to RE and maximal sprint speed that translate to improved performance. Moreover, these activities, plus MST and specific strengthening of tissues vulnerable to injury, are important for lowering the risk of overuse injury.

Accepted

## REFERENCES

1. Abe D, Yanagawa K, Yamanobe K, and Tamura K. Assessment of middle-distance running performance in sub-elite young runners using energy cost of running. *Eur J Appl Physiol Occup Physiol* 77: 320-325, 1998.
2. Aderem J, and Louw QA. Biomechanical risk factors associated with iliotibial band syndrome in runners: a systematic review. *BMC Musculoskelet Disord* 16: 356, 2015.
3. Almarwaey OA, Jones AM, and Tolfrey K. Physiological correlates with endurance running performance in trained adolescents. *Med Sci Sports Exerc* 35: 480-487, 2003.
4. Augustsson J, ThomeÉ R, HÖrnstedt P, Lindblom J, Karlsson J, and Grimby G. Effect of pre-exhaustion exercise on lower-extremity muscle activation during a leg press exercise. *J Strength Cond Res* 17: 411-416, 2003.
5. Baar K. Training for endurance and strength: lessons from cell signaling. *Med Sci Sports Exerc* 38: 1939-1944, 2006.
6. Baar K. Using molecular biology to maximize concurrent training. *Sports Med* 44 Suppl 2: S117-125, 2014.
7. Baar K. Minimizing injury and maximizing return to play: lessons from engineered ligaments. *Sports Med* 47: 5-11, 2017.
8. Bazyley CD, Abbott HA, Bellon CR, Taber CB, and Stone MH. Strength training for endurance athletes: Theory to practice. *Strength Cond J* 37: 1-12, 2015.
9. Beattie K, Carson BP, Lyons M, Rossiter A, and Kenny IC. The effect of strength training on performance indicators in distance runners. *J Strength Cond Res* 31: 9-23, 2017.
10. Behringer M, Heede Av, Matthews M, and Mester J. Effects of strength training on motor performance skills in children and adolescents: a meta-analysis. *Pediatr Exerc Sci* 23: 186-206, 2011.
11. Bergeron MF, Mountjoy M, Armstrong N, Chia M, Côté J, Emery CA, Faigenbaum A, Hall G, Kriemler S, and Léglise M. International Olympic Committee consensus statement on youth athletic development. *Br J Sports Med* 49: 843-851, 2015.
12. Berryman N, Maurel DB, and Bosquet L. Effect of plyometric vs. dynamic weight training on the energy cost of running. *J Strength Cond Res* 24: 1818-1825, 2010.
13. Bertelsen M, Hulme A, Petersen J, Brund RK, Sørensen H, Finch C, Parner ET, and Nielsen R. A framework for the etiology of running-related injuries. *Scand J Med Sci Sports* 27: 1170-1180, 2017.
14. Birat A, Bourdier P, Piponnier E, Blazevich AJ, Maciejewski H, Duche P, and Ratel S. Metabolic and fatigue profiles are comparable between prepubertal children and well-trained adult endurance athletes. *Front Physiol* 9: 387, 2018.
15. Blagrove RC, Brown N, Howatson G, and Hayes PR. Strength and conditioning habits of competitive distance runners. *J Strength Cond Res*, 2017. [e-pub ahead of print]
16. Blagrove RC, Howatson G, and Hayes PR. Effects of strength training on the physiological determinants of middle-and long-distance running performance: a systematic review. *Sports Med* 48: 1117-1149, 2018.
17. Blagrove RC, Howe LP, Cushion EJ, Spence A, Howatson G, Pedlar CR, and Hayes PR. Effects of strength training on postpubertal adolescent distance runners. *Med Sci Sports Exerc*, 2018. [e-pub ahead of print]
18. Bluett KA, De Ste Croix MB, and Lloyd RS. A preliminary investigation into concurrent aerobic and resistance training in youth runners. *Isokinet Exerc Sci* 23: 77-85, 2015.
19. Bohm S, Mersmann F, and Arampatzis A. Human tendon adaptation in response to mechanical loading: a systematic review and meta-analysis of exercise intervention studies on healthy adults. *Sports Med Open* 1: 7, 2015.
20. Brandon LJ. Physiological factors associated with middle distance running performance. *Sports Med* 19: 268-277, 1995.
21. Brenner JS. Sports specialization and intensive training in young athletes. *Pediatr* 138: e20162148, 2016.

22. Bulbulian R, Wilcox AR, and Darabos BL. Anaerobic contribution to distance running performance of trained cross-country athletes. *Med Sci Sports Exerc* 18: 107-113, 1986.
23. Chuter VH, and de Jonge XAJ. Proximal and distal contributions to lower extremity injury: a review of the literature. *Gait Posture* 36: 7-15, 2012.
24. Cole AS, Woodruff ME, Horn MP, and Mahon AD. Strength, power, and aerobic exercise correlates of 5-km cross-country running performance in adolescent runners. *Pediatr Exerc Sci* 18: 374-384, 2006.
25. Cormie P, McGuigan MR, and Newton RU. Influence of strength on magnitude and mechanisms of adaptation to power training. *Med Sci Sports Exerc* 42: 1566-1581, 2010.
26. Côté J, Lidor R, and Hackfort D. ISSP position stand: To sample or to specialize? Seven postulates about youth sport activities that lead to continued participation and elite performance. *Int J Sport Exerc Psychol* 7: 7-17, 2009.
27. Cunningham LN. Relationship of running economy, ventilatory threshold, and maximal oxygen consumption to running performance in high school females. *Res Q Exerc Sport* 61: 369-374, 1990.
28. Dellagrana RA, Guglielmo LG, Santos BV, Hernandez SG, da Silva SG, and de Campos W. Physiological, anthropometric, strength, and muscle power characteristics correlates with running performance in young runners. *J Strength Cond Res* 29: 1584-1591, 2015.
29. Denadai BS, de Aguiar RA, de Lima LC, Greco CC, and Caputo F. Explosive training and heavy weight training are effective for improving running economy in endurance athletes: A systematic review and meta-analysis. *Sports Med* 47: 545-554, 2017.
30. DiFiori JP, Benjamin HJ, Brenner JS, Gregory A, Jayanthi N, Landry GL, and Luke A. Overuse injuries and burnout in youth sports: a position statement from the American Medical Society for Sports Medicine. *Br J Sports Med* 48: 287-288, 2014.
31. Doma K, Deakin GB, and Bentley DJ. Implications of impaired endurance performance following single bouts of resistance training: an alternate concurrent training perspective. *Sports Med* 47: 2187-2200, 2017.
32. Duffey MJ, Martin DF, Cannon DW, Craven T, and Messier SP. Etiologic factors associated with anterior knee pain in distance runners. *Med Sci Sports Exerc* 32: 1825-1832, 2000.
33. Enright K, Morton J, Iga J, and Drust B. The effect of concurrent training organisation in youth elite soccer players. *Eur J Appl Physiol* 115: 2367-2381, 2015.
34. Fernhall B, Kohrt W, Burkett LN, and Walters S. Relationship between the lactate threshold and cross-country run performance in high school male and female runners. *Pediatr Exerc Sci* 8: 37-47, 1996.
35. Fletcher JR, Esau SP, and MacIntosh BR. Changes in tendon stiffness and running economy in highly trained distance runners. *Eur J Appl Physiol* 110: 1037-1046, 2010.
36. Fletcher JR, and MacIntosh BR. Running economy from a muscle energetics perspective. *Front Physiol* 8: 433, 2017.
37. Fort-Vanmeerhaeghe A, Romero-Rodriguez D, Lloyd RS, Kushner A, and Myer GD. Integrative neuromuscular training in youth athletes. Part II: Strategies to prevent injuries and improve performance. *Strength Cond J* 38: 9-27, 2016.
38. Franettovich MS, Honeywill C, Wyndow N, Crossley KM, and Creaby MW. Neuromotor control of gluteal muscles in runners with achilles tendinopathy. *Med Sci Sports Exerc* 46: 594-599, 2014.
39. Frederic M, Cookingham CL, Chaudhari AM, Dowdell BC, Oestreicher N, and Sahrman SA. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin J Sport Med* 10: 169-175, 2000.
40. Granacher U, Lesinski M, Büsch D, Muehlbauer T, Prieske O, Puta C, Gollhofer A, and Behm DG. Effects of resistance training in youth athletes on muscular fitness and athletic performance: a conceptual model for long-term athlete development. *Front Physiol* 7: 164, 2016.
41. Harries SK, Lubans DR, and Callister R. Resistance training to improve power and sports performance in adolescent athletes: A systematic review and meta-analysis. *J Sci Med Sport* 15: 532-540, 2012.



42. Johnston RE, Quinn TJ, Kertzer R, and Vroman NB. Strength training in female distance runners: impact on running economy. *J Strength Cond Res* 11: 224-229, 1997.
43. Kearney PE, and Hayes PR. Excelling at youth level in competitive track and field athletics is not a prerequisite for later success. *J Sports Sci*, 2018. [e-pub ahead of print]
44. Klusemann MJ, Pyne DB, Fay TS, and Drinkwater EJ. Online video-based resistance training improves the physical capacity of junior basketball athletes. *J Strength Cond Res* 26: 2677-2684, 2012.
45. Lauersen JB, Bertelsen DM, and Andersen LB. The effectiveness of exercise interventions to prevent sports injuries: a systematic review and meta-analysis of randomised controlled trials. *Br J Sports Med* 48: 871-877, 2014.
46. Lesinski M, Prieske O, and Granacher U. Effects and dose-response relationships of resistance training on physical performance in youth athletes: a systematic review and meta-analysis. *Br J Sports Med* 50: 781-795, 2016.
47. Lloyd RS, Cronin JB, Faigenbaum AD, Haff GG, Howard R, Kraemer WJ, Micheli LJ, Myer GD, and Oliver JL. National Strength and Conditioning Association position statement on long-term athletic development. *J Strength Cond Res* 30: 1491-1509, 2016.
48. Lloyd RS, Faigenbaum AD, Stone MH, Oliver JL, Jeffreys I, Moody JA, Brewer C, Pierce KC, McCambridge TM, and Howard R. Position statement on youth resistance training: the 2014 International Consensus. *Br J Sports Med* 48: 498-505, 2014.
49. Lloyd RS, and Oliver JL. The youth physical development model: A new approach to long-term athletic development. *Strength Cond J* 34: 61-72, 2012.
50. Mahon A, Del Corral P, Howe C, Duncan G, and Ray M. Physiological correlates of 3-kilometer running performance in male children. *Int J Sports Med* 17: 580-584, 1996.
51. Makhlof I, Castagna C, Manzi V, Laurencelle L, Behm DG, and Chaouachi A. Effect of sequencing strength and endurance training in young male soccer players. *J Strength Cond Res* 30: 841-850, 2016.
52. Malina RM. Physical growth and biological maturation of young athletes. *Exerc Sport Sci Rev* 22: 280-284, 1994.
53. Marta C, Marinho D, Barbosa T, Izquierdo M, and Marques M. Effects of concurrent training on explosive strength and VO<sub>2</sub>max in prepubescent children. *Int J Sports Med* 34: 888-896, 2013.
54. Matos N, and Winsley RJ. Trainability of young athletes and overtraining. *J Sports Sci Med* 6: 353, 2007.
55. McKeon PO, and Fouchet F. Freeing the foot: integrating the foot core system into rehabilitation for lower extremity injuries. *Clin Sports Med* 34: 347-361, 2015.
56. McNarry M, and Jones A. The influence of training status on the aerobic and anaerobic responses to exercise in children: A review. *Eur J Sport Sci* 14: S57-S68, 2014.
57. Messier SP, Edwards DG, Martin DF, Lowery RB, Cannon DW, James MK, Curl WW, Read HM, Jr., and Hunter DM. Etiology of iliotibial band friction syndrome in distance runners. *Med Sci Sports Exerc* 27: 951-960, 1995.
58. Mikkola J, Rusko H, Nummela A, Pollari T, and Hakkinen K. Concurrent endurance and explosive type strength training improves neuromuscular and anaerobic characteristics in young distance runners. *Int J Sports Med* 28: 602-611, 2007.
59. Millet GP, Jaouen B, Borrani F, and Candau R. Effects of concurrent endurance and strength training on running economy and VO<sub>2</sub> kinetics. *Med Sci Sports Exerc* 34: 1351-1359, 2002.
60. Moesch K, Elbe AM, Hauge ML, and Wikman JM. Late specialization: the key to success in centimeters, grams, or seconds (cgs) sports. *Scand J Med Sci Sports* 21: e282-e290, 2011.
61. Mucha MD, Caldwell W, Schlueter EL, Walters C, and Hassen A. Hip abductor strength and lower extremity running related injury in distance runners: A systematic review. *J Sci Med Sport* 20: 349-355, 2017.
62. Myer GD, Faigenbaum AD, Chu DA, Falkel J, Ford KR, Best TM, and Hewett TE. Integrative training for children and adolescents: techniques and practices for reducing sports-related injuries and enhancing athletic performance. *Phys Sports Med* 39: 74-84, 2011.

63. Myer GD, Jayanthi N, DiFiori JP, Faigenbaum AD, Kiefer AW, Logerstedt D, and Micheli LJ. Sports specialization, part II: alternative solutions to early sport specialization in youth athletes. *Sports Health* 8: 65-73, 2016.
64. Myer GD, Kushner AM, Faigenbaum AD, Kiefer A, Kashikar-Zuck S, and Clark JF. Training the developing brain, part I: cognitive developmental considerations for training youth. *Curr Sports Med Rep* 12: 304-310, 2013.
65. Noakes TD. Implications of exercise testing for prediction of athletic performance: a contemporary perspective. *Med Sci Sports Exerc* 20: 319-330, 1988.
66. Paavolainen L, Hakkinen K, Hamalainen I, Nummela A, and Rusko H. Explosive-strength training improves 5-km running time by improving running economy and muscle power. *J Appl Physiol* 86: 1527-1533, 1999.
67. Paavolainen LM, Nummela AT, and Rusko HK. Neuromuscular characteristics and muscle power as determinants of 5-km running performance. *Med Sci Sports Exerc* 31: 124-130, 1999.
68. Pellegrino J, Ruby BC, and Dumke CL. Effect of plyometrics on the energy cost of running and MHC and titin isoforms. *Med Sci Sports Exerc* 48: 49-56, 2016.
69. Piacentini MF, De Ioannon G, Comotto S, Spedicato A, Vernillo G, and La Torre A. Concurrent strength and endurance training effects on running economy in master endurance runners. *J Strength Cond Res* 27: 2295-2303, 2013.
70. Potdevin FJ, Alberty ME, Chevutschi A, Pelayo P, and Sidney MC. Effects of a 6-week plyometric training program on performances in pubescent swimmers. *J Strength Cond Res* 25: 80-86, 2011.
71. Ramirez-Campillo R, Alvarez C, Henriquez-Olguin C, Baez EB, Martinez C, Andrade DC, and Izquierdo M. Effects of plyometric training on endurance and explosive strength performance in competitive middle- and long-distance runners. *J Strength Cond Res* 28: 97-104, 2014.
72. Ramírez-Campillo R, Henríquez-Olguín C, Burgos C, Andrade DC, Zapata D, Martínez C, Álvarez C, Baez EI, Castro-Sepúlveda M, and Peñailillo L. Effect of progressive volume-based overload during plyometric training on explosive and endurance performance in young soccer players. *J Strength Cond Res* 29: 1884-1893, 2015.
73. Ramírez-Campillo R, Meylan C, Alvarez C, Henríquez-Olguín C, Martínez C, Cañas-Jamett R, Andrade DC, and Izquierdo M. Effects of in-season low-volume high-intensity plyometric training on explosive actions and endurance of young soccer players. *J Strength Cond Res* 28: 1335-1342, 2014.
74. Reardon J. Optimal pacing for running 400-and 800-m track races. *Am J Phys* 81: 428-435, 2013.
75. Santos A, Marinho D, Costa A, Izquierdo M, and Marques M. The effects of concurrent resistance and endurance training follow a specific detraining cycle in young school girls. *J Hum kinet* 29: 93-103, 2011.
76. Santos AP, Marinho DA, Costa AM, Izquierdo M, and Marques MC. The effects of concurrent resistance and endurance training follow a detraining period in elementary school students. *J Strength Cond Res* 26: 1708-1716, 2012.
77. Schumann M, Mykkanen OP, Doma K, Mazzolari R, Nyman K, and Hakkinen K. Effects of endurance training only versus same-session combined endurance and strength training on physical performance and serum hormone concentrations in recreational endurance runners. *Appl Physiol Nutr Metabol* 40: 28-36, 2015.
78. Scofield KL, and Hecht S. Bone health in endurance athletes: runners, cyclists, and swimmers. *Curr Sports Med Rep* 11: 328-334, 2012.
79. Shaw AJ, Ingham SA, and Folland JP. The valid measurement of running economy in runners. *Med Sci Sports Exerc* 46: 1968-1973, 2014.
80. Solomon ML, Briskin SM, Sabatina N, and Steinhoff JE. The pediatric endurance athlete. *Curr Sports Med Rep* 16: 428-434, 2017.
81. Spurrs RW, Murphy AJ, and Watsford ML. The effect of plyometric training on distance running performance. *Eur J Appl Physiol* 89: 1-7, 2003.

82. Steib S, Rahlf AL, Pfeifer K, and Zech A. Dose-response relationship of neuromuscular training for injury prevention in youth athletes: a meta-analysis. *Front Physiol* 8: 920, 2017.
83. Tammelin T, Näyhä S, Hills AP, and Järvelin M-R. Adolescent participation in sports and adult physical activity. *Am J Prev Med* 24: 22-28, 2003.
84. Tenforde AS, Kraus E, and Fredericson M. Bone stress injuries in runners. *Phys Med Rehabil Clin N Am* 27: 139-149, 2016.
85. Thiel C, Foster C, Banzer W, and De Koning J. Pacing in Olympic track races: competitive tactics versus best performance strategy. *J Sports Sci* 30: 1107-1115, 2012.
86. Turner AM, Owings M, and Schwane JA. Improvement in running economy after 6 weeks of plyometric training. *J Strength Cond Res* 17: 60-67, 2003.
87. Unnithan V, Timmons J, Paton J, and Rowland T. Physiologic correlates to running performance in pre-pubertal distance runners. *Int J Sports Med* 16: 528-533, 1995.
88. von Rosen P, Floström F, Frohm A, and Heijne A. Injury patterns in adolescent elite endurance athletes participating in running, orienteering, and cross-country skiing. *Int J Sports Phys Ther* 12: 822, 2017.
89. Warden SJ, Davis IS, and Fredericson M. Management and prevention of bone stress injuries in long-distance runners. *J Orthop Sports Phys Ther* 44: 749-765, 2014.
90. Wilson JM, Marin PJ, Rhea MR, Wilson SM, Loenneke JP, and Anderson JC. Concurrent training: a meta-analysis examining interference of aerobic and resistance exercises. *J Strength Cond Res* 26: 2293-2307, 2012.
91. Wong P-I, Chamari K, and Wisløff U. Effects of 12-week on-field combined strength and power training on physical performance among U-14 young soccer players. *J Strength Cond Res* 24: 644-652, 2010.